

## **STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid**

### **Rite Aid Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredient (in each 30 mL)**

Bismuth subsalicylate 525 mg

#### **Purpose**

Upset stomach reliever/Antidiarrheal

#### **Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - fullness
  - belching

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

#### **Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### **Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

**When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) right away.

**Directions**

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL (1 dose) every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
  - do not exceed 8 doses (240 mL) in 24 hours
  - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each 30 mL contains:** potassium 8 mg sodium 9 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium

- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

\*Compare to the active ingredient in Pepto-Bismol®

REGULAR STRENGTH

STOMACH RELIEF

BISMUTH SUBSALICYLATE 525 mg

UPSET STOMACH RELIEVER

ANTIDIARRHEAL

relieves:

heartburn • indigestion • nausea

upset stomach • diarrhea

alcohol free • sugar free

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company, Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

### **Package Label**

**Drug Facts** (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ mL = milliliter ■ shake well before using  
 ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours  
 ■ use until diarrhea stops but not more than 2 days ■ children under 12 years of age: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: potassium 8 mg, sodium 9 mg ■ each 30 mL contains: salicylate 206 mg  
 ■ low sodium ■ keep tightly closed ■ protect from freezing  
 ■ avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

**Questions or comments?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**DISTRIBUTED BY:** RITE AID 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com PLD-C4928 LB007773

**SATISFACTION GUARANTEE:**  
 If you're not satisfied, we'll happily refund your money.



NDC 11822-2910-8

**FREE** ALCOHOL FREE  
 F R M SUGAR FREE

Compare to the active ingredient in **Pepto-Bismol®**\*

# REGULAR STRENGTH STOMACH RELIEF

**BISMUTH SUBSALICYLATE 525 mg**

**UPSET STOMACH RELIEVER  
 ANTIDIARRHEAL**

relieves:  
 heartburn • indigestion • nausea  
 upset stomach • diarrhea



8 FL OZ (237 mL)

**Drug Facts**

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

**Active ingredient (in each 30 mL dose)** Purpose  
 Bismuth subsalicylate 525 mg..... Upset stomach reliever/Antidiarrheal

**Uses** relieves ■ diarrhea ■ travelers' diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ gas ■ indigestion ■ nausea ■ fullness ■ belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert: Contains salicylate. Do not take if you are:** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use** if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use** if you have  
 ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use** if you are taking any drug for ■ diabetes ■ gout ■ arthritis  
 ■ anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

## RITE AID Regular Strength Stomach Relief

## STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-2910
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-2910-8	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/27/2019	
2	NDC:11822-2910-7	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/27/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	12/27/2019	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 2/2023

Rite Aid Corporation