

**COLD FLU SEVERE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated**  
**HEB**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**HEB - 1163 - 2019-1004**

**Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- for the temporary relief of the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not exceed recommended dosage**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-

800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 4 hours</li><li>▪ swallow whole - do not crush, chew, or dissolve</li><li>▪ do not take more than 10 caplets in 24 hours</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>▪ ask a doctor</li></ul>

### Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

### Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

### Questions or comments?

1-844-705-4384

### PRINCIPAL DISPLAY PANEL

Compare to Tylenol Cold + Flu Severe Caplets active ingredients

NDC 37808-663-05

HEB

Cold & Flu Severe

Acetaminophen / Pain Reliever/Fever Reducer

Dextromethorphan HBr / Cough Suppressant

Guaifenesin / Expectorant

Phenylephrine HCl / Nasal Decongestant

Cold & Flu

COOL TASTE

For Adults

Actual Size

Relief of:

Head & Body Aches, Fever, Sore Throat, Cough

Nasal Congestion, Mucus & Chest Congestion

24 CAPLETS

**Drug Facts (continued)**

**Other information**

- store in 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

**Inactive ingredients** acetylsalicylic acid, croscarmellose sodium, croscopolone, D&C yellow #10

**Questions or comments?** 1-844-705-4394

**Drug Facts (continued)**

aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, methylcellulose, polyethylene glycol, polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

**Drug Facts (continued)**

**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions), or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**Directions**

do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 4 hours
- swallow whole - do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours

children under 12 years

- ask a doctor

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Keep out of reach of children.**

**If pregnant or breast-feeding,** ask a health professional before use. These could be signs of a serious condition.

- cough comes back or occurs with rash or headache that lasts
- new symptoms occur
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- part, nasal congestion, or cough gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur

**Stop use and ask a doctor if**

- when using this product do not exceed recommended dosage
- if you are taking the blood thinning drug warfarin

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Drug Facts (continued)**

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- part, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts
- new symptoms occur

**Drug Facts (continued)**

**Directions**

do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 4 hours
- swallow whole - do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours

children under 12 years

- ask a doctor

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Keep out of reach of children.**

**If pregnant or breast-feeding,** ask a health professional before use. These could be signs of a serious condition.

- cough comes back or occurs with rash or headache that lasts
- new symptoms occur
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- part, nasal congestion, or cough gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur

**Stop use and ask a doctor if**

- when using this product do not exceed recommended dosage
- if you are taking the blood thinning drug warfarin

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Drug Facts (continued)**

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- part, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts
- new symptoms occur



**DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN**



DISTRIBUTED BY H-E-B®  
SAN ANTONIO, TX 78204

**Drug Facts (continued)**

**Active ingredients (in each caplet) Purpose**

- Acetaminophen 325 mg: Pain reliever/fever reducer
- Dextromethorphan HBr 10 mg: Cough suppressant
- Guaifenesin 200 mg: Expectorant
- Phenylephrine HCl 5 mg: Nasal decongestant

**Uses**

- for the temporary relief of the following cold/flu symptoms: minor aches and pains, headache, sore throat
- nasal congestion
- cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

**Drug Facts (continued)**

Compare to Tylenol® Cold + Flu Severe Caplets active ingredients\*

NDC 37808-663-05

**H-E-B**

# Cold & Flu SEVERE

Acetaminophen / Pain Reliever/Fever Reducer  
Dextromethorphan HBr / Cough Suppressant  
Guaifenesin / Expectorant  
Phenylephrine HCl / Nasal Decongestant

**Cold & Flu**

For Adults

**Relief of:**

- Head & Body Aches • Fever • Sore Throat • Cough
- Nasal Congestion • Mucus & Chest Congestion



**24 CAPLETS**

NC

NC

NC

# COLD FLU SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-663
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, PREGELATINIZED CORN</b> (UNII: O8232NY3SJ)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	AAA;1136
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-663-05	2 in 1 CARTON	05/01/2018	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/01/2018	

**Labeler - HEB (007924756)**

Revised: 12/2022

HEB