

**EQUATE ASPIRIN- aspirin tablet, chewable**  
**Wal-Mart Stores Inc**

-----  
**Wal-Mart Chewable Aspirin Drug Facts**

**Active ingredient (in each chewable tablet)**

Aspirin 81 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

for the temporary relief of minor aches and pains

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

## **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

## **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

## **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
  - you experience any of the following signs of stomach bleeding:
    - feel faint
    - vomit blood
    - have bloody or black stools
    - have stomach pain that does not get better
    - pain gets worse or lasts for more than 10 days
    - fever gets worse or lasts for more than 3 days
    - redness or swelling is present
    - new symptoms occur
    - ringing in the ears or loss of hearing occurs
- These could be signs of a serious condition

## **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

## **Other information**

- store at 20-25°C (68-77°F)

### **Inactive ingredients**

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

### **Questions or comments?**

**1-888-287-1915**

### **Principal Display Panel**

3 PACK

equate™

Compare to Bayer® Chewable Aspirin active ingredient

Chewable Aspirin

Pain Reliever (NSAID)

LOW DOSE

Aspirin Regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

Orange Flavor


Actual Size

3 Bottles Inside

81 mg

108 TABLETS

ACTUAL SIZE



**3 PACK**

**equate™**

NDC 49035-467-08



# Chewable Aspirin

Pain Reliever (NSAID)

**LOW DOSE**

Aspirin Regimen  
Talk to your doctor or other healthcare provider before using this product for your heart.



Orange Flavor



Actual Size



3 Bottles Inside

**81 mg**

**108 TABLETS**

A2002 ZE C13

**Drug Facts**

**Active ingredient (in each chewable tablet)** ..... **Purpose**  
 Aspirin 81 mg (NSAID)\* ..... Pain reliever  
 \*nonsteroidal anti-inflammatory drug

**Uses** for the temporary relief of minor aches and pains

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older ■ have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after to surgery unless directed by a doctor

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING



Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

**DISTRIBUTED BY:**  
**Walmart Inc.,**  
 Bentonville, AR 72716  
 \*\*This product is not manufactured or distributed by Bayer Aktiengesellschaft, owner of the registered trademark Bayer®.



**Drug Facts (continued)**

- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor


**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

**Questions or comments?** 1-888-287-1915



**EQUATE ASPIRIN**

aspirin tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-467
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

**Product Characteristics**

Color	ORANGE	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	L467
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-467-68	1 in 1 CARTON	03/19/2003	
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:49035-467-08	3 in 1 CARTON	03/19/2003	
2		36 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/19/2003	

**Labeler** - Wal-Mart Stores Inc (051957769)

Revised: 9/2024

Wal-Mart Stores Inc