# ASPIRIN- aspirin tablet, film coated CHAIN DRUG MARKETING ASSOCIATION INC

-----

# **Quality Choice 44-157**

# Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)\*
\*nonsteroidal anti-inflammatory drug

## **Purpose**

Pain reliever/fever reducer

#### Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

# Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

# Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

## Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

# If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

#### Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

## Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

#### Questions or comments?

1-800-426-9391

# **Principal Display Panel**

NDC 63868-352-10

QC® QUALITY CHOICE

†Compare to the active ingredient in Genuine BAYER® Aspirin

Aspirin (NSAID)

Pain Reliever Fever Reducer 325 mg

**Coated Tablets** 

100 Tablets

actual size

# TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.

50844 REV0122C15712

Distributed by CDMA, Inc. 43157 W Nine Mile Novi, MI 48375 www.qualitychoice.com Questions: 800-935-2362



# Adhesive Only

*luestions or comments?* 1-800-426-9391

STOP PEELING

Stop use and ask a doctor if an allergic reaction occurs. polyethylene glycol, propylene glycol ■ store at 25°C (77°F); excursions permitted between 15°-30°C If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or ■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days Seek medical help right away.

you experience any of the following signs of stomach bleeding: 

feel faint 
vomit blood Ask a doctor or pharmacist before use if you are taking a nrescrintion drug for gout diabetes arthritis Other information ■children under 12 years: do not use unless directed by a adults and children 12 years and over take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 drink a full (eep out of reach of children. In case of overdose, get medical doctor because it may cause problems in the unborn child or ater in pregnancy unless definitely directed to do so by a ıelp or contact a Poison Control Center right away complications during hese could be signs of a serious condition. redness or swelling is present new symptoms occur (59°-86°F) ■ use by expiration date on package tablets in 24 hours unless directed by a doctor have stomach pain that does not get better have bloody or black stools glass of water with each dose do not take more than directed delivery.

■ stomach bleeding warning applies to you■ you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis,

or kidney disease ■ you have asthma

Ask a doctor before use if

■ you are taking a diuretic

reliever/tever reducer if you have ever had a any of its ingredients Do not use ■ if you are allergic to aspirin or any other pain

if you have ever had an allergic reaction to this product or

Drug Facts (continued

#### **Quality Choice 44-157**

#### **ASPIRIN**

aspirin tablet, film coated

<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-352
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76F) (ASPIRIN - UNII:R16CO5Y76F)	AS PIRIN	325 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: 08232NY3SJ)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;157;ASPIRIN	
Contains				

F	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-352- 10	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
2	NDC:63868-352- 03	300 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	02/10/1996		

# Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		038154464	pack(63868-352)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		832867837	manufacture(63868-352)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(63868-352)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63868-352)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>

LNK International, Inc. 967626305 pack(63868-352)

Revised: 11/2024 CHAIN DRUG MARKETING ASSOCIATION INC