

PRESSURE AND PAIN PE- acetaminophen and phenylephrine hydrochloride tablet, coated
Chain Drug Marketing Association

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

QCH - 1120 - 2019-1004

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - nasal congestion
- promotes sinus drainage
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 4 hours▪ do not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #6, hypromellose, lactose, magnesium stearate, povidone, pregelatinized starch, propylene glycol, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

PRINCIPAL DISPLAY PANEL

NDC 63868-971-24

QUALITY CHOICE

†Compare to Active Ingredients in SUDAFED PE® Pressure + Pain

Pressure + Pain PE

Pain Reliever / Fever Reducer, Nasal Decongestant

Acetaminophen, Phenylephrine HCl

For Relief of:

Sinus Headache

Sinus Pressure & Congestion

24 Caplets

SEE NEW WARNINGS INFORMATION & DIRECTIONS

N C

N C

N C

Drug Facts (continued)

Active ingredients (in each caplet) Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
- sinus congestion and pressure
- minor aches and pains
- nasal congestion
- temporarily reduces fever

Warnings

Liver warnings: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

Drug Facts

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children

- take 2 caplets every 4 hours
- do not take more than 10 caplets in 24 hours

children under 12 years

- ask a doctor

QC QUALITY CHOICE

Maximum Strength, Non-Drowsy

Pressure & Pain PE

Pain Reliever | Fever Reducer, Nasal Decongestant

6

35515195673

1

QC QUALITY CHOICE

NDC 63868-971-24

Compare to Active Ingredients in SUDAFED PE® Pressure + Pain

Maximum Strength, Non-Drowsy

Pressure + Pain PE

Pain Reliever | Fever Reducer, Nasal Decongestant

Acetaminophen, Phenylephrine HCl

For Relief of:

- Sinus Headache
- Sinus Pressure & Congestion

24 Caplets

SEE NEW WARNINGS INFORMATION & DIRECTIONS

Distributed by C.D.M.A., Inc.®
 131577 Anne Arundel
 Nov, MD 21113
 www.qualitychoice.com
 Questions: 248-449-9300

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

F112006QCH_R1

Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #6, hypromellose, lactose,

Drug Facts (continued)

magnesium stearate, povidone, pregelatinized starch, propylene glycol, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

acetaminophen and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-971
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1114
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-971-24	2 in 1 CARTON	08/29/2007	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/29/2007	

Labeler - Chain Drug Marketing Association (011920774)

Revised: 10/2019

Chain Drug Marketing Association