

DULCOLAX- bisacodyl tablet, coated
Mechanical Servants, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dulcolax®

Drug Facts

Active ingredient (in each tablet)

Bisacodyl USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

take with a glass of water

adults and children 12 years and over	1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	1 tablet in a single daily dose

children under 6 years of age | ask a doctor

Other information

- store at 20-25°C (68-77°F)
- protect from excessive humidity

Inactive ingredients

acacia, acetylated monoglyceride, carnauba wax, cellulose acetate phthalate, corn starch, dibutyl phthalate, docusate sodium, gelatin, glycerin, iron oxides, kaolin, lactose, magnesium stearate, methylparaben, pharmaceutical glaze, polyethylene glycol, povidone, propylparaben, Red No. 30 lake, sodium benzoate, sorbitan monooleate, sucrose, talc, titanium dioxide, white wax, Yellow No. 10 lake

Questions

1-888-285-9159 (English/Spanish)

Or visit **www.DULCOLAX.com**

Repackaged and distributed by Convenience Valet® Melrose Park, IL 60160

PRINCIPAL DISPLAY PANEL - 5 mg Pouch Package

CONVENIENCE

CV

VALET®

DRINKING CUP

INCLUDED

***Dulcolax*®**

LAXATIVE

(bisacodyl USP 5 mg)

C6A

3 TABLETS

CONVENIENCE VALET CV

DRINKING CUP INCLUDED


Dulcolax

LAXATIVE
(bisacodyl USP 5 mg)

C6A

3 TABLETS

Dulcolax



0 9 4 1 8 7 0 6 7 4 1 3

Dulcolax

LAXATIVE
(bisacodyl USP 5 mg)

Drug Facts

Active ingredient (in each tablet)	Purpose
Bisacodyl USP 5 mg	Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

6741-09-08.FB Repackaged and distributed by Convenience Valet®
Mekrose Park, IL 60160 C6A-CVDF-6741

CB-DS-E-6741 ▼ LIFT PANEL UP ▼

Dulcolax

LAXATIVE TABLETS
(bisacodyl USP, 5 mg)

Drug Facts

Active ingredient (in each tablet)	Purpose
Bisacodyl USP 5 mg	Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

6741-09-08.FB Repackaged and distributed by Convenience Valet®
Mekrose Park, IL 60160 C6A-CVDF-6741

CB-DS-F-6741 ▼ LIFT PANEL UP ▼

Drug Facts (continued)

When using this product, do not chew or crush tablet(s)

when using this product • do not chew or crush tablet(s)
 • do not use within 1 hour after taking an antacid or milk
 • it may cause stomach discomfort, faintness and cramps
Stop use and ask a doctor if • you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
 • you need to use a laxative for more than 1 week
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions take with a glass of water

adults and children 12 years and over	1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Hinge

Hinge

Drug Facts (continued)

Other information

- store at 20-25°C (68-77°F)
- protect from excessive humidity

Inactive ingredients acacia, acetylated monoglyceride, carnauba wax, cellulose acetate phthalate, corn starch, dibutyl phthalate, docusate sodium, gelatin, glycerin, iron oxides, kaolin, lactose, magnesium stearate, methylparaben, pharmaceutical glaze, polyethylene glycol, povidone, propylparaben, Red No. 30 lake, sodium benzoate, sorbitan monooleate, sucrose, talc, titanium dioxide, white wax, Yellow No. 10 lake

Questions 1-888-285-9159 (English/Spanish)
 Or visit www.DULCOLAX.com

Individually sealed. Do not use if seal is broken.
 DULCOLAX does not supply store brands.
 Boehringer Ingelheim Consumer Health Care Products Division of Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877. Made in Mexico.
 Copyright © 2008, Boehringer Ingelheim Pharmaceuticals, Inc.
 All rights reserved.

DULCOLAX

bisacodyl tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:29485-6741(NDC:0597-0012)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
bisacodyl (UNII: 10X0709Y6I) (bisacodyl - UNII:10X0709Y6I)	bisacodyl	5 mg

Inactive Ingredients

Ingredient Name	Strength
acacia (UNII: 5C5403N26O)	
carnauba wax (UNII: R12CBM0EIZ)	
cellulose (UNII: F2O5O2O19F)	
starch, corn (UNII: O8232NY3SJ)	
dibutyl phthalate (UNII: 2286E5R2KE)	
docosate sodium (UNII: F05Q2T2JA0)	
gelatin (UNII: 2G86QN327L)	
glycerin (UNII: PDC6A3C0OX)	
kaolin (UNII: 24H4NWX5CO)	
lactose (UNII: J2B2A4N98G)	
magnesium stearate (UNII: 70097M6B0)	
methylparaben (UNII: A2I8C7HI9T)	
shellac (UNII: 46N107B71O)	
polyethylene glycols (UNII: 3WJQ0SDW1A)	
povidone (UNII: FZ989GH94E)	
propylparaben (UNII: Z8IX2SC1OH)	
sodium benzoate (UNII: OJ245FE5EU)	
sorbitan monooleate (UNII: 06XEA2VD56)	
sucrose (UNII: C151H8M554)	
talc (UNII: 7SEV7J4R1U)	
titanium dioxide (UNII: 15FIX9V2JP)	
white wax (UNII: 7G1J5DA97F)	
diacetylated monoglycerides (UNII: 5Z17386USF)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	12;BI
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29485-6741-3	1 in 1 PACKAGE		
1		3 in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	01/01/2010	

Labeler - Mechanical Servants, Inc. (005530951)

Establishment

Name	Address	ID/FEI	Business Operations
Boehringer Ingelheim Promeco S.A. de C.V.		812579472	MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
Mechanical Servants, Inc.		005530951	RELABEL

Revised: 1/2012

Mechanical Servants, Inc.