

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled  
Meijer, Inc.**

-----  
**Nighttime Cold and Flu**

***Drug Facts***

***Active ingredients (in each softgel)***

Acetaminophen 325 mg

Dextromethorphan hydrobromide 15 mg

Doxylamine succinate 6.25 mg

***Purpose***

Pain reliever/fever reducer

Cough suppressant

Antihistamine

***Uses***

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat n headache n minor aches and pains
- fever
- runny nose and sneezing

***Warnings***

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, or occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur n avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed - **see Overdose warning.**
- do not exceed 4 doses per 24 hours

---

adults & children 12 years & over

2 softgels with water every 6 hours

children 4 to under 12 years

ask a doctor

children under 4 years

do not use

---

# When using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

## Other information

- store between 15°C to 30°C (59°F to 86°F)
- avoid excessive heat

## Inactive Ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, myglyol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

## Questions or comments?

1-888-333-9792

## Principal Display Carton

**meijer**

NDC 79481-002-7

COMPARE TO VICKS®  
NYQUIL® COLD & FLU  
LIQUICAPS®  
ACTIVE INGREDIENT!

**Nighttime Cold & Flu**

PAIN RELIEVER/FEVER REDUCER,  
COUGH SUPPRESSANT  
ANTIHISTAMINE &  
NASAL DECONGESTANT

Acetaminophen | 325 mg  
Dextromethorphan HBr | 15 mg  
Doxylamine Succinate | 6.25 mg

**24**  
SOFTGELS\*\*  
(\*\*Liquid-filled Capsules)

RELIEVES: ACHEs, FEVER,  
SORE THROAT, COUGH,  
RUNNY NOSE, SNEEZING

Actual Size

**meijer**

**Nighttime Cold & Flu**

**Drug Facts**

**Active ingredients (in each softgel)**

Active ingredients	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

**Uses** Temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, or occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma

**Drug Facts (continued)**

■ cough that occurs with too much phlegm (mucus)

■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are:**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

**When using this product**

- do not use more than directed
- drowsiness may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed - see **Overdose warning**.
- do not exceed 4 doses per 24 hours

Adults and children 12 years and over: 2 softgels with water every 6 hours

Children 4 to under 12 years: ask a doctor

Children under 4 years: do not use

**When using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

EXP. DATE: \_\_\_\_\_  
LOT NO: \_\_\_\_\_

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION  
TAMPER EVIDENT- DO NOT USE IF PACKAGING IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

how2recycle.info  
MULTI-LAYER  
PAPER  
BOX  
TRAY

This product is not manufactured or distributed by the Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and LiquiCaps®

MADE IN INDIA

Drug Facts (continued)  
Other information  
Ask a doctor before use if you have:  
■ severe asthma (COPD) ■ need a prescription  
■ severe liver disease ■ severe kidney disease

Active ingredients (D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, myglyol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide)

Questions or comments? 1-888-333-9792

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com  
ORIG 06/23

OUR QUALITY GUARANTEE  
The Meijer Family  
www.meijer.com/QUALITYGUARANTEE

PID 467803  
7 60236 30881 2

## NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-0027
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	902;215;AP02
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0027-2	2 in 1 CARTON	09/15/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	09/15/2022	

**Labeler** - Meijer, Inc. (006959555)

Revised: 12/2024

Meijer, Inc.