

**BRANTUSSIN DM- dextbrompheniramine maleate, dextromethorphan hbr,  
phenylephrine hcl liquid  
Brandywine Pharmaceuticals, LLC**

-----

**BranTussin DM**

**Active Ingredient**

Dexbrompheniramine Maleate 2 mg  
Dextromethorphan Hydrobromide 15 mg  
Phenylephrine Hydrochloride 7.5 mg

**Uses**

temporarily relieves these symptoms due to the common cold, hayfever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- cough due to minor throat and
- bronchial irritation
- nasal congestion
- reduces swelling of nasal passages

**Purposes**

Antihistamine

Cough Suppressant

Nasal Decongestant

**Warnings**

Do not exceed recommended dosage.

**Do not use this product**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional condition, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

- trouble urinating due to an enlarged prostate gland
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

### **Keep this and all drugs out of the reach of children**

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

### **When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- be careful when driving a motor vehicle or operating machinery

### **Ask a doctor or pharmacist before use**

if you are taking sedatives or tranquilizers.

### **If pregnant or breastfeeding**

ask a health professional before use.

### **Directions**

#### **Do not exceed recommended dosage**

---

Adults and children 12 years of age and over	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 6 teaspoonfuls in 24
Children 6 to under 12 years of age	1/2 teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 3 teaspoonfuls in 24 hours
Children under 6 years of age	Consult a doctor.

---

## Other Information

Store at 59° - 86°F (15° - 30°C)

## Inactive Ingredients

Anhydrous Citric Acid, Glycerin, Flavor, Propylene Glycol, Purified Water, Sodium Benzoate, Sodium Citrate Dihydrate, Sodium Saccharin, Sorbitol Solution.

## Questions? Comments?

Serious side effects associated with use of this product may be reported to this number. Call 1-800-882-1041 Mon. - Fri. (8 a.m. to 5 p.m. CST).

## Package Label



Drug Facts	
<b>Active Ingredients</b>	<b>Purpose</b>
<b>(in each 5 mL teaspoonful)</b>	
Dextrompheniramine Maleate, 2 mg.....	Antihistamine
Dextromethorphan HBr, 15 mg.....	Cough Suppressant
Phenylephrine HCl, 7.5 mg.....	Nasal Decongestant
<b>Uses:</b> temporarily relieves these symptoms due to the common cold; hay fever (allergic rhinitis) or other upper respiratory allergies:	
<ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ itching of the nose or throat</li> <li>■ itchy, watery eyes</li> <li>■ cough due to minor throat and bronchial irritation</li> <li>■ nasal congestion</li> <li>■ reduces swelling of nasal passages</li> </ul>	
<b>Warnings</b>	
Do not exceed recommended dosage. Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional condition, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	

Drug Facts (continued)	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> <li>■ a breathing problem such as emphysema or chronic bronchitis</li> <li>■ glaucoma</li> <li>■ trouble urinating due to an enlarged prostate gland</li> <li>■ a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema</li> <li>■ a cough that occurs with too much phlegm (mucus)</li> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes mellitus</li> </ul>	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.	
<b>When using this product</b>	
<ul style="list-style-type: none"> <li>■ excitability may occur, especially in children</li> <li>■ may cause marked drowsiness</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> </ul>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>■ nervousness, dizziness, or sleeplessness occur</li> <li>■ cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.</li> <li>■ new symptoms occur</li> </ul>	

Drug #	
If pregnant health prc	
Keep this reach of c	
In case of medical h	
Control Cx	
<b>Direct</b>	
Do not use	
Take unde	
healthcar	
Adults and 12 years of over:	
Children 6-12 years of	
Children ur years of ag	
<b>Other Info</b>	
Store at 15°	
Inactive In	
Citric Acid, Sodium Cit, Water, Sorb Flavor, Suc	
<b>Question</b>	
Serious side product may Call 800-314- Mon - Fri 8-	

## BRANTUSSIN DM

dexbrompheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71321-700
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	7.5 mg in 5 mL

<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 5 mL
<b>DEXBROMPHENIRAMINE MALEATE</b> (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII:75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	2 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71321-700-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/14/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/14/2022	

**Labeler** - Brandywine Pharmaceuticals, LLC (080581956)

Revised: 1/2024

Brandywine Pharmaceuticals, LLC