

**BACTIMICINA CHILDRENS ALLERGY- diphenhydramine hydrochloride liquid  
DLC Laboratories, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**BACTIMICINA<sup>®</sup> CHILDREN'S ALLERGY**

***Drug Facts***

**Active ingredient (in each 5 mL = 1 teaspoonful)**

Diphenhydramine HCl 12.5 mg

**Purpose**

Antihistamine

**Uses**

- Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat

**Warnings**

**Do not use**

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if the child has**

- a breathing problem such as chronic bronchitis
- glaucoma

**Ask a doctor or pharmacist before use if the child is** taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- find right dose on chart below
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours
- mL = milliliter; tsp = teaspoonful

<b>Age (yr)</b>	<b>Dose (mL or tsp)</b>
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL (1 tsp) to 10 mL (2 tsp)

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

## Other information

- **each 5 mL (1 tsp) contains:** sodium 2.3 mg
- store at room temperature. Protect from light. Store in outer carton until contents used.
- dosage cup provided
- **do not use if printed bottle wrap is missing or broken**

## Inactive ingredients

citric acid, D&C red no. 33, FD&C red no. 40, flavor, glycerin, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

## Questions

1-800-858-3889

Manufactured by  
DLC Laboratories, Inc.  
Paramount, CA 90723 USA

## PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton

4-6 Hour Relief

Voltee Para

Español

TRUSTED SINCE 1978

New

Look

Children's

Bactimicina®

Ages 6+ Years

Antihistamine • Liquid

Allergy

Diphenhydramine HCL (Antihistamine)

Sneezing • Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Alcohol Free

Cherry Flavor

4 FL OZ (118 mL) Dosing Cup Included

Do not use if printed bottle wrap is missing or broken.

**Drug Facts**

**Active ingredient** *Purpose*  
(in each 5 mL = 1 teaspoonful)  
Diphenhydramine HCl 12.5 mg.....Antihistamine

**Uses**

- Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat

**Warnings**  
Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

4-6 Hour Relief

TRUSTED SINCE 1978

**Children's Bactimicina**

Antihistamine • Liquid Ages 6+ Years

**Allergy**

Diphenhydramine HCl (Antihistamine)

Sneezing • Runny Nose  
Itchy, Watery Eyes • Itchy Throat or Nose

Alcohol Free

Cherry Flavor

4 FL OZ (118 mL)

Manufactured by / Fabricado por  
DLC Laboratories, Inc.  
Paramount, CA 90723 USA  
Questions / Preguntas  
1-800-858-3889  
dlclabs.com © 2023 DLC Laboratories, Inc.

@delacruzproducts @DLCLaboratories

**Drug Facts (continued)**

Ask a doctor before use if the child has

- a breathing problem such as chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

Peel Here  
Levante Aquí

**Drug Facts**  
**Datos Medicinales**

**Drug Facts (continued)**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- find right dose on chart below
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours
- mL = milliliter; tsp = teaspoonful

Age (yr)	Dose (mL or tsp)
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL (1 tsp) to 10 mL (2 tsp)

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**

- each 5 mL (1 tsp) contains: sodium 2.3 mg
- store at room temperature. Protect from light. Store in outer carton until contents used.
- dosage cup provided

**Inactive ingredients**  
citric acid, D&C red no. 33, FD&C red no. 40, flavor, glycerin, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

**Datos Medicinales**

**Ingrediente activo** *Propósito*  
(en cada 5 ml = 1 cucharadita)  
Difenhidramina HCl 12.5 mg.....Antihistamínico

**Usos** ■ alivio temporal de estos síntomas causados por la alergia al polen u otras alergias de las vías respiratorias superiores:

- goteo nasal
- estornudos
- lagrimeo y picazón en los ojos
- picazón en la nariz o la garganta

**Advertencias**  
No usar  
■ para hacer que un niño duerma

**Datos Medicinales (continuación)**

- con ningún otro producto que contenga difenhidramina, aunque sea de uso tópico en la piel

Consulte con un médico antes de usar este producto si el niño

- padece problemas respiratorios como bronquitis crónica
- padece glaucoma

Consulte con un médico o farmacéutico antes de usar este producto si el niño está tomando sedantes o tranquilizantes

**Al usar este producto**

- puede presentarse una marcada somnolencia
- los sedantes y los tranquilizantes pueden aumentar la somnolencia
- puede causar nerviosismo, particularmente en niños

**Manténgase fuera del alcance de los niños.** En caso de sobredosis, pida ayuda médica o contactese con un Centro de Control de Envenenamientos inmediatamente.

**Instrucciones**

- encuentre la dosis correcta en la tabla de abajo
- tomar cada 4 a 6 horas, o según lo que indique un médico
- no tome más de 6 dosis en 24 horas
- ml = mililitro; cdtá. = cucharadita

Edad (años)	Dosis (ml o cdtá.)
niños menores de 2 años	no lo use
niños de 2 a 5 años	no lo use a menos que lo indique un médico
niños de 6 a 11 años	5 ml (1 cdtá.) a 10 ml (2 cdtá.)

**Atención:** Use únicamente el vaso dosificador incluido que fue diseñado especialmente para usarse con este producto. No use ningún otro dosificador.

**Otra información**

- cada cucharadita (5 ml) contiene: sodio 2.3 mg
- consérvese a temperatura ambiente. Proteger de la luz. Guarde en la caja hasta terminar el contenido.
- incluye vaso dosificador

**Ingredientes inactivos**  
ácido cítrico, D&C rojo no. 33, FD&C rojo no. 40, saborizante, glicerina, benzoato de sodio, citrato de sodio, sorbitol, sucralosa, agua, goma xantana

P0186-AYX

XAY-8910d

# BACTIMICINA CHILDRENS ALLERGY

diphenhydramine hydrochloride liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:24286-1565
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24286-1565-4	1 in 1 CARTON	06/17/2015	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/17/2015	

**Labeler** - DLC Laboratories, Inc. (093351930)

**Establishment**

Name	Address	ID/FEI	Business Operations
DLC Laboratories, Inc.		093351930	manufacture(24286-1565) , label(24286-1565)

Revised: 7/2023

DLC Laboratories, Inc.