

**TUBERCULINUM RESIDUUM- tuberculinum residuum pellet
Boiron**

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

TUBERCULINUM RESIDUUM 15C MD

TUBERCULINUM RESIDUUM 15C

(**contains 0.443 mg of the active ingredient per pellet)

Relieves stiffness and pain associated with minor arthritis*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breast-feeding ask a health professional before use.

Keep out of reach of children.

Do not use if pellet dispenser seal is broken.

Contains approx 80 quick dissolving pellets per tube.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

1-800-BOIRON-1 (1-800-264-7661),

BoironUSA.com Info@boiron.com

Distributed by Boiron, Inc. Newtown Square, PA 19073

Tuberculinum residuum 15 C
 Relieves stiffness and pain associated with minor arthritis*

BOIRON®
 HOMEOPATHIC MEDICINE Made in France

HPUS NDC 0220-5065-41
 Do not use if pellet dispenser seal is broken

LOT: EXP:
 3 06965 06541 3

Contains approx. 80 pellets. US
 Peel for Drugs Facts and instructions for use.

Tuberculinum residuum 15 C

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE. NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

Drug Facts

Active ingredient*: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: ■ Do not use if pellet dispenser seal is broken.

Drug Facts (continued)

Inactive ingredients: lactose, sucrose

Questions or comments?
 BoironUSA.com Info@Boiron.com
 1-800-BOIRON-1 (1-800-264-7661)
 *C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.
 Distributed by Boiron Inc.
 Newtown Square, PA 19073

How to dispense pellets?

1 Turn tube upside down.
 2 Twist until 5 pellets are dispensed into cap.
 3 Carefully remove cap and use it to pour pellets under the tongue.

TUBERCULINUM RESIDUUM

tuberculinum residuum pellet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-5065
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Boiron of

Ingredient Name		Basis of Strength	Strength	
TUBERCULINUM RESIDUUM (UNII: X4R8E2P3V3) (TUBERCULINUM RESIDUUM - UNII:X4R8E2P3V3)		TUBERCULINUM RESIDUUM	15 [hp_C] in 15 [hp_C]	
Inactive Ingredients				
Ingredient Name			Strength	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)				
SUCROSE (UNII: C151H8M554)				
Product Characteristics				
Color	white	Score		
Shape	ROUND	Size	4mm	
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-5065-41	15 [hp_C] in 1 TUBE; Type 0: Not a Combination Product	10/14/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved homeopathic		10/14/2022		

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment

Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-5065)

Revised: 1/2023

Boiron