

**COLD AND FLU RELIEF NIGHTTIME, MULTI-SYMPTOM- acetaminophen,  
dextromethorphan hbr, doxylamine succinate capsule, liquid filled  
ARMY AND AIR FORCE EXCHANGE SERVICE**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Exchange Select 44-660**

***Active ingredients  
(in each liquid-filled capsule)***

Acetaminophen 325 mg  
Dextromethorphan HBr 15 mg  
Doxylamine succinate 6.25 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine

***Uses***

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - minor aches and pains
  - fever
  - headache
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not take more than directed**
- do not take more than 8 capsules per 24 hours
- adults and children 12 years and over: take 2 capsules with water every 6 hours
- children under 12 years: ask a doctor

***Other information***

- protect from heat, humidity and light
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol

### ***Questions or comments?***

**1-800-426-9391**

### **Principal Display Panel**

exchange<sup>®</sup>select<sup>™</sup>

**Compare to the Active Ingredients of  
Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu LiquiCaps<sup>®</sup> \***

**NightTime  
Cold & Flu Relief  
MULTI-SYMPTOM**

**Acetaminophen,**  
Dextromethorphan HBr, Doxylamine succinate

- Pain Reliever/Fever Reducer
- Cough Suppressant
- Antihistamine

Actual Size  
**24 Liquid Caps**

□ quality value

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu LiquiCaps<sup>®</sup>.

**Product of China  
Packaged and Quality Assured in the USA**

**"SATISFACTION GUARANTEED OR YOUR MONEY BACK"**

Manufactured For your Military Exchanges

Distributed by: LNK International, Inc.

Hauppauge, NY 11788

1-800-426-9391

50844 REV0418D66008

exchange select

NightTime

Cold & Flu Relief  
MULTI-SYMPTOM

exchange select

Compare To The Active Ingredients of  
Vicks® NyQuil® Cold & Flu LiquiCaps®\*

NightTime

Cold &amp; Flu Relief

MULTI-SYMPTOM

Acetaminophen,  
Dextromethorphan HBr, Doxylamine succinate

- Pain Reliever/Fever Reducer
- Cough Suppressant
- Antihistamine



Actual Size

24 Liquid Caps

quality  
valueNo print area  
Lot no. & Exp. dateTAMPER EVIDENT: DO NOT USE IF  
PACKAGE IS OPENED OR IF BLISTER  
UNIT IS TORN, BROKEN OR SHOWS  
ANY SIGNS OF TAMPERING

\*Satisfaction Guaranteed or Your Money Back\*  
Manufactured For Your Military Exchanges  
Distributed by: LNK International, Inc.  
Hauptpaug, NY 11788  
1-800-426-9391

\*This product is not manufactured or  
distributed by The Procter & Gamble Company,  
owner of the registered trademark Vicks®  
NyQuil® Cold & Flu LiquiCaps®.

Product of China. Packaged and  
Quality Assured in the USA

50844 REV0418D66008

## Drug Facts (continued)

**Drug Facts (continued)**

■ Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F). ■ See end flap for expiration date and lot number

**Inactive ingredients** D & C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol

**Questions or comments?** 1-800-426-9391

## Drug Facts (continued)

**Other information** ■ Protect from heat, humidity and light

■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**

■ Do not take more than directed

■ Do not take more than 8 capsules per 24 hours

■ Adults and children 12 years and over: take 2 capsules with water every 6 hours

■ Children under 12 years: ask a doctor

## Drug Facts

KEEP OUTER PACKAGE FOR  
COMPLETE PRODUCT INFORMATION

## Active ingredients

(in each liquid-filled capsule)

Acetaminophen 325 mg ..... Pain reliever/fever reducer

Dextromethorphan HBr 15 mg ..... Cough suppressant

Doxylamine succinate 6.25 mg ..... Antihistamine

## Uses

■ Temporarily relieves common cold and flu symptoms:

■ sore throat ■ minor aches and pains

■ headache ■ fever ■ runny nose and sneezing

■ cough due to minor throat and bronchial irritation

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI, ask a doctor or pharmacist before taking this product.

■ If you have ever had an allergic reaction to this product or any of its ingredients ■ To make a child sleepy

## Drug Facts (continued)

■ With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

■ a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema

■ difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

■ taking the blood thinning drug warfarin

■ taking sedatives or tranquilizers

**When using this product** ■ Do not exceed recommended dosage

■ avoid alcoholic beverages

■ marked drowsiness may occur

■ alcohol, sedatives, and tranquilizers may increase drowsiness

■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

■ pain or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

B-0066-660-08-R  
REV0418D66008

## Exchange Select 44-660

**COLD AND FLU RELIEF NIGHTTIME, MULTI-SYMPTOM**

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55301-660
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

**Product Characteristics**

<b>Color</b>	GREEN (clear)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	660
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-660-08	2 in 1 CARTON	03/01/2015	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL	part341	03/01/2015	

**Labeler** - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(55301-660)

Revised: 9/2019

ARMY AND AIR FORCE EXCHANGE SERVICE