

**DAYTIME NIGHTTIME SEVERE COLD AND FLU RELIEF- acetaminophen,
dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl
Rite Aid Corporation**

Rite Aid Corporation Daytime Nighttime Severe Cold & Flu Relief Drug Facts

**Nighttime Severe Cold & Flu
Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 44 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C

yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Daytime Severe Cold & Flu **Active ingredients - (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours

- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

COMBO PACK

SEE NEW WARNINGS

Compare to the active ingredients of Vicks® DayQuil® Severe Cold & Flu and Vicks® NyQuil® Severe Cold & Flu

MAXIMUM STRENGTH

daytime

severe

cold & flu relief

acetaminophen

phenylephrine HCl

dextromethorphan HBr

guaifenesin

pain reliever/fever reducer

nasal decongestant

cough suppressant

expectorant

NON-DROWSY

ALCOHOL FREE

ANTIHISTAMINE FREE

multi-symptom relief

12 FL OZ (355 mL)

MAXIMUM STRENGTH

nighttime

severe

cold & flu relief

acetaminophen

dextromethorphan HBr

doxylamine succinate

phenylephrine HCl

pain reliever/fever reducer

nasal decongestant

cough suppressant
 antihistamine
 multi-symptom relief
 10% ALCOHOL
 12 FL OZ (355 mL)

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECK AND IS BROKEN OR MISSING

*These products are not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe Cold & Flu and Vicks® NyQuil® Severe Cold & Flu.

Drug Facts (continued)
Inactive ingredients: butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9280

RITE AID PHARMACY

SEE NEW WARNINGS

*Compare to the active ingredients of Vicks® DayQuil® Severe Cold & Flu and Vicks® NyQuil® Severe Cold & Flu

MAXIMUM STRENGTH daytime severe cold & flu relief

acetaminophen
 phenylephrine HCl
 dextromethorphan HBr
 guaifenesin

pain reliever/fever reducer
 nasal decongestant
 cough suppressant
 expectorant

NON-DROWSY ALCOHOL FREE ANTIHISTAMINE FREE

multi-symptom relief

12 FL OZ (355 mL)

COMBO PACK

MAXIMUM STRENGTH nighttime severe cold & flu relief

acetaminophen
 dextromethorphan HBr
 doxylamine succinate
 phenylephrine HCl

pain reliever/fever reducer
 nasal decongestant
 cough suppressant
 antihistamine

multi-symptom relief

10% ALCOHOL

12 FL OZ (355 mL)

IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.

SATISFACTION GUARANTEED

DISTRIBUTED BY:
 RITE AID
 30 HUNTER LANE
 CAMP HILL, PA 17011

Drug Facts (continued)
Inactive ingredients: alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-800-719-9280

CODE AREA

RA# 372997

Nighttime Severe Cold & Flu		Daytime Severe Cold & Flu	
Drug Facts	Drug Facts	Drug Facts	Drug Facts
Active ingredients (in each 30 mL)	Active ingredients (in each 15 mL)	Active ingredients (in each 30 mL)	Active ingredients (in each 15 mL)
Acetaminophen HBr 20 mg.....Pain reliever/fever reducer	Acetaminophen 325 mg.....Pain reliever/fever reducer	Dextromethorphan HBr 20 mg.....Cough suppressant	Dextromethorphan HBr 10 mg.....Cough suppressant
Doxylamine succinate 1.25 mg.....Antihistamine	Guafenesin 200 mg.....Expectorant	Phenylephrine HCl 10 mg.....Nasal decongestant	Phenylephrine HCl 5 mg.....Nasal decongestant
Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ minor aches and pains ■ headache ■ sore throat ■ runny nose and sneezing ■ cough to help you sleep ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ fever ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose	Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ headache ■ fever ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ sore throat ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ promotes nasal and/or sinus drainage	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients	Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients	Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ breathing problems such as emphysema or chronic bronchitis ■ thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ a sodium-restricted diet	Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ diabetes
Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin	Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin	When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ be careful when driving a motor vehicle or operating machinery ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks	When using this product ■ do not use more than directed
Stop use and ask a doctor if ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ you get nervous, dizzy or sleepless ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ take only as directed—see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs	Directions ■ take only as directed—see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs	adults & children 12 yrs & over 30 mL every 4 hrs	adults & children 12 yrs & over 30 mL every 4 hrs
children 4 to under 12 yrs ask a doctor	children 6 to under 12 yrs 15 mL every 4 hrs	children under 4 yrs do not use	children 4 to under 6 yrs ask a doctor
children under 4 yrs do not use	children under 4 yrs do not use		
Other information ■ each 30 mL contains sodium 44 mg ■ store at 20-25°C (68-77°F)	Other information ■ each 15 mL contains sodium 6 mg ■ store at 20-25°C (68-77°F). Do not refrigerate. ►		

20 09 20098 :



DAYTIME NIGHTTIME SEVERE COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl kit

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:11822-0328

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0328-1	1 in 1 CARTON; Type 0: Not a Combination Product	07/24/2015	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

Part 1 of 2

NIGHTTIME SEVERE COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

Product Information

Item Code (Source) NDC:11822-0511

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 355W5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	GREEN	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0511-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/23/2015	

Part 2 of 2

DAYTIME SEVERE COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Item Code (Source)	NDC:11822-0710
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	

XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color	ORANGE (clear)	Score	
Shape		Size	
Flavor	FRUIT, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0710-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/18/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/24/2015	

Labeler - Rite Aid Corporation (014578892)

Revised: 11/2024

Rite Aid Corporation