

**EXTRA STRENGTH PAIN RELIEF- acetaminophen, aspirin, caffeine tablet  
BETTER LIVING BRANDS, LLC**

-----  
**370R\_Albertsons\_21130-037\_Acetaminophen, Aspirin and Caffeine  
250/250/65 mg tablets**

***Drug Facts***

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*\*)

Caffeine 65 mg

\*\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever

Pain reliever

Pain reliever aid

***Uses***

Temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives

- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are taking**

- a prescription drug for diabetes, gout, or arthritis
- any other drug or are under a doctor's care for any serious condition

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

- feel faint

- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
  - ringing in the ears or loss of hearing occurs
  - painful area is red or swollen
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - any new symptoms occur

These could be signs of a serious condition

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

### ***Other information***

- store at 20°- 25°C (68°- 77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

### ***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?* call 1-877-290-4008**

Lot No.: Varnish Omit  
Exp. Date:



RD 23308  
S5954



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

| Active ingredients (in each caplet) | Purposes          |
|-------------------------------------|-------------------|
| Acetaminophen 250 mg                | Pain reliever     |
| Aspirin 250 mg (NSAID)**            | Pain reliever     |
| Caffeine 65 mg                      | Pain reliever aid |

Uses

Temporarily relieves minor aches and pains due to:  
 ■ headache ■ a cold ■ arthritis ■ muscular aches  
 ■ toothache ■ premenstrual and menstrual cramps

Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ hives ■ rash ■ If a skin reaction occurs, stop use and seek medical help right away.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Stomach bleeding warning:** This product contains an NSAID.

Drug Facts (continued)

which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed  
**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.  
**Do not use** ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
**Ask a doctor before use if** ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma  
**Ask a doctor or pharmacist before use if you are taking** ■ a prescription drug for diabetes, gout, or arthritis ■ any other drug or are under a doctor's care for any serious condition  
**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools

Drug Facts (continued)

■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur  
 These could be signs of a serious condition  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions** ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years; ask a doctor  
**Other information** ■ store at 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package  
**Inactive ingredients:** carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide  
**Questions or comments?** 1-877-290-4008

Compare to Excedrin® Extra Strength active ingredients!  
 NOC 20190-037-742

**Extra Strength Pain Relief**

**ACETAMINOPHEN 250 mg**  
**ASPIRIN 250 mg (NSAID)**  
**CAFFEINE 65 mg**

Actual Size

**24 CAPLETS\***  
 (\*Caplet-shaped tablets)

Pain Reliever/Pain Reliever Aid  
 - Headache Relief

Signature Care  
 Quality Guaranteed

E3144R  
Varnish/Print Omit Area

This product is not a manufacturer or distributor of  
 Gastrointestinal Consumer Health Care Holdings (GCHC), LLC  
 owner of the registered trademark Excedrin® Extra Strength.  
 P.O. BOX 997, PLEASANTON, CA 94566-0099  
 1-888-723-8279  
 3708-1123

PAPER PLASTIC  
 BOX BOTTLE

how2recycle.info

NO IT'S NOT  
 LOVE IT







Quality Guaranteed

NDC 2130-037-02

# Extra Strength Pain Relief

**ACETAMINOPHEN 250 mg**  
**ASPRIN 250 mg (NSAID)**  
**CAFFEINE 65 mg**

Pain Reliever/Pain Reliever Aid  
- Headache Relief

**24 CAPLETS\***  
(\* Capsule-shaped tablets)

**READ AND KEEP CARTON FOR COMPLETE PRODUCT INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

## Drug Facts

### Active Ingredients (in each caplet)

- Acetaminophen 250 mg..... Pain reliever
- Aspirin 250 mg (NSAID)\*..... Pain reliever
- Caffeine 65 mg..... Pain reliever aid
- \*\*nonsteroidal anti-inflammatory drug

Use temporarily relieves minor aches and pains due to: ■ headache ■ a cold ■ arthritis ■ muscular aches ■ to relieve ■ premenstrual and menstrual cramps ■ **Warnings** See carton for complete warnings

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Keep out of reach of children.**

**Directions:** ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

**Other information:** ■ store at 20°-25° C (68°-77°F); excursions permitted between 15°-30° C (59°-86°F) ■ use by expiration date on package

**Questions or comments?** 1-877-294-4008

SS954  
DISTRIBUTED BY: BETTER LIVING BRANDS, LLC  
P.O. BOX 98, PLEASANTON, CA 94666-0098  
1-888-723-3929 370R 0223



RD 23054

**LOVE IT**  
**OR IT'S ON US!**

Lot No.:  
Exp. Date:  
Varnish Omit Area

Varnish/Print Omit Area

Exp. Date: Lot No.:

FD 23 054  
37M 0223  
1-888-723-3929  
P.O. BOX 98  
PLEASANTON, CA 94666-0098  
© 2011 B.L.B.

**RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active Ingredients (in each caplet)**

Acetaminophen 250 mg..... Pain reliever  
Aspirin 250 mg (NSAID)\*..... Pain reliever  
Caffeine 65 mg..... Pain reliever aid

**Use**

Temporarily relieves minor aches and pains due to: ■ headache ■ a cold ■ arthritis ■ muscular aches ■ to relieve ■ premenstrual and menstrual cramps

**Warnings**

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Liver warning:** Acetaminophen may cause severe liver damage. Do not use more than directed. Do not use with other drugs containing acetaminophen. Severe liver damage may occur if you take: ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Drug Facts (continued)**

**Directions:** ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

**Other information:** ■ store at 20°-25° C (68°-77°F); excursions permitted between 15°-30° C (59°-86°F) ■ use by expiration date on package

**Questions or comments?** 1-877-294-4008

**Drug Facts (continued)**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-525-0222) right away. Seek medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions:** ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

**Other information:** ■ store at 20°-25° C (68°-77°F); excursions permitted between 15°-30° C (59°-86°F) ■ use by expiration date on package

**Questions or comments?** 1-877-294-4008

Varnish Omit

Compare to Brand? (with imprint)

NDC 2130-037-02

**Extra Strength Pain Relief**

**ACETAMINOPHEN 250 mg**  
**ASPRIN 250 mg (NSAID)**  
**CAFFEINE 65 mg**

Pain Reliever/Pain Reliever Aid  
- Headache Relief

100 CAPLETS\*  
(\* Capsule-shaped tablets)

E6011AR5

This product is not manufactured or distributed by Signature Care. Contact the manufacturer for more information.

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**



Varnish Omit

Signature Care Quality Guaranteed



**STARCH, CORN** (UNII: O8232NY3SJ)

**CARNAUBA WAX** (UNII: R12CBM0EIZ)

**POVIDONE** (UNII: FZ989GH94E)

**SODIUM STARCH GLYCOLATE TYPE A POTATO** (UNII: 5856J3G2A2)

**STEARIC ACID** (UNII: 4ELV7Z65AP)

### Product Characteristics

|                 |                                 |                     |          |
|-----------------|---------------------------------|---------------------|----------|
| <b>Color</b>    | white                           | <b>Score</b>        | no score |
| <b>Shape</b>    | CAPSULE (Capsule shaped tablet) | <b>Size</b>         | 18mm     |
| <b>Flavor</b>   |                                 | <b>Imprint Code</b> | TCL;370  |
| <b>Contains</b> |                                 |                     |          |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:21130-037-01 | 1 in 1 CARTON                                      | 06/01/2023           |                    |
| 1 |                  | 100 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |
| 2 | NDC:21130-037-42 | 1 in 1 CARTON                                      | 06/01/2023           |                    |
| 2 |                  | 24 in 1 BOTTLE; Type 0: Not a Combination Product  |                      |                    |
| 3 | NDC:21130-037-20 | 200 in 1 BOTTLE; Type 0: Not a Combination Product | 07/17/2023           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013                                     | 06/01/2023           |                    |

**Labeler** - BETTER LIVING BRANDS, LLC (009137209)

**Registrant** - TIME CAP LABORATORIES, INC (037052099)

### Establishment

| Name                       | Address | ID/FEI    | Business Operations    |
|----------------------------|---------|-----------|------------------------|
| TIME CAP LABORATORIES, INC |         | 037052099 | manufacture(21130-037) |

Revised: 3/2024

BETTER LIVING BRANDS, LLC