

IBUPROFEN- ibuprofen tablet
BETTER LIVING BRANDS, LLC

601R Albertsons_21130-261_Ibuprofen Tablets USP 200 mg (Brown round tablets)

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 1. headache
 2. toothache
 3. backache
 4. menstrual cramps
 5. the common cold
 6. muscular aches
 7. minor pain of arthritis

- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin,

ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 1. feel faint
 2. vomit blood
 3. have bloody or black stools
 4. have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 1. chest pain
 2. trouble breathing
 3. weakness in one part or side of body
 4. slurred speech
 5. leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? 1-877-290-4008

NDC 21130-261-01

Signature care
Quality Guaranteed

Ibuprofen
Tablets USP, 200 mg
Pain Reliever/
Fever Reducer
(NSAID)
100 TABLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Ibuprofen 200 mg (NSAID)Pain reliever/Fever reducer, nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ● headache ● toothache ● backache ● menstrual cramps
 ● the common cold ● muscular aches ● minor pain of arthritis
 ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions ■ **do not take more than directed** ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ● if pain or fever does not respond to 1 tablet, 2 tablets may be used ● do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ● ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Questions or comments? 1-877-290-4008
 DISTRIBUTED BY: BETTER LIVING BRANDS LLC
 P.O. BOX 98, PLEASANTON, CA 94566-0098
 1-888-723-3929 Made in India 601R 0223

PLASTIC BOTTLE
 how2recycle.info
 RD 23054

LOVE IT OR IT'S ON-US
 Lot No.:
 Exp. Date:

Varnish Omit Area



Quality Guaranteed

NDC 21130-261-05

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer
(NSAID)

50 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID) *Pain reliever,
Fever reducer

* nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains, due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

Drug Facts (continued under label)

Lot No.:

Exp. Date:

Varnish Omit Area



- take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- If pregnant or breast-feeding,** ask a health professional before use.
- Keep out of reach of children.**
- Directions**
- do not take more than directed
 - the smallest effective dose should be used
 - adults and children 12 years and over:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
 - children under 12 years: ask a doctor
- Questions or comments?**
1-877-290-4008
- DISTRIBUTED BY:**
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0009
41-888-723-3929
RD 23054
Made in India
601R 0223 S5934
- LOVE IT**
OR IT'S ON US +
- PLASTIC BOTTLE**
how2recycle.info

Adhesive Area

Lot No.:
Exp. Date:



Some here for more information
Visit www.betterliving-brands.com
or call 1-800-99-
BETTER LIVING BRANDS, LLC
P.O. Box 99
PLEASANTON, CA 94566-0099
4-86-86
1-86-22-3833
© 2013
All trademarks are the property of their respective owners.

**LOVE IT
OR IT'S ON US!**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet) **Purpose**
Ibuprofen 200 mg (NSAID)* Pain reliever/
Fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
● headache ● toothache
● backache ● menstrual cramps ● the common cold ● muscular aches ● minor pain of arthritis ■ temporarily reduces fever

Warnings
Allergic alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness/itching ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Directions ■ do not take more than directed tablets and:
● the smallest effective dose should be used
● adults and children 12 years and over:
■ take 1 tablet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
● children under 12 years:
■ ask a doctor



Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use:
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor or health care provider if:
■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you're on a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are:
■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug
When using this product:
■ take with food or milk if stomach upset occurs
■ stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding: ● heart burn ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better

Directions ■ do not take more than directed tablets and:
● the smallest effective dose should be used
● adults and children 12 years and over:
■ take 1 tablet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
● children under 12 years:
■ ask a doctor

Other information ■ read all warnings and directions before use
■ store at 20-25°C (68-77°F)
■ see also www.betterliving-brands.com for complete directions
■ see also www.betterliving-brands.com for complete directions
■ see also www.betterliving-brands.com for complete directions

Questions or comments? 1-877-994-008
Pregnant or breastfeeding? See the label for complete directions.
Pregnant or breastfeeding? See the label for complete directions.
Pregnant or breastfeeding? See the label for complete directions.

Active ingredients ibuprofen, hydroxyethylcellulose, croscarmellose sodium, iron oxide red, polyethylene glycol, polyvinyl alcohol, piperazine stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide
Directions before use
■ store at 20-25°C (68-77°F)
Drug Facts (continued)

Compare to Advil® Tablets active ingredient! NDC 21130-261-05

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/
Fever Reducer
(NSAID)



ACTUAL SIZE
50 TABLETS



Quality Guaranteed

Varnish Omit

Varnish Omit

9286 Varnish/Print Omit Area

Adhesive Area

- take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**
- Directions**
 - do not take more than directed
 - the smallest effective dose should be used
 - adults and children 12 years and over:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
 - children under 12 years: ask a doctor

Questions or comments?

1-877-290-4008

DISTRIBUTED BY:

BETTER LIVING BRANDS LLC

P.O. BOX 99

PLEASANTON, CA 94566-0099

+1-888-723-3929

Made in India

601R 0223 55934



RD 23054
 +1-888-723-3929
 Made in India
 601R 0223 55934

NDC 21130-261-42



Ibuprofen
 Tablets USP, 200 mg
 Pain Reliever/Fever Reducer
 (NSAID)

24 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)* Pain reliever/
Fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems

Drug Facts (continued under label)

Lot No.:

Exp. Date:

Varnish Omit Area



Varnish Omit
Lot No.:
Exp Date:



SS864
RD 2364
LOVE IT OR IT'S ON US+
P.O. BOX 89
LETTERBIDGE PARK, LLC
PLEASANTON, CA 94688-0089
408-753-3669
www.lovetheproduct.com
© 2011
All rights reserved. No part of this publication may be reproduced without the prior written permission of the publisher.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts
Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID) Pain reliever/
Fever reducer
Nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and
pains due to:
● headache ● toothache
● backache ● menstrual cramps ● the
common cold ● muscle aches ● minor pain
of arthritis ● Temporarily reduces fever

Warnings
Able to get a heart attack or stroke. NSAIDs, except
aspirin, increase the risk of heart attack, heart
failure, and stroke. These can be fatal. The risk is
higher if you use more than directed or for longer
than directed.
Do not use
● if you have ever had an allergic reaction to any
other pain reliever, fever reducer,
● aspirin, or a heart/surgery
● aspirin or a heart/surgery
Ask a doctor before use if:
● so much to be feeling
warming up to you ● you have problems or
serious side effects from taking pain relievers or
fever reducers ● you have a history of stomach
problems, such as heartburn ● you have high
blood pressure, heart disease, liver cirrhosis,
kidney disease, asthma, or had a stroke ● you
are taking a diuretic

Directions
● do not take more than directed
● the smallest effective dose should be used
● take 1 tablet every 4 to 6 hours
while symptoms persist
● if pain or fever does not respond
to 1 tablet, 2 tablets may be used
● do not exceed 6 tablets in 24
hours, unless directed by a
doctor
● ask a doctor
children under
12 years:
● ask a doctor



how2recycle.com

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except
aspirin, increase the risk of heart attack, heart
failure, and stroke. These can be fatal. The risk is
higher if you use more than directed or for longer
than directed.
Do not use
● if you have ever had an allergic reaction to any
other pain reliever, fever reducer,
● aspirin, or a heart/surgery
● aspirin or a heart/surgery
Ask a doctor before use if:
● so much to be feeling
warming up to you ● you have problems or
serious side effects from taking pain relievers or
fever reducers ● you have a history of stomach
problems, such as heartburn ● you have high
blood pressure, heart disease, liver cirrhosis,
kidney disease, asthma, or had a stroke ● you
are taking a diuretic

Directions
● do not take more than directed
● the smallest effective dose should be used
● take 1 tablet every 4 to 6 hours
while symptoms persist
● if pain or fever does not respond
to 1 tablet, 2 tablets may be used
● do not exceed 6 tablets in 24
hours, unless directed by a
doctor
● ask a doctor
children under
12 years:
● ask a doctor

Directions
● do not take more than directed
● the smallest effective dose should be used
● take 1 tablet every 4 to 6 hours
while symptoms persist
● if pain or fever does not respond
to 1 tablet, 2 tablets may be used
● do not exceed 6 tablets in 24
hours, unless directed by a
doctor
● ask a doctor
children under
12 years:
● ask a doctor

Directions
● do not take more than directed
● the smallest effective dose should be used
● take 1 tablet every 4 to 6 hours
while symptoms persist
● if pain or fever does not respond
to 1 tablet, 2 tablets may be used
● do not exceed 6 tablets in 24
hours, unless directed by a
doctor
● ask a doctor
children under
12 years:
● ask a doctor

Directions
● do not take more than directed
● the smallest effective dose should be used
● take 1 tablet every 4 to 6 hours
while symptoms persist
● if pain or fever does not respond
to 1 tablet, 2 tablets may be used
● do not exceed 6 tablets in 24
hours, unless directed by a
doctor
● ask a doctor
children under
12 years:
● ask a doctor

Drug Facts (continued)
Directions to use:
● store at 20°-25°C (68°-77°F)
Inactive ingredients: calcium silicate, iron oxide red,
magnesium stearate, microcrystalline cellulose,
polyethylene glycol, polyvinyl alcohol
pregelatinized starch, talc, titanium dioxide
Questions or comments? 1-877-293-4008
TALK TO YOUR DOCTOR OR THE PHARMACEUTICAL RESEARCH AND MANUFACTURING INDUSTRY OF AMERICA

Compare to
Advil® Tablets
active ingredient!
NDC 21130-261-42

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/
Fever Reducer
(NSAID)



ACTUAL SIZE
24 TABLETS



Quality Guaranteed

Varnish
Omit

Varnish
Omit

926 E Varnish/Print Omit Area

VALUE PACK



Quality Guaranteed

Compare to Advil® Tablets active ingredient* NDC 2130-261-56

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/
Fever Reducer
(NSAID)



ACTUAL SIZE

1000 TABLETS

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)* Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



RD 23054

Scan for more information

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® tablets

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94698-0099
Lot No. 55624
Made in India 601 R 0223 55624

LOVE IT OR IT'S ON US!

Varnish Omit Area
Exp. Date:

Adhesive Area

Other information read all warnings and directions before use. Store at 20-25°C (68-77°F).
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.
Questions or comments? 1-877-290-4008

Directions do not take more than directed. The smallest effective dose should be used.
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
ask a doctor

Warnings Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters.
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear.

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you, you have problems or serious side effects from taking pain relievers or fever reducers, you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic.

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear.

Other information read all warnings and directions before use. Store at 20-25°C (68-77°F).
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.
Questions or comments? 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



Compare to Advil® Tablets active ingredient* NDC 2130-261-53

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/
Fever Reducer
(NSAID)



ACTUAL SIZE
500 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts
Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)* Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis.
temporarily reduces fever.

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters.
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you, you have problems or serious side effects from taking pain relievers or fever reducers, you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic.

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear.

Other information read all warnings and directions before use. Store at 20-25°C (68-77°F).
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.
Questions or comments? 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® tablets

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94698-0099
Lot No. 55624
Made in India 601 R 0223 55624



Scan for more information

LOVE IT OR IT'S ON US!

Varnish Omit Area
Exp. Date:

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	114
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-261-54	250 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2023	
2	NDC:21130-261-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2023	
3	NDC:21130-261-56	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/22/2023	
4	NDC:21130-261-05	1 in 1 CARTON	08/30/2023	
4		50 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:21130-261-01	1 in 1 CARTON	08/30/2023	
5		100 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:21130-261-42	1 in 1 CARTON	09/15/2023	
6		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	06/12/2023	

Labeler - BETTER LIVING BRANDS, LLC (009137209)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(21130-261)