

IBUPROFEN- ibuprofen capsule, liquid filled
BETTER LIVING BRANDS, LLC

Albertsons Ibuprofen Capsules USP, 200 mg and Ibuprofen Capsules USP, 200 mg (mini capsules)

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to

200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F).

Inactive ingredients

ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments?



NDC 21130-361-80

Ibuprofen

CAPSULES 200 mg

Pain Reliever/Fever Reducer
(NSAID)

80 SOFTGELS**
(**Liquid-filled capsules)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*	Pain reliever/
(present as the free acid and potassium salt)	Fever reducer
*nonsteroidal anti-inflammatory drug	

Uses ■ temporarily relieves minor aches and pains due to:
headache, toothache, backache, menstrual cramps, the
common cold, muscular aches, minor pain of arthritis
■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Questions or comments? 1-877-290-4008 RD 23048

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0099
1-888-723-3929
Made in India
610T 0223 S5934



0419623204

Lot No.:
Exp. Date:

Varnish Omit Area

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule) Purpose
Sodium ibuprofen (NSAID) * Pain reliever/fever reducer
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

Drug Facts (continued)

- hives
 - facial swelling
 - asthma (wheezing)
 - stomach
 - skin redness
 - rash
 - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, even aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

- Do not use**
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
 - right before or after heart surgery
- Ask a doctor before use if**
- stomach bleeding warning applies to you
 - you have problems or serious side effects from taking pain relievers or fever reducers
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 - you are taking a diuretic
- Ask a doctor or pharmacist before use if you are**
- under a doctor's care for any serious condition
 - taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
 - taking any other drug
- When using this product**
- take with food or milk if stomach upset occurs

Drug Facts (continued)

- Stop use and ask a doctor if**
- you experience any of the following signs of stomach bleeding:
 - belly pain
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
- If pregnant or breastfeeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
- Directions** do not take more than directed
- the smallest effective dose should be used
 - adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
 - children under 12 years: ask a doctor

Other information

- each capsule contains potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Drug Facts (continued)

Inactive ingredients: ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, sodium, sorbitan monoleate, sorbitol sorbitan

Questions or comments? 1-877-290-4000

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Advil® and Cellego®. Cellego® is the registered trademark of Liquid-Gels®.

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
1-800-723-3929
Made in India
6107 0223

DO NOT USE IF SEAL UNDER BOTTLE CAP
IMPRINTED WITH "SEALED FOR YOUR
PROTECTION" IS BROKEN OR MISSING



Varnish/Print
Omit Area
Lot No.:
Exp Date:



Quality Guaranteed

Varnish Omit

Ibuprofen
CAPSULES 200 mg
Pain Reliever/Fever Reducer
(NSAID)

Compare to
Advil® Liqui-Gels
active ingredient*

NDC 21130-361-80



80 SOFTGELS**
(**Liquid-filled capsule)

Varnish Omit

Varnish/Print Omit Area

C6011AR3

stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Questions or comments?
1-877-290-4008

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
1-888-723-3929
Made in India
610T 0223 S5934

RD 23048

 **PLASTIC BOTTLE**
how2recycleinfo

OR IT'S ON US *

1/16"
Margin



Quality
Guaranteed

NDC 21130-361-23

Ibuprofen

CAPSULES 200 mg

Pain Reliever/Fever Reducer
(NSAID)

20 SOFTGELS**
(**Liquid-filled capsules)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient
(in each capsule)

Solubilized ibuprofen equal to Pain reliever/200 mg ibuprofen (NSAID)* Fever reducer (present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had

Drug Facts (continued under label)

Lot No.: _____
 Price Here  Exp Date: _____

Varnish Omit Area

Outside



IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-361
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SHELLAC (UNII: 46N107B71O)	

AMMONIA (UNII: 5138Q19F1X)	
GELATIN (UNII: 2G86QN327L)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
WATER (UNII: 059QF0KO0R)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
SORBITOL (UNII: 506T60A25R)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-361-23	1 in 1 CARTON	10/15/2024	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-361-80	1 in 1 CARTON	10/15/2024	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	10/15/2024	

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-070
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
Inactive Ingredients				
Ingredient Name			Strength	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)				
SORBITOL (UNII: 506T60A25R)				
AMMONIA (UNII: 5138Q19F1X)				
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)				
GELATIN (UNII: 2G86QN327L)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SHELLAC (UNII: 46N107B71O)				
WATER (UNII: 059QF0KO0R)				
FERROUS OXIDE (UNII: G7036X8B5H)				
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)				
Product Characteristics				
Color	green	Score	no score	
Shape	CAPSULE	Size	13mm	
Flavor		Imprint Code	62	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-070-80	1 in 1 CARTON	10/15/2024	
1		80 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA079205	10/15/2024	

Labeler - BETTER LIVING BRANDS, LLC (009137209)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

MARKSANS PHARMA LIMITED		925822975	manufacture(21130-361, 21130-070)
-------------------------	--	-----------	-----------------------------------

Revised: 10/2024

BETTER LIVING BRANDS, LLC