

IBUPROFEN- ibuprofen capsule, liquid filled

Costco Wholesale

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F).
- swallow whole, do not crush, chew, or dissolve

Inactive ingredients

FD&C blue #1, FD&C yellow#6, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO ADVIL® LIQUI-GELS® active ingredient†

Ibuprofen

Solubilized Ibuprofen Capsules, 200 mg

Liquid Softgels

Pain Reliever/Fever Reducer **(NSAID)**

Safety Sealed

SOFTGELS**

(**LIQUID-FILLED CAPSULES)

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Distributed by:

PL Developments

Westbury, NY 11590 U.S.A.

For: Costco Wholesale Corporation

P.O.Box 34535

Seattle, WA 98124-1535 USA

1-800-774-2678

www.costco.com

Product Label

<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="text-align: left; padding: 2px;">Drug Facts</th> <th style="text-align: left; padding: 2px;">Purpose</th> </tr> <tr> <td style="padding: 2px;"> Active ingredient (in each capsule) Solubilized ibuprofen (as the 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) (nonsteroidal anti-inflammatory drug) </td> <td style="padding: 2px;"> Pain reliever Fever reducer </td> </tr> <tr> <td colspan="2" style="padding: 2px;"> Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscle aches ■ minor pain of arthritis ■ backaches ■ menstrual cramps ■ the common cold ■ temporarily reduces fever </td> </tr> <tr> <td colspan="2" style="padding: 2px;"> Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters If a allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ take had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take 3 or more alcoholic drinks every day while using this product Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or unusual side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug </td> </tr> <tr> <th style="text-align: left; padding: 2px;">Drug Facts (continued)</th> <td></td> </tr> <tr> <td colspan="2" style="padding: 2px;"> When using this product ■ take with food or milk, if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ heart burn ■ vomit blood ■ have blood or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ sudden dizziness ■ fainting ■ weakness in one part or side of body ■ pain in jaw or tooth more than 10 days ■ fever, pain, or rash more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear ■ If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. ■ Age and of most of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). </td> </tr> <tr> <td colspan="2" style="padding: 2px;"> Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 3 capsules, 3 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor </td> </tr> <tr> <td colspan="2" style="padding: 2px;"> Other information ■ each capsule contains: potassium 20 mg ■ store at 25-30°C (77-86°F) ■ excipient excipients list above 40°C (104°F) ■ read all warnings and directions before use. Keep carton. ■ swallow whole. Do not crush, chew, or dissolve. Inactive ingredients: PEG 400, 1, PEG 400, yellow no. 6, gelatin, sucrose (vegetarian), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, croscarmellose, hydroxypropyl methylcellulose, sodium acetate. </td> </tr> <tr> <td colspan="2" style="padding: 2px;"> Questions or comments? Call 1-877-733-3688 Monday-Friday 8AM-5PM EST *This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®. </td> </tr> </table>	Drug Facts	Purpose	Active ingredient (in each capsule) Solubilized ibuprofen (as the 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) (nonsteroidal anti-inflammatory drug)	Pain reliever Fever reducer	Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscle aches ■ minor pain of arthritis ■ backaches ■ menstrual cramps ■ the common cold ■ temporarily reduces fever		Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters If a allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ take had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take 3 or more alcoholic drinks every day while using this product Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or unusual side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug		Drug Facts (continued)		When using this product ■ take with food or milk, if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ heart burn ■ vomit blood ■ have blood or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ sudden dizziness ■ fainting ■ weakness in one part or side of body ■ pain in jaw or tooth more than 10 days ■ fever, pain, or rash more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear ■ If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. ■ Age and of most of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).		Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 3 capsules, 3 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor		Other information ■ each capsule contains: potassium 20 mg ■ store at 25-30°C (77-86°F) ■ excipient excipients list above 40°C (104°F) ■ read all warnings and directions before use. Keep carton. ■ swallow whole. Do not crush, chew, or dissolve. Inactive ingredients: PEG 400, 1, PEG 400, yellow no. 6, gelatin, sucrose (vegetarian), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, croscarmellose, hydroxypropyl methylcellulose, sodium acetate.		Questions or comments? Call 1-877-733-3688 Monday-Friday 8AM-5PM EST *This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.		<p style="text-align: center;">KIRKLAND Signature</p> <p style="text-align: center;">COMPARE TO ADVIL® LIQUI-GELS® active ingredient†</p> <p style="text-align: center; font-size: 2em;">IBUPROFEN</p> <p style="text-align: center;">Solubilized Ibuprofen Capsules, 200mg</p> <p style="text-align: center; border: 1px solid blue; border-radius: 10px; display: inline-block; padding: 2px;">Liquid Softgels</p> <p style="text-align: center; font-size: 1.5em;">Pain Reliever Fever Reducer (NSAID)</p> <p style="text-align: center; font-size: 1.5em;">Safety Sealed</p> <p style="text-align: center;">two 180 count bottles TOTAL</p> <p style="text-align: center; font-size: 2em; color: blue;">360 SOFTGELS** (**LIQUID-FILLED CAPSULES)</p> <p style="text-align: right;">NDC 63981-750-36 ITM. / ART. 196119</p>	
Drug Facts	Purpose																			
Active ingredient (in each capsule) Solubilized ibuprofen (as the 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) (nonsteroidal anti-inflammatory drug)	Pain reliever Fever reducer																			
Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscle aches ■ minor pain of arthritis ■ backaches ■ menstrual cramps ■ the common cold ■ temporarily reduces fever																				
Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters If a allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ take had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take 3 or more alcoholic drinks every day while using this product Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or unusual side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug																				
Drug Facts (continued)																				
When using this product ■ take with food or milk, if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ heart burn ■ vomit blood ■ have blood or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ sudden dizziness ■ fainting ■ weakness in one part or side of body ■ pain in jaw or tooth more than 10 days ■ fever, pain, or rash more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear ■ If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. ■ Age and of most of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).																				
Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 3 capsules, 3 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor																				
Other information ■ each capsule contains: potassium 20 mg ■ store at 25-30°C (77-86°F) ■ excipient excipients list above 40°C (104°F) ■ read all warnings and directions before use. Keep carton. ■ swallow whole. Do not crush, chew, or dissolve. Inactive ingredients: PEG 400, 1, PEG 400, yellow no. 6, gelatin, sucrose (vegetarian), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, croscarmellose, hydroxypropyl methylcellulose, sodium acetate.																				
Questions or comments? Call 1-877-733-3688 Monday-Friday 8AM-5PM EST *This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.																				

PLD-HS29A FCM004D

Lot No: 18V10092a

Exp. Date: 0 96619 19691 3

KIRKLAND Ibuprofen 200 mg Softgels

IBUPROFEN
 ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63981-750
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	AT146
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63981-750-36	2 in 1 BOX	07/31/2018	
1		360 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206999	07/31/2018	

Labeler - Costco Wholesale (103391843)

