MINOXIDIL FOR WOMEN- minoxidil aerosol, foam BosleyMD

Minoxidil (for Women)

Drug Facts

Drug Facts

Active ingredient

Minoxidil 5% w/w (without propellant)

Purpose

Hair regrowth treatment for women

Use

to regrow hair on the top of the scalp (see pictures on side of this carton)

Warnings

For external use only.

Extremely Flammable: Avoid fire, flame, or smoking during and immediately following application.

Do not use if

- your degree of hair loss is different than that shown on side of this carton because this product may not work for you
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- your hair loss is associated with childbirth
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have

heart disease

When using this product

- do not use more than directed
- do not apply on other parts of the body

- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 3 months with **once-daily** use. For some women, you may need to use this product **once a day** for at least 6 months before you see results.
- the amount of hair regrowth is different for each person. This product will not work for all women.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 6 months

Pregnant or breast-feeding

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- see enclosed leaflet for complete directions on how to use
- apply half a capful **once daily** directly to the scalp in the hair loss area
- massage into scalp with fingers, then wash hands well
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- store at controlled room temperature 20° to 25°C (68° to 77°F)
- contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120°F (49°C).

Inactive ingredients

butane, butylated hydroxytoluene, cetyl alcohol, citric acid, glycerin, isobutane, lactic acid, polysorbate 60, propane, purified water, SD alcohol 40-B, stearyl alcohol

Question?

Call 1-800-BOSLEY1 (1800-267-5391

Principal display panel

Minoxidil

Topical Aerosol,

5% (For Women)

HAIR REGROWTH TREATMENT

CLINICALLY PROVEN TO HELP REGROW HAIR

Once-a-Day

Foam for Women

Reactivates

Hair Follicles

Unscented

Read and keep carton and information leaflet for full product information

NOT FOR USE BY MEN

Distributed by: BosleyMD

800.BOSLEY1 bosleymd.com

© 2024 Bosley, Inc. All rights reserved.

Manufactured for Scientific Hair Research LLC,

Los Angeles, CA 90045

Package label











4: APPLY TO SCALP, NOT HAIR Part your hair to expose hair less area. Massage foam into scalp, not hair. Repeat until all hair loss areas have been covered.



5: CLOSE CAP AND WASH HAMDS Snap cap back into place. Be sure arrows do not line up, so cap remaises child resistant. Wash hands and any surface thoroughly after use.

Distributed by: BosleyMD 800.BOSLEY1 bosleymd.com © 2024 Bosley, Inc. All rights reserved. Manufactured for Scientific Hair Research LLC, Los Angeles, CA 90045 Product of Italy





BOSLEY MD Hair regrowth treatment for women

MINOXIDIL FOR WOMEN

minoxidil aerosol, foam

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:69113-502 Route of Administration TOPICAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1) MINOXIDIL 5965120SH1) MINOXIDIL 50 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
BUTANE (UNII: 6LV4FOR43R)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
ISOBUTANE (UNII: BXR49TP611)	
LACTIC ACID (UNII: 33X04XA5AT)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
PROPANE (UNII: T75W9911L6)	
WATER (UNII: 059QF0KO0R)	
ALCOHOL (UNII: 3K9958V90M)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	

Product Characteristics					
Color	white	Score			
Shape		Size			
Flavor		Imprint Code			
Contains					

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:69113-502- 01	1 in 1 PACKAGE	08/29/2025				
1		60 g in 1 CAN; Type 0: Not a Combination Product					
2	NDC:69113-502- 02	2 in 1 PACKAGE	08/29/2025				
2		60 g in 1 CAN; Type 0: Not a Combination Product					

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA208092	08/29/2025		

Labeler - BosleyMD (961897647)

Revised: 8/2025 BosleyMD