ECONAZOLE NITRATE- econazole nitrate cream Preferred Pharmaceuticals Inc.

Econazole Nitrate Cream 1%

Rx Only

FOR TOPICAL, DERMATOLOGIC USE ONLY – NOT FOR OPHTHALMIC, ORAL, OR INTRAVAGINAL USE.

Prescribing Information

DESCRIPTION

Econazole Nitrate Cream 1% contains the antifungal agent, econazole nitrate 1% in a water miscible base consisting of pegoxol 7 stearate, peglicol 5 oleate, mineral oil, benzoic acid, butylated hydroxyanisole, and purified water. The white to off-white soft cream is for topical use only.

Chemically, econazole nitrate is 1-[2-{(4-chloro-phenyl) methoxy}-2-(2,4-dichlorophenyl)ethyl]-1H-imidazole mononitrate. Its structure is as follows:

CLINICAL PHARMACOLOGY

After topical application to the skin of normal subjects, systemic absorption of econazole nitrate is extremely low. Although most of the applied drug remains on the skin surface, drug concentrations were found in the stratum corneum which, by far, exceeded the minimum inhibitory concentration for dermatophytes. Inhibitory concentrations were achieved in the epidermis and as deep as the middle region of the dermis. Less than 1% of the applied dose was recovered in the urine and feces.

Microbiology

Econazole nitrate has been shown to be active against most strains of the following microorganisms, both *in vitro* and in clinical infections as described in the INDICATIONS AND USAGE section.

Dermatophytes	Yeasts	
Epidermophyton floccosum	Candida albicans	
Microsporum audouini	Malassezia furfur	
Microsporum canis		
Microsporum gypseum		
Trichophyton mentagrophytes		
Trichophyton rubrum		
Trichophyton tonsurans		

Econazole nitrate exhibits broad-spectrum antifungal activity against the following organisms *in vitro*, **but the clinical significance of these data is unknown.**

Dermatophytes	Yeasts
Trichophyton verrucosum	Candida guillermondii
	Candida parapsilosis
	Candida tropicalis

INDICATIONS AND USAGE

ECONAZOLE NITRATE CREAM 1% is indicated for topical application in the treatment of tinea pedis, tinea cruris, and tinea corporis caused by *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton tonsurans*, *Microsporum canis*, *Microsporum audouini*, *Microsporum gypseum*, and *Epidermophyton floccosum*, in the treatment of cutaneous candidiasis, and in the treatment of tinea versicolor.

CONTRAINDICATIONS

ECONAZOLE NITRATE CREAM 1% is contraindicated in individuals who have shown hypersensitivity to any of its ingredients.

WARNINGS

ECONAZOLE NITRATE CREAM 1% is not for ophthalmic use.

PRECAUTIONS

General

If a reaction suggesting sensitivity or chemical irritation should occur, use of the medication should be discontinued.

For external use only. Avoid introduction of ECONAZOLE NITRATE CREAM 1% into the eyes.

Carcinogenicity Studies

Long-term animal studies to determine carcinogenic potential have not been performed.

Fertility (Reproduction)

Oral administration of econazole nitrate in rats has been reported to produce prolonged gestation. Intravaginal administration in humans has not shown prolonged gestation or other adverse reproductive effects attributable to econazole nitrate therapy.

Pregnancy

Pregnancy Category C

Econazole nitrate has not been shown to be teratogenic when administered orally to mice, rabbits or rats. Fetotoxic or embryotoxic effects were observed in Segment I oral studies with rats receiving 10 to 40 times the human dermal dose. Similar effects were observed in Segment II or Segment III studies with mice, rabbits and/or rats receiving oral doses 80 or 40 times the human dermal dose. Econazole nitrate should be used in the first trimester of pregnancy only when the physician considers it essential to the welfare of the patient. The drug should be used during the second and third trimesters of pregnancy only if clearly needed.

Nursing Mothers

It is not known whether econazole nitrate is excreted in human milk. Following oral administration of econazole nitrate to lactating rats, econazole and/or metabolites were excreted in milk and were found in nursing pups. Also, in lactating rats receiving large oral doses (40 or 80 times the human dermal dose), there was a reduction in post partum viability of pups and survival to weaning; however, at these high doses, maternal toxicity was present and may have been a contributing factor. Caution should be exercised when econazole nitrate is administered to a nursing woman.

ADVERSE REACTIONS

During clinical trials, approximately 3% of patients treated with econazole nitrate 1% cream reported side effects thought possibly to be due to the drug, consisting mainly of burning, itching, stinging, and erythema. One case of pruritic rash has also been reported.

OVERDOSE

Overdosage of econazole nitrate in humans has not been reported to date. In mice, rats, guinea pigs and dogs, the oral LD_{50} values were found to be 462, 668, 272, and >160 mg/kg, respectively.

To report **SUSPECTED ADVERSE REACTIONS**, contact Teligent Pharma, Inc. at 1-856-697-1441, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

Sufficient ECONAZOLE NITRATE CREAM 1% should be applied to cover affected areas once daily in patients with tinea pedis, tinea cruris, tinea corporis, and tinea versicolor, and twice daily (morning and evening) in patients with cutaneous candidiasis. Early relief of symptoms is experienced by the majority of patients and clinical improvement may be seen fairly soon after treatment is begun; however, candidal infections and tinea cruris and corporis should be treated for two weeks and tinea pedis for one month in order to reduce the possibility of recurrence. If a patient shows no clinical improvement after the treatment period, the diagnosis should be redetermined. Patients with tinea versicolor usually exhibit clinical and mycological clearing after two weeks of treatment.

HOW SUPPLIED

ECONAZOLE NITRATE CREAM 1% is supplied in tubes of;

15 grams (NDC68788-7046-1)

30 grams (NDC68788-7046-3)

Store at controlled room temperature 20° - 25°C (68° - 77°F).

Teligent Pharma, Inc.

Buena, New Jersey 08310


Rev. 01/2016

Relabeled By: Preferred Pharmaceuticals Inc.

PRINCIPAL DISPLAY PANEL - 30 g Tube Label NDC 68788-7406-3 ECONAZOLE NITRATE CREAM 1% 30 grams

Rx ONLY

FOR TOPICAL USE ONLY NOT FOR OPHTHALMIC USE




Generic for: Spectazole
Each gm contains: Econazole nitrate 1%

Pkg Size: Exp Date: Lot#: Batch#:

Ins: Mfg: Teligent Pharma, Inc. Prod#:

Warning

See package insert for full prescribing information. Do not use it sail has been punctured or is not visible. Store at 10 - 25 - (26 - 77 f. x. Only. For topical use only. Keep this and all medication out of the Feach of children.



ECONAZOLE NITRATE

econazole nitrate cream

Product Information

Product TypeHUMAN PRESCRIPTION DRUGItem Code (Source)NDC:68788-7406(NDC:52565-022)Route of AdministrationTOPICAL

Active Ingredient/Active Moiety

8		
Ingredient Name	Basis of Strength	Strength
Econazole Nitrate (UNII: H438 WYN10E) (Econazole - UNII:6Z1Y2V4A7M)	Econazole Nitrate	10 mg in 1 g

Inactive Ingredients Ingredient Name Strength Pegoxol 7 Stearate (UNII: 3EW5AXE5X5) PEG-5 Oleate (UNII: 0240 V77G50) Mineral Oil (UNII: T5L8T28FGP) Benzoic Acid (UNII: 8SKN0B0MIM) Butylated Hydroxyanisole (UNII: REK4960K2U) Water (UNII: 059QF0KO0R)

				Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date					
1 ND	OC:68788-7406-3	1 in 1 CARTON	06/06/2018						
1		30 g in 1 TUBE; Type 0: Not a Combination Product							
2 ND	DC:68788-7406-1	15 g in 1 TUBE; Type 0: Not a Combination Product	09/05/2019						

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA076574	06/06/2018			

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment				
Name	Address	ID/FEI	Business Operations	
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-7406)	

Revised: 5/2020 Preferred Pharmaceuticals Inc.