

**MULTI SYMPTOM COLD AND MULTI SYMPTOM NIGHTTIME COLD CHILDRENS-  
dextromethorphan hbr, guaifenesin, phenylephrine hci, acetaminophen,  
diphenhydramine hci, phenylephrine hci  
CVS Pharmacy**

-----

**DRUG FACTS**

**Active ingredients for Nighttime (in each 10 mL)**

Acetaminophen 325 mg

Diphenhydramine HCl 12.5 mg

Phenylephrine HCl 5 mg

**Active ingredients for Daytime (in each 5 mL)**

Dextromethorphan HBr 5 mg

Guaifenesin 100 mg

Phenylephrine HCl 2.5 mg

**Purpose for Nighttime**

**Pain reliever / fever reducer**

Antihistamine / Cough suppressant

Nasal Decongestant

**Purpose for Daytime**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

**Nighttime**

- temporarily relieves these common cold and flu symptoms
- minor aches and pains
- headache
- sore throat
- sneezing
- runny nose
- nasal congestion

- cough
- controls cough to help your child get to sleep
- temporarily reduces fever

### **Daytime**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make cough more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep
  - nasal congestion due to a cold
  - stuffy nose

### **Warnings**

#### **NIGHTTIME**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

#### **Nighttime**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- in a child who is taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product
- with any other drug containing diphenhydramine, even one used on skin.
- If your child is allergic to acetaminophen or any of the inactive ingredients in this product

- to make a child sleepy

### **Daytime**

- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression,psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

### **Ask a doctor before use if you have**

#### **Nighttime**

- liver disease
- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

#### **Daytime**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

### **Ask a doctor or pharmacist before use if you are**

#### **Nighttime**

- taking the blood thinning drug warfain
- taking sedative or tranquilizers

### **When using these products**

#### **Nighttime**

- **do not use more than directed**
- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizer may increase drowsiness

#### **Daytime**

- **do not use more than directed**

### **Stop use and ask a doctor if**

#### **Nighttime**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or last more than 5 days

- redness or swelling is present
- new symptoms occurs
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious conditions.

**Daytime**

- nervousness, dizziness or sleeplessness occur
- symptoms do not better within 7 days or occur with fever
- cough lasts more than 7 days, comes back or occur with a fever, rash, or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Nighttime**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any sign or symptoms.

**Daytime**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

**Nighttime**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- dose as follows or as directed by a doctor
- **children 6 to under 12 years of age:** 10 mL every 4 hours
- **children under 6 years of age:** do not use

**Daytime**

- do not take more than 6 doses in any 24-hours period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device
- keep dosing cup with product
- dose as follows, or as directed by a doctor
- mL = milliliter

Age	Dose
-----	------

children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

## Other information

### Nighttime

- **each 10 mL contains:** sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

### Daytime

- **each 5 mL contains:** 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

### Nighttime

citric acid, disodium EDTA, FD&C Blue #1, FD&C red #40, Flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### Daytime

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose xanthan gum

## Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

## Principal Display Panel

Compare to Mucinex® Children's Multi-Symptom Cold & Night Time Multi-Symptom Cold active ingredient†

### Children's

#### **NIGHTTIME MULTI-SYMP TOM COLD**

Acetaminophen 325 mg Pain Reliever, Fever Reducer

Diphenhydramine HCl 12.5 mg Antihistamine, Cough Suppressant

Phenhydramine HCl 5 mg Nasal Decongestant

Relieves:

Nasal and chest congestion

Reduces Fever

Soothes Cough and Sore Throat

No Added Alcohol

Berry Flavor

Ages 6-11 years

Dosage Cup Included

Ages 6 - 11 Years

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

†This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Children's Multi-Symptom Cold & Night Time Multi-Symptom Cold.

**DAYTIME**

Children's

**DAYTIME MULTI-SYMPTOM COLD**

Dextromethorphan HBr 5 mg Cough Suppressant

Guaifenesin 100 mg Expectorant

Phenylephrine HCl 2.5 mg Nasal Decongestant

Relieves: Stuffy nose, Cough, Chest congestion

Breaks up Mucus

No Added Alcohol

Berry Flavor

Dosage cup included

Ages 4 to 11 years

**FL OZ (mL)**

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

1-800-SHOP CVS

**Product Label**

**Daytime Multi-Symptom Cold**

**Drug Facts (continued)**

**Other information**

- each 5 mL contains sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl paraben, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Nighttime Multi-Symptom Cold**

**Drug Facts (continued)**

with any other drug containing diphenhydramine even one used on the skin

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product
- to make a child sleepy

**Ask a doctor before use if your child has**

- liver disease
- kidney disease
- high blood pressure
- diabetes
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if your child is**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- do not use more than directed
- irritability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- neurotosis, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- do not use if your child is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Nighttime Multi-Symptom Cold**

**Drug Facts (continued)**

**Other information**

- each 10 mL contains sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl paraben, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

**\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Children's Multi-Symptom Cold & Night Time Multi-Symptom Cold.**

**HANGER WARNING: DO NOT USE IF PRINTED SECURITY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTSIDE CHILDREN FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**PARENTS:**  
Learn about best medicine uses.  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woodstock, NJ 08899  
© 2016 CVS Pharmacy  
CVS.com® 1-800-SHOP CVS  
V-36981

100% money back guarantee  
CVS.com/returnpolicy

PAPER BOX PLASTIC BOTTLE PLASTIC CUP  
Recycled and Recyclable

**COMBO PACK**  
NDC 69842-820-08

**Children's DAYTIME MULTI-SYMPТОМ COLD**  
Dextromethorphan HBr, 5 mg  
Cough Suppressant  
Guaifenesin, 100 mg  
Expectorant  
Phenylephrine HCl, 2.5 mg  
Nasal Decongestant

**Children's NIGHTTIME MULTI-SYMPТОМ COLD**  
Acetaminophen, 325 mg  
Pain Reliever, Fever Reducer  
Diphenhydramine HCl, 12.5 mg  
Antihistamine, Cough Suppressant  
Phenylephrine HCl, 5 mg  
Nasal Decongestant

Compare to Mucinex® Children's Multi-Symptom Cold & Night Time Multi-Symptom Cold active ingredients!

Relieves: Stuffy nose, Cough, Chest congestion  
Breaks Up Mucus / No Added Alcohol

Relieves Nasal Congestion / Reduces Fever  
Soothes Cough & Sore Throat / No Added Alcohol

**TWO — 4 FL OZ (118 mL) BOTTLES**  
**TOTAL 8 FL OZ (236 mL)**

dozing cup included

#177011

PLD-DS4M  
PLD-DS5M F02894Z

Lit No.  
Exp. Date

## CVS Children's Daytime Multi-Symptom Cold Nighttime Multi-Symptom Cold Berry Flavor

### MULTI SYMPTOM COLD AND MULTI SYMPTOM NIGHTTIME COLD CHILDRENS

dextromethorphan hbr, guaifenesin, phenylephrine hci, acetaminophen, diphenhydramine hci, phenylephrine hci kit

#### Product Information

**Product Type** HUMAN OTC DRUG **Item Code (Source)** NDC:69842-820

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-820-08	1 in 1 KIT; Type 0: Not a Combination Product	02/28/2017	

#### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	118 mL
Part 2	1 BOTTLE, PLASTIC	118 mL

# MULTI SYMPTOM COLD DAYTIME CHILDRENS

dextromethorphan hbr, guaifenesin, phenylephrine hci liquid

## Product Information

**Item Code (Source)** NDC:69842-607

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE CALCIUM DISODIUM</b> (UNII: 25IH6R4SGF)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-607-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/28/2017	

## Part 2 of 2

### MULTI SYMPTOM NIGHTTIME COLD CHILDRENS

acetaminophen, diphenhydramine hci, phenylephrine hci liquid

## Product Information

Item Code (Source)	NDC:69842-606
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE CALCIUM DISODIUM</b> (UNII: 25IH6R4SGF)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-606-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/28/2017	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/28/2017	

**Labeler** - CVS Pharmacy (062312574)

Revised: 9/2025

CVS Pharmacy