

**WOMENS LAXATIVE- bisacodyl tablet, delayed release**  
**Marc Glassman, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Womens Laxative**

**Active ingredient**

Bisacodyl USP 5mg

**Purpose**

Stimulant Laxative

**Uses**

- for relief of occasional constipation and irregularity
- this product usually causes bowel movements in 6 to 12 hours

**Warnings**

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

**When using this product**

- do not chew or crush
- do not take this product within 1 hour of taking antacid or milk
- you may have stomach discomfort, faintness or cramps

**Stop use and ask a doctor if**

- rectal bleeding or failure to have a bowel movement after use of a laxative

These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

**Pregnancy/Breastfeeding**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children**

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a doctor or Poison Control Center (1-800-222-1222) right away.

## Directions

- take with water
- **Adults and children 12 years of age and older:** take 1 to 3 tablets in a single dose, once daily
- **Children 6 to 12 years of age:** take 1 tablet once daily
- **Children under 6 years of age:** consult a doctor

## Other information

- store between 68° to 77°F (20° to 25°C)
- do not use if blister unit is broken or torn
- Calcium content: 16.33mg per tablet

## Inactive ingredients

acacia, calcium sulfate anhydrous, carnauba wax, corn starch, D&C red #7 calcium lake, hypromellose phthalate, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, povidone, shellac, silicon dioxide, sodium starch glycolate, stearate acid, sugar, talc, titanium dioxide, triacetin

## Principal Display Panel

The image shows the principal display panel for Marc's Women's Laxative Bisacodyl 5 mg tablets. The panel is divided into several horizontal sections. At the top, the text "Women's Laxative Bisacodyl 5 mg" is written in pink, oriented upside down. Below this is a red rectangular logo with the word "Marc's" in white, featuring a small green leaf icon. The next section is a pink band containing the NDC number "NDC 68998-232-85" on the left and the text "Compare to the active Ingredient In Correctol®" on the right. Below the pink band is another red rectangular logo with the word "Marc's" in white and a green leaf icon. The main body of the panel is white and features the text "ENTERIC COATED" in pink, followed by "Women's Laxative" in large pink letters. Below this is a pink horizontal bar with the text "Bisacodyl 5 mg" in yellow. At the bottom, the text "For gentle, more natural relief of simple constipation" is written in pink. In the bottom right corner, there is a small illustration of a pink, oval-shaped tablet.

30 Tablets



**Women's Laxative  
Bisacodyl 5 mg**

**Drug Facts**

**Active Ingredient (in each tablet)** **Purpose**  
Bisacodyl USP 5mg.....Stimulant Laxative

**Uses** ■ for relief of occasional constipation and irregularity  
■ this product usually causes bowel movements in 6 to 12 hours

**Warnings**  
Do not use if you cannot swallow without chewing.  
**Ask a doctor before use if you have** ■ abdominal pain, nausea or vomiting  
■ a sudden change in bowel habits that persists over a period of 2 weeks  
**When using this product** ■ do not chew or crush ■ do not take this product within 1 hour of taking antacid or milk  
■ you may have stomach discomfort, faintness or cramps  
**Stop use and ask a doctor if** ■ rectal bleeding or failure to have a bowel movement after use of a laxative These could be signs of a serious condition.  
■ you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a doctor or Poison Control Center (1-800-222-1222) right away.

**Directions** ■ take with water ■ Adults and children 12 years of age and older: take 1 to 3 tablets in a single dose, once daily ■ Children 6 to 12 years of age: take 1 tablet once daily ■ Children under 6 years of age: consult a doctor

**Other information**  
■ store between 68° to 77°F (20° to 25°C) ■ do not use if blister unit is broken or torn ■ Calcium content: 16.33mg per tablet

**Inactive ingredients** acacia, calcium sulfite anhydrous, carnauba wax, corn starch, D&C red #7 calcium lake, hypromellose phthalate, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, povidone, shellac, silicon dioxide, sodium starch glycolate, stearate acid, sugar, talc, titanium dioxide, triacetin

\*This product is not manufactured or distributed by Bayer Consumer Healthcare, owner of the registered trademark Correctol®.  
LB1242 R1116

**Distributed by:** Marc Glassman, Inc.  
West 130th Street, Cleveland, OH 44130  
 0 94442 00143 5

Womens Laxative

**WOMENS LAXATIVE**

bisacodyl tablet, delayed release

**Product Information**

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68998-232
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg
--	-----------	------

### Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
KAOLIN (UNII: 24H4NWX5CO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

### Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	TCL;057
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68998-232-85	2 in 1 CARTON	07/01/2014	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	07/01/2014	

**Labeler** - Marc Glassman, Inc. (094487477)