

**EXTRA FAST COLTALIN- acetaminophen, chlorpheniramine maleate,
phenylephrine hydrochloride tablet
FORTUNE PHARMACAL COMPANY LIMITED**

DRUG FACTS

Active ingredients(in each tablet)

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Phenylephrine hydrochloride 5 mg

Purpose

Pain reliever-fever reducer

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold, the flu, or hay fever:
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sinus congestion and pressure
 - nasal congestion due to a cold or hay fever
 - runny nose
 - sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever
- temporarily reduces fever
- promotes nasal and/or sinus drainage

Warnings

Liver warning:This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 12 tablets in 24 hours, which the maximum daily amount for this product
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know whether a prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has

- heart disease
- high blood pressure
- diabetes
- glaucoma
- liver disease
- thyroid disease
- a high fever
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if the user is

- taking tranquilizers or sedatives
- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**
- may cause excitability especially in children
- use caution when driving or operating machinery
- avoid alcoholic beverages
- may cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect

Stop use and ask a doctor if

- pain or nasal congestion gets worse or lasts more than 5 days (children) or 7 days (adults)
- sore throat persists for more than 2 days
- fever gets worse or lasts more than 3 days
- nasal congestion is accompanied by fever
- nervousness, dizziness, or sleeplessness occur
- redness or swelling is present
- new symptoms occur
- any of the following occurs (these could be signs of a serious condition):
 - severe sore throat
 - sore throat is accompanied or followed by fever, headache, rash, nausea or vomiting

Do not give to children under 6 years of age unless directed by a doctor.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- adults and children 12 years of age and over: 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses (12 tablets) in 24 hours, or as directed by a doctor
- children 6 to under 12 years of age: 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses (5 tablets) in 24 hours, or as directed by a doctor
- children under 6 of age : consult a doctor

Other information

- keep tightly sealed
- store between below 25 C (77 F) in a dry place, away from direct sunlight

Inactive ingredients

colloidal silicon dioxide, corn starch, FD C yellow no. 6 aluminum lake, magnesium stearate, povidone K30, sodium starch glycolate

Questions or comments?(888) 221-3496 M-F 9 am to 5 pm

- you may also use this number to report serious adverse events associated with the use of this product

EXTRA FAST COLTALIN

Safely Sealed

NDC 51467-010-01

Acetaminophen - Pain Reliever-Fever Reducer
Chlorpheniramine maleate - Antihistamine
Phenylephrine hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief

- Minor Aches & Pain, Sore Throat
- Headache, Fever
- Nasal Congestion, Sinus Pressure/Drainage
- Sneezing, Runny Nose
- Itchy, Watery Eyes

24 Tablets

In Individually Sealed Strip

Aspirin Free



Product code

EXTRA FAST SAFELY SEALED • NDC 51467-010-01

COLTALIN®

GOLD & FLU

速效幸福傷風感冒素

Acetaminophen - Pain Reliever - Fever Reducer
Chlorpheniramine Maleate - Antihistamine
Phenylephrine Hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief
傷風、流感、花粉熱 舒緩各種症狀

- Minor Aches & Pain, Sore Throat 輕微疼痛、喉嚨痛
- Headache, Fever 發熱、頭痛
- Nasal Congestion, Sinus Pressure/Drainage 鼻塞、鼻竇充血
- Sneezing, Runny Nose 打噴嚏、流鼻涕
- Itchy, Watery Eyes 眼睛發癢、流眼淚



24
Tablets

In Individually Sealed Strip
獨立鋁箔密封包裝

ASPIRIN FREE

FORTUNE

MIX
Packaging
FSC FBCXXXX
PRINTED WITH SOY

24 Tablets/H

速效幸福傷風感冒素

Please refer to enclosed package insert for Chinese translation of drug facts.
請參閱內附之說明書以查看中文藥品資料。

Distributed by:
SOLSTICE MEDICINE COMPANY
美國蘇氏藥業有限公司
215 W. Ann Street
Los Angeles, CA 90012 U.S.A.
Tel: (888) 221-3496
Fax: (323) 221-2448
www.solsticemed.com



Pharma code

Drug Facts (continued)

• may cause drowsiness, alcohol, sedatives, and tranquilizers

Stop use and ask a doctor if

- pain or nasal congestion gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- sore throat persists for more than 2 days
- nasal congestion is accompanied by fever
- nervousness, dizziness, or sleepiness occur
- any of the following occurs (these could be signs of a serious condition):
 - drowsiness or swelling is present
 - new symptoms occur
- severe sore throat
- sore throat is accompanied or followed by fever, headache, rash, nausea or vomiting

Do not give to children under 6 years of age unless directed by a doctor.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Forming a habit may not notice any signs or symptoms.

Directions

take 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses (12 tablets) in 24 hours, or as directed by a doctor

adults and children 12 years of age and over

take 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses (5 tablets) in 24 hours, or as directed by a doctor

children 6 to under 12 years of age

take 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses (5 tablets) in 24 hours, or as directed by a doctor

consult a doctor

children under 6 years of age

Other information

- keep tightly sealed
- store below 25°C (77°F) in a dry place, away from direct sunlight

Inactive ingredients colloidal silicon dioxide, corn starch, FD&C yellow no. 6 aluminum lake, magnesium stearate, povidone K30, sodium starch glycolate

Questions or comments? (888) 221-3496 M-F 9 am to 5 pm

• you may also report serious side effects to this phone number

Drug Facts

Active Ingredients (in each tablet)

Acetaminophen 325 mg, Pain reliever- fever reducer

Chlorpheniramine maleate 2 mg, Antihistamine

Phenylephrine hydrochloride 5 mg, Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold, the flu, or hay fever:
 - sinus congestion and pressure
 - nasal congestion
 - runny nose
- sneezing, itching of the nose or throat, and itchy, watery eyes
- temporarily reduces fever
- promotes nasal and/or sinus drainage

Warnings

Liver Warnings: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 12 tablets in 24 hours, which is the maximum daily amount for this product
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug
- if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has

- heart disease
- high blood pressure
- diabetes
- glaucoma
- liver disease
- thyroid disease
- a high fever
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if the user is

- taking tranquilizers or sedatives
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- may cause excitability especially in children
- use caution when driving or operating machinery
- avoid alcoholic beverages

Manufactured by:
FORTUNE PHARMACEUTICAL CO., LTD.
香港福藥有限公司
Hong Kong
Made in China

Do not use if individually sealed foil unit dose pouches are torn, broken, or show evidence of being opened.
Keep outer carton for complete warnings and product information.
COLTALIN®
Cold & Flu
EXTRA FAST

Version no.

EXTRA FAST COLTALIN

Safely Sealed

NDC 51467-010-02

Acetaminophen - Pain Reliever-Fever Reducer
Chlorpheniramine maleate - Antihistamine
Phenylephrine hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief


- Minor Aches & Pain, Sore Throat
- Headache, Fever
- Nasal Congestion, Sinus Pressure/Drainage
- Sneezing, Runny Nose
- Itchy, Watery Eyes

36 Tablets

In Individually Sealed Strip

Aspirin Free





EXTRA FAST COLTALIN

acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51467-010
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	12mm

Flavor		Imprint Code	COLTALIN	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51467-010-01	2 in 1 BOX	04/04/2017	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:51467-010-02	3 in 1 BOX	04/04/2017	
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	04/04/2017		

Labeler - FORTUNE PHARMACAL COMPANY LIMITED (686280561)

Registrant - FORTUNE PHARMACAL COMPANY LIMITED (686280561)

Establishment			
Name	Address	ID/FEI	Business Operations
FORTUNE PHARMACAL COMPANY LIMITED		686280561	manufacture(51467-010)

Revised: 11/2025

FORTUNE PHARMACAL COMPANY LIMITED