

**AWQYYSL PET MOTION SICKNESS- maropitant citrate tablet**  
**Hong Kong Xile International Trading Limited**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**Xile, AWQYYSL Pet Motion Sickness Tablets, 16 / 24 / 60mg \*10 tablets each box**

This product is for veterinary use only. Do not administer to humans.  
Keep out of reach of children.  
Do not exceed the recommended dosage.

Prevention of Acute Vomiting: Used to prevent vomiting associated with various causes, including surgery, chemotherapy, and gastrointestinal disorders.  
Treatment of Motion Sickness: Effective in managing vomiting induced by motion sickness in dogs.  
Suitable for dogs, cats, and other pets.

Select the appropriate tablet strength and dosage based on your pet's weight: For prevention of acute vomiting, dispense whole or half tablets in strength(s) that most closely result in a 2 mg/kg dose.

**Administration Time:**

Administer the tablet 2 hour before travel for optimal effectiveness. Once daily, up to 5 days.

**Administration Method:**

Place the tablet directly into your pet's mouth.

Maropitant Citrate, 10% w/w

Lactose

Magnesium Stearate

Microcrystalline Cellulose

Mild side effects such as drowsiness or dry mouth may occur in some pets. These symptoms typically resolve on their own and do not require treatment. If severe side effects occur, stop use immediately and consult a veterinarian.

Keep out of reach of children.

Store at 15-30°C (59-86°F), protect from light.

Do not use in pets with a known allergy to any of the ingredients.



#### AWQYYSL-Maropitant Citrate Tablets Product Brochure

Thank you for choosing AWQYYSL Pet Motion Sickness Tablets! To ensure the safe and effective use of this product for your pet, please read this instruction manual carefully before use.

**Product Name:** AWQYYSL Tablets

**Active Ingredient:** [Maropitant Citrate]

**Properties:** Light orange round tablets

#### Pharmacological action:

##### 1. Inhibition of Substance P:

Maropitant citrate works by selectively inhibiting the binding of substance P to the NK1 receptors in the central nervous system (CNS). Substance P is a key neurotransmitter involved in the emetic (vomiting) reflex.

##### 2. Central and Peripheral Actions:

**Central Action:** By blocking NK1 receptors in the brainstem's vomiting center (area postrema), maropitant effectively prevents the initiation of the vomiting reflex.

**Peripheral Action:** It also acts on NK1 receptors in the gastrointestinal (GI) tract, reducing the sensitivity of the gut to emetic stimuli.

##### 3. Broad-Spectrum Antiemetic:

Maropitant citrate is effective against a wide range of emetic stimuli, including those induced by chemotherapy, toxins, and motion sickness. This makes it a versatile antiemetic agent.

##### 4. Onset and Duration:

The drug has a rapid onset of action, typically within 1-2 hours after administration, and provides sustained antiemetic effects for up to 24 hours.

##### 5. Safety Profile:

Maropitant citrate is generally well-tolerated in dogs, with a favorable safety profile. Common side effects are mild and may include pain at the injection site, lethargy, and diarrhea.

#### Clinical Applications:

**Prevention of Acute Vomiting:** Used to prevent vomiting associated with various causes, including surgery, chemotherapy, and gastrointestinal disorders.

**Treatment of Motion Sickness:** Effective in managing vomiting induced by motion sickness in dogs.

**Available Strengths:** (1) 16mg (2) 24mg (3) 60mg

**Shelf Life:** 36 months. 2 days for half a tablet opened.

**Expiration Date:** [See bottom of package]

**Storage Conditions:** Store at 15-30°C (59-86°F), protect from light

**Suitable For:** Dogs, cats, and other pets

#### Directions for Use:

##### Dosage Guidelines:

Select the appropriate tablet strength and dosage based on your pet's weight: For prevention of acute vomiting, dispense whole or half tablets in strength(s) that most closely result in a 2 mg/kg dose:

Dog body weight		Number of Tablets		
Pounds	Kilograms	16mg	24mg	60mg
8	4	1/2		
15	8	1		
25	12		1	
50	24		2	
65	30			1
130	60			2

##### Administration Time:

Administer the tablet 2 hour before travel for optimal effectiveness. once daily, up to 5 days.

**Administration Method:** Place the tablet directly into your pet's mouth

##### Precautions:

This product is for veterinary use only. Do not administer to humans. Keep out of reach of children. Do not exceed the recommended dosage.

If your pet experiences any adverse reactions, discontinue use immediately and consult a veterinarian. This product is not a substitute for professional veterinary care. If motion sickness symptoms persist or worsen, seek veterinary advice. Store in a cool, dry place away from direct sunlight.

##### Contraindications:

Do not use in pets with a known allergy to any of the ingredients.

##### Possible Side Effects:

Mild side effects such as drowsiness or dry mouth may occur in some pets. These symptoms typically resolve on their own and do not require treatment. If severe side effects occur, stop use immediately and consult a veterinarian.

##### Disclaimer:

This instruction manual is for reference only. For specific dosage and usage guidance, consult your veterinarian.

**FDA Registered with NDC#**

##### Manufacturer Name:

Hong Kong Xile International Trading Limited

**Address:** 16 Shing Yip Street Kwun Tong, Hong Kong

**Email:** zhangpanqiye@163.com

**AWQYYSL** Caring for Your Pet's Well-Being, Every Step of the Way."



## Drug Information

**Active Ingredient:** (Per chewable tablet) Maropitant Citrate, USP 16mg

**Purpose:**

AWQYSL tablets are used to prevent and alleviate symptoms of motion sickness in animals caused by travel in cars, boats, airplanes, or other modes of transportation, such as nausea, vomiting, dizziness, and other related symptoms.

**Directions for Use:**

Administer at least two hours before travel to prevent vomiting caused by an empty stomach; however, do not feed a full meal before travel.

For animals 8 weeks of age and older.

Administer once daily at a dose of 2.0 mg per kilogram of body weight, for up to 5 consecutive days.

To prevent acute vomiting, a full or half tablet may be given, not exceeding two doses per day.

**Warnings:**

If your dog has any chronic medical conditions, consult a veterinarian before use.

Consult a veterinarian or pharmacist if your dog is currently taking any other medications.

Not suitable for pregnant animals or those with liver dysfunction.

Do not wrap the product in food for administration, as this may delay dissolution and affect efficacy.

Not for human use. Keep out of reach of children. In case of accidental ingestion, seek medical advice.

**Additional Information:** Store at room temperature between 59°F-86°F (15°C-30°C).

Protect from moisture. Check the expiration date and batch number on the seal.

**Inactive Ingredients:** Lactose, Magnesium Stearate, Microcrystalline Cellulose.

**Manufacturer Information:**

**Manufacturer Name:** Hong Kong Xile International Trading Limited

**Email:** zhangpanqiye@163.com

**Address:** 16 Shing Yip Street Kwun Tong, Hong Kong

See package insert for complete dosage information and complete product information.



AWQYSL

10 TABLETS (16.0mg each)



Alleviating motion sickness in animals  
Travel anxiety

**No side effect**

- Vomiting
- Drooling
- Shivering
- Pacemaking
- Restlessness or anxiety
- Panning

AWQYSL  
Maropitant Citrate Tablets

#1  
RECOMMENDED  
BY VETERINARIANS  
BRAND

For oral use in animals only  
Keep this and all drugs out of the reach of children

AWQYSL  
Maropitant Citrate Tablets



## Drug Information

**Active Ingredient:** (Per chewable tablet) Maropitant Citrate, USP 24mg

**Purpose:**

AWQYSL tablets are used to prevent and alleviate symptoms of motion sickness in animals caused by travel in cars, boats, airplanes, or other modes of transportation, such as nausea, vomiting, dizziness, and other related symptoms.

**Directions for Use:**

Administer at least two hours before travel to prevent vomiting caused by an empty stomach; however, do not feed a full meal before travel.

For animals 8 weeks of age and older.

Administer once daily at a dose of 2.0 mg per kilogram of body weight, for up to 5 consecutive days.

To prevent acute vomiting, a full or half tablet may be given, not exceeding two doses per day.

**Warnings:**

If your dog has any chronic medical conditions, consult a veterinarian before use.

Consult a veterinarian or pharmacist if your dog is currently taking any other medications.

Not suitable for pregnant animals or those with liver dysfunction.

Do not wrap the product in food for administration, as this may delay dissolution and affect efficacy.

Not for human use. Keep out of reach of children. In case of accidental ingestion, seek medical advice.

**Additional Information:** Store at room temperature between 59°F-86°F (15°C-30°C).

Protect from moisture. Check the expiration date and batch number on the seal.

**Inactive Ingredients:** Lactose, Magnesium Stearate, Microcrystalline Cellulose.

**Manufacturer Information:**

**Manufacturer Name:** Hong Kong Xile International Trading Limited

**Email:** zhangpanqiye@163.com

**Address:** 16 Shing Yip Street Kwun Tong, Hong Kong

See package insert for complete dosage information and complete product information.



AWQYSL

10 TABLETS (24.0mg each)



Alleviating motion sickness in animals  
Travel anxiety

- Vomiting
- Drooling
- Shivering
- Pacing
- Restlessness or anxiety
- Panning

**No side effect**

Maropitant Citrate Tablets

AWQYSL

#1 RECOMMENDED  
BY VETERINARIANS  
BRAND

For oral use in animals only  
Keep this and all drugs out of the reach of children

Maropitant Citrate Tablets

AWQYSL

## Drug Information

**Active Ingredient:** (Per chewable tablet) Maropitant Citrate, USP 60mg

**Purpose:**

AWQYSL tablets are used to prevent and alleviate symptoms of motion sickness in animals caused by travel in cars, boats, airplanes, or other modes of transportation, such as nausea, vomiting, dizziness, and other related symptoms.

**Directions for Use:**

Administer at least two hours before travel to prevent vomiting caused by an empty stomach; however, do not feed a full meal before travel.

For animals 8 weeks of age and older.

Administer once daily at a dose of 2.0 mg per kilogram of body weight, for up to 5 consecutive days.

To prevent acute vomiting, a full or half tablet may be given, not exceeding two doses per day.

**Warnings:**

If your dog has any chronic medical conditions, consult a veterinarian before use.

Consult a veterinarian or pharmacist if your dog is currently taking any other medications.

Not suitable for pregnant animals or those with liver dysfunction.

Do not wrap the product in food for administration, as this may delay dissolution and affect efficacy.

Not for human use. Keep out of reach of children. In case of accidental ingestion, seek medical advice.

**Additional Information:** Store at room temperature between 59°F-86°F (15°C-30°C).

Protect from moisture. Check the expiration date and batch number on the seal.

**Inactive Ingredients:** Lactose, Magnesium Stearate, Microcrystalline Cellulose.

**Manufacturer Information:**

**Manufacturer Name:** Hong Kong Xile International Trading Limited

**Email:** zhangpanqiye@163.com

**Address:** 16 Shing Yip Street Kwun Tong, Hong Kong

See package insert for complete dosage information and complete product information.



AWQYSL

10 TABLETS (60.0mg each)



Alleviating motion sickness in animals  
Travel anxiety

**No side effect**

- Vomiting
- Drooling
- Shivering
- Pacemaking
- Restlessness or anxiety
- Panning

AWQYSL  
Maropitant Citrate Tablets

#1  
RECOMMENDED  
BY VETERINARIANS  
BRAND

**For oral use in animals only**  
Keep this and all drugs out of the reach of children

AWQYSL  
Maropitant Citrate Tablets

## AWQYSL PET MOTION SICKNESS

maropitant citrate tablet

### Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:85209-101
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>Maropitant Citrate</b> (UNII: LXN6S3999X) (MAROPITANT - UNII:4XE2T9H4DH)	MAROPITANT	10 mg in 100 mg

Inactive Ingredients	
Ingredient Name	Strength
<b>Lactose</b> (UNII: J2B2A4N98G)	
<b>Magnesium stearate</b> (UNII: 70097M6I30)	
<b>Microcrystalline Cellulose</b> (UNII: OP1R32D61U)	

Product Characteristics			
<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	5mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85209-101-21	1600 mg in 1 BAG		
2	NDC:85209-101-31	2400 mg in 1 BAG		
3	NDC:85209-101-71	6000 mg in 1 BAG		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/26/2025	

**Labeler** - Hong Kong Xile International Trading Limited (765202372)

**Registrant** - Hong Kong Xile International Trading Limited (765202372)

Establishment			
Name	Address	ID/FEI	Business Operations
Hong Kong Xile International Trading Limited		765202372	label, manufacture, api manufacture

Revised: 6/2025

Hong Kong Xile International Trading Limited