

PROHEART 6- moxidectin
Zoetis Inc.

ProHeart® 6 (moxidectin)
For Extended-Release Injectable Suspension for Dogs

CAUTION

Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

ProHeart 6 (moxidectin) for extended-release injectable suspension consists of two separate vials: One vial contains 10% moxidectin sterile microspheres and the second vial contains a specifically formulated sterile vehicle for constitution with the microspheres. No other diluent should be used. A clear or translucent appearance of the vehicle is normal. Each mL of constituted drug product contains 3.4 mg moxidectin, 3.1% glyceryl tristearate, 2.4% hydroxypropyl methylcellulose, 0.87% sodium chloride, 0.17% methylparaben, 0.02% propylparaben and 0.001% butylated hydroxytoluene. Hydrochloric acid is used to adjust pH. The constituted product may appear as a hazy to milky suspension.

INDICATIONS

ProHeart 6 is indicated for use in dogs six months of age and older for the prevention of heartworm disease caused by *Dirofilaria immitis* for 6 months.
ProHeart 6 is indicated for the treatment of existing larval and adult hookworm (*Ancylostoma caninum* and *Uncinaria stenocephala*) infections.

DOSAGE AND ADMINISTRATION

Always provide Client Information Sheet and review with owners before administering ProHeart 6. The owner should be advised to observe their dog for adverse drug events including those described on the sheet.

Frequency of Treatment

ProHeart 6 prevents the development of heartworm disease caused by *D. immitis* for six months. It should be administered within one month of the dog's first exposure to mosquitoes. Follow-up treatments may be given every six months if the dog has continued exposure to mosquitoes and if the dog continues to be healthy without weight loss. When replacing another heartworm preventive product, ProHeart 6 should be given within one month of the last dose of the former medication.

ProHeart 6 eliminates the larval and adult stages of *A. caninum* and *U. stenocephala* present at the time of treatment. However, persistent effectiveness has not been

established for this indication. Re-infection with *A. caninum* and *U. stenocephala* may occur sooner than 6 months.

Dose

The recommended subcutaneous dose is 0.05 mL of the constituted suspension/kg body weight (0.0227 mL/lb.). This amount of suspension will provide 0.17 mg moxidectin/ kg bodyweight (0.0773 mg/lb.).

To ensure accurate dosing, calculate each dose based on the dog's weight at the time of treatment. Do not overdose growing puppies in anticipation of their expected adult weight. The following dosage chart may be used as a guide.

DOSAGE CHART

Dog Wt. Dose Volume			Dog Wt. Dose Volume		
lb	kg	mL/Dog	lb	kg	mL/Dog
11	5	0.25	77	35	1.75
22	10	0.50	88	40	2.00
33	15	0.75	99	45	2.25
44	20	1.00	110	50	2.50
55	25	1.25	121	55	2.75
66	30	1.50	132	60	3.00

Injection Technique

The two-part sustained release product must be mixed at least 30 minutes prior to the intended time of use (see **CONSTITUTION PROCEDURES** for initial mixing instructions). Once constituted, **swirl the bottle gently before every use to uniformly re-suspend the microspheres**. Withdraw 0.05 mL of suspension/kg body weight into an appropriately sized syringe fitted with an 18G or 20G hypodermic needle. Dose promptly after drawing into dosing syringe. If administration is delayed, gently roll the dosing syringe prior to injection to maintain a uniform suspension and accurate dosing.

Using aseptic technique, inject the product subcutaneously in the left or right side of the dorsum of the neck cranial to the scapula. No more than 3 mL should be administered in a single site. The location(s) of each injection (left or right side) should be noted so that prior injection sites can be identified and the next injection can be administered on the opposite side.

RISK MINIMIZATION ACTION PLAN

The ProHeart 6 and ProHeart 12 Risk Minimization Action Plan (RiskMAP) provides educational materials to the veterinarian, veterinary staff, and the dog owner explaining the risks and proper use of ProHeart 6 and ProHeart 12. ProHeart 6 and ProHeart 12 are the same formulation, but ProHeart 6 is three times less concentrated than ProHeart 12. ProHeart 6 and ProHeart 12 are for use in dogs only and are available through a restricted distribution program to veterinarians that have completed the RiskMAP training and certification module.

The ProHeart 6 and ProHeart 12 web-based training and certification module is available at <http://www.proheart6.com>. This website has important information on the safe and effective use of ProHeart 6 and ProHeart 12 for veterinarians.

Only veterinarians and veterinary technicians/assistants that have completed the training and are certified can administer ProHeart 6 and ProHeart 12. Veterinarians are expected to report all adverse events that occur in animals or humans to the manufacturer. Important safety information is included below:

CONTRAINDICATIONS

ProHeart 6 is contraindicated in animals previously found to be hypersensitive to this drug or ProHeart 12.

HUMAN WARNINGS

Not for human use. Keep this and all drugs out of the reach of children.

May be slightly irritating to the eyes. May cause slight irritation to the upper respiratory tract if inhaled. May be harmful if swallowed. If contact with the eyes occurs, rinse thoroughly with water for 15 minutes and seek medical attention immediately. If accidental ingestion occurs, contact a Poison Control Center or a physician immediately. The Safety Data Sheet (SDS) contains more detailed occupational safety information.

WARNINGS

Anaphylactic and anaphylactoid reactions may occur in some dogs following administration of ProHeart 6 alone or with vaccines. In some cases, these reactions have resulted in death following administration of moxidectin microspheres (see POST-APPROVAL EXPERIENCE). Anaphylactic and anaphylactoid reactions should be treated immediately with the same measures used to treat hypersensitivity reactions to vaccines and other injectable products.

Always provide Client Information Sheet and review with owners before administering ProHeart 6. The owner should be advised to observe their dog for adverse drug events including those described on the sheet.

Do not administer ProHeart 6 to dogs who are sick, debilitated, underweight or who have a history of weight loss.

PRECAUTIONS

Prior to administration of ProHeart 6, the health of the patient should be assessed by a thorough medical history, physical examination and diagnostic testing as indicated (see **WARNINGS**).

Caution should be used when administering ProHeart 6 in dogs with pre-existing allergic

disease, including food allergy, atopy, and flea allergy dermatitis. (see **WARNINGS**). Caution should be used when administering ProHeart 6 concurrently with vaccinations. Adverse reactions, including anaphylaxis, have been reported following the concomitant use of moxidectin microspheres and vaccinations (see **WARNINGS** and **POST-APPROVAL EXPERIENCE**).

ProHeart 6 should not be used more frequently than every 6 months.

The effectiveness of ProHeart 6 has not been evaluated in dogs less than 6 months of age.

Prior to administration of ProHeart 6, dogs should be tested for existing heartworm infections. Infected dogs should be treated with an adulticide to remove adult heartworms. ProHeart 6 is not effective against adult *D. immitis*.

Caution should be used when administering ProHeart 6 to heartworm positive dogs (see **ADVERSE REACTIONS**).

ADVERSE REACTIONS

In field studies, the following adverse reactions were observed in dogs treated with ProHeart 6: anaphylaxis, vomiting, diarrhea (with and without blood), listlessness, weight loss, seizures, injection site pruritus, and elevated body temperature. Dogs with clinically significant weight loss (>10%) were more likely to experience a severe adverse reaction.

In a laboratory effectiveness study, dogs with 4- and 6-month-old heartworm infections experienced vomiting, lethargy and bloody diarrhea. These signs were more severe in the dogs with 4-month-old heartworm infections, including one dog that was recumbent and required supportive care, than in the dogs with older (6-month-old) infections.

Post-Approval Experience (Rev. 2018)

The following adverse events are based on post-approval adverse drug experience reporting. Not all adverse reactions are reported to FDA/CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using these data. The following adverse events are listed in decreasing order of frequency by body system.

Immune: anaphylaxis and/or anaphylactoid reactions, urticaria, head/facial edema, pruritus, pale mucous membranes, collapse, cardiovascular shock, erythema, immune-mediated hemolytic anemia, immune-mediated thrombocytopenia (signs reflected in other system categories could be related to allergic reactions, i.e. gastrointestinal, dermatologic, and hematologic)

Gastrointestinal: vomiting (with or without blood), diarrhea with or without blood, hypersalivation

General: depression, lethargy, anorexia, fever, weight loss, weakness

Dermatological: injection site pruritus/swelling, erythema multiforme

Neurological: seizures, ataxia, trembling, hind limb paresis

Hematological: leukocytosis, anemia, thrombocytopenia

Respiratory: dyspnea, tachypnea, coughing

Hepatic: elevated liver enzymes, hypoproteinemia, hyperbilirubinemia, hepatopathy

Urinary: elevated BUN, elevated creatinine, hematuria, polydipsia, polyuria

Cardiopulmonary signs such as coughing and dyspnea may occur in heartworm positive dogs.

In some cases, death has been reported as an outcome of the adverse events listed above.

For a copy of the Safety Data Sheet (SDS) or to report suspected adverse reactions, contact Zoetis at 1-888-963-8471. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or www.fda.gov/reportanimalae.

INFORMATION FOR DOG OWNERS

Always provide Client Information Sheet and review with owners before administering ProHeart 6. Owners should be advised of the potential for adverse reactions, including anaphylaxis, and be informed of the clinical signs associated with drug toxicity (see **WARNINGS, ADVERSE REACTIONS** and **POST-APPROVAL EXPERIENCE** sections.)

Owners should be advised to contact their veterinarian immediately if signs of toxicity are observed. The vast majority of patients with drug related adverse reactions have recovered when the signs are recognized and veterinary care, if appropriate, is initiated.

CLINICAL PHARMACOLOGY

Moxidectin is a semi-synthetic methoxime derivative of nemadectin which is a fermentation product of *Streptomyces cyaneogriseus* subspecies *noncyanogenus*. Moxidectin is a pentacyclic 16-membered lactone macrolide.

Moxidectin has activity resulting in paralysis and death of affected parasites. The stage of the canine heartworm affected at the recommended dose rate of 0.5 mg/kg (0.227 mg/lb) is the tissue larval stage.

The larval and adult stages of the canine hookworms, *A. caninum* and *U. stenocephala*, are susceptible.

Following injection with ProHeart 6, peak moxidectin blood levels will be observed approximately 7-14 days after treatment. At the end of the 6-month dosing interval, residual drug plasma concentrations are negligible. Accordingly, little or no drug accumulation is expected to occur with repeated administrations.

ANIMAL SAFETY

General Safety: ProHeart 6 has been administered to a wide variety of healthy dogs six months of age and older, including a wide variety of breeds, pregnant and lactating females, breeding males, and ivermectin-sensitive collies. In clinical studies, two geriatric dogs with a history of weight loss after the initial ProHeart 6 injection died within a month of the second 6 month injection. A third dog who was underweight for its age and breed

and who had a history of congenital problems experienced lethargy following the initial injection of ProHeart 6. The dog never recovered and died 3 months later (see **WARNINGS**).

ProHeart 6 administered at 3 times the recommended dose in dogs with patent heartworm infections and up to 5 times the recommended dose in ivermectin-sensitive collies did not cause any adverse reactions. ProHeart 6 administered at 3 times the recommended dose did not adversely affect the reproductive performance of male or female dogs.

ProHeart 6 administered up to 5 times the recommended dose in 7-8 month old puppies did not cause any systemic adverse effects.

In well controlled clinical field studies, ProHeart 6 was used in conjunction with a variety of veterinary products including anthelmintics, antiparasitics, antibiotics, analgesics, steroids, non-steroidal anti-inflammatory drugs (NSAIDs), anesthetics and flea control products.

Injection Site Reactions: Injection site observations were recorded during effectiveness and safety studies. In clinical studies, ProHeart 6 was administered at six-month intervals to client-owned dogs under field conditions. There were no reports of injection site reactions in these field studies and evaluations of the injection sites revealed no abnormalities.

In a laboratory safety study, ProHeart 6 was administered at 1, 3 and 5 times the recommended dose to 7-8 month old puppies. Injection sites were clipped to facilitate observation. Slight swelling/edema at the injection site was observed in some dogs from all treated groups. These injection site reactions appeared as quickly as 8 hours post injection and lasted up to 3 weeks. A three-year repeated injection study was conducted to evaluate the safety of up to 6 injections of ProHeart 6 administered at the recommended dose (0.17 mg/kg) every 6 months. Mild erythema and localized deep subcuticular thickening were seen in dogs that received four injections in the same area on the neck and in one dog that received two injections in the same area on the neck. Microscopic evaluation on the injection sites from all dogs 6 months after the last injection consistently showed mild granulomatous panniculitis with microvacuolation. The only adverse reaction seen that was not related to the injection site was weight loss in one dog.

Some dogs treated with ProHeart 6 in laboratory effectiveness studies developed transient, localized inflammatory injection site reactions. These injection site reactions were visible grossly for up to 3 weeks after injection. Histologically, well-defined granulomas were observed in some dogs at approximately 5 months after injection.

CONSTITUTION PROCEDURES

The two-part ProHeart 6 product must be mixed at least 30 minutes prior to the intended time of use.

Items needed to constitute ProHeart 6:	<ul style="list-style-type: none">• Microspheres• Sterile 20 mL syringe for transfer	<ul style="list-style-type: none">• Enclosed vent needle (25G)• Transfer needle (18G)	<ul style="list-style-type: none">• Sterile Vehicle
--	---	--	---

or 20G)

Constitution of the 20 mL vial product (17.7 mL when constituted).



1. Shake the microsphere vial to break up any aggregates prior to constitution.
 2. Using an 18G or 20G needle and sterile syringe withdraw 17.0 mL of the unique sterile vehicle from the vial. There is more sterile vehicle supplied than the 17.0 mL required.
 3. Insert the enclosed 25G vent needle into the microsphere vial.
 4. Slowly transfer the sterile vehicle into the microsphere vial through the stopper using the transfer needle and syringe.
 5. Once the sterile vehicle has been added, remove the vent and transfer needles from the microsphere vial. Discard unused sterile vehicle and needles.
 6. Shake the microsphere vial vigorously until a thoroughly mixed suspension is produced. The constituted product may appear as a hazy to milky suspension.
 7. Record the time and date of mixing on the microsphere vial.
 8. Allow suspension to stand for at least 30 minutes to allow large air bubbles to dissipate.
 9. **Before every use, gently swirl the mixture to achieve uniform suspension.** The constituted product may appear as a hazy to milky suspension. The microspheres and vehicle will gradually separate on standing.
 10. Use a 1 mL or 3 mL syringe and an 18G or 20G needle for dosing. Dose promptly after drawing into dosing syringe. If administration is delayed, gently roll the dosing syringe prior to injection to maintain a uniform suspension and accurate dosing.
 11. Refrigerate the unused product. The constituted product remains stable for 8 weeks in a refrigerator. Avoid direct sunlight.
-

STORAGE INFORMATION

Store the unconstituted product at or below 25°C (77°F). Do not expose to light for extended periods of time. After constitution, the product is stable for 8 weeks stored under refrigeration at 2° to 8°C (36° to 46°F).

HOW SUPPLIED

ProHeart 6 is available in the following three package sizes.

1. 1-Pack

20 mL vial product:
1 - 10% moxidectin
sterile microspheres
- 598 mg/vial
1 - Sterile vehicle - 17

2. 5-Pack

20 mL vial product:
5 - 10% moxidectin
sterile microspheres
- 598 mg/vial
5 - Sterile vehicle - 17

3. 10-Pack

20 mL vial product:
10 - 10% moxidectin
sterile microspheres
- 598 mg/vial
10 - Sterile vehicle - 17

Client Information Sheet

Client Information Sheet

Review this information with your veterinarian each time your dog receives ProHeart 6. When your veterinarian prescribes medicine for your dog, it's sometimes hard to remember all the information discussed during the visit. This sheet is provided as a summary, but if you have any additional questions you should speak with your veterinarian.

ProHeart® 6
(moxidectin)

What is ProHeart 6?



ProHeart 6 is a medication that prevents heartworm disease in dogs 6 months of age and older. Your veterinarian gives it to your dog via an injection and it protects your dog from developing heartworm disease for 6 full months. ProHeart 6 also treats common hookworm infections your dog may have at the time of injection.

What should I discuss with my veterinarian before using ProHeart 6?



- **General health** - ProHeart 6 should only be given to healthy dogs.
- **Changes in behavior or health**, including weight loss.
- **Allergies** - Past or present, uncontrolled allergies, including food, flea or skin allergies.
- **Past problems with or reactions to vaccines or medications.**
- **Current medications, supplements or special diets**, including those you can get without a prescription.

What is heartworm disease and how can my dog get it?



Heartworm disease is a serious, potentially fatal illness where worms grow in a dog's heart and lungs. Mosquitoes spread heartworm disease when they bite a dog. A single bite from an infected mosquito can put your dog at risk for developing heartworm disease.

What are hookworms and how could my dog get them?



Hookworms are common parasites that can live in a dog's intestines. Your dog can get hookworms by eating hookworm larvae (young hookworms) that may be in the dirt or they can get hookworms through skin contact.

What possible ProHeart 6 side effects could happen to my dog?



It is important to contact your veterinarian if you observe *any* signs of illness in your dog. Severe reactions require emergency treatment by your veterinarian. Watch your dog for the following possible signs of illness:

- **Allergic reactions** - Allergic symptoms such as swelling of the face, itching, hives and/or inflamed skin. Allergic reactions may occur when ProHeart 6 is given alone or with vaccines. Some allergic reactions can be severe, such as difficulty breathing or collapse.
- **Vomiting and/or diarrhea** - Either with or without blood.
- **Seizures**
- **Change in your dog's appetite or activity level**

Most reactions occur within the first 24 hours of receiving ProHeart 6; severe allergic reactions may occur in the first two hours.

IF YOU NOTICE ANY SIGNS OF ILLNESS, OR ANYTHING OUT OF THE ORDINARY AFTER YOUR DOG RECEIVES PROHEART 6, CONTACT YOUR VETERINARIAN IMMEDIATELY.

In some cases, these events may be serious and may cause death.

If you have any questions about ProHeart 6, talk to your veterinarian.

[Veterinarian: Please place contact information below]

Practice Name: _____

Vet Name: _____

Phone: _____

To obtain additional information, visit ProHeart6.com or call 1-888-963-8471.
Approved by FDA under NADA # 141-189

Distributed by: Zoetis Inc. Kalamazoo, MI 49007

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted.

© 2020 Zoetis Services LLC. All rights reserved. PRO-00309

zoetis

zoetis


Distributed by Zoetis Inc., Kalamazoo, MI 49007

Approved by FDA under NADA # 141-189


Revised: November 2021

40036951

PRINCIPAL DISPLAY PANEL - 1 Vial Kit Carton




PRINCIPAL DISPLAY PANEL - 5 Vial Kit Carton



PRINCIPAL DISPLAY PANEL - 10 Vial Kit Carton

For Use In Dogs

10 Vials of Moxidectin Sterile Microspheres 598 mg
10 Vials of Sterile Vehicle 17 mL
10 Vent Needles



PROHEART 6

moxidectin kit

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-3670
---------------------	--------------------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-3670-1	1 in 1 CARTON		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	10 VIAL, MULTI-DOSE	10 mL
Part 2	1 VIAL, MULTI-DOSE	17 mL

Part 1 of 2

PROHEART

moxidectin injection

Product Information

Route of Administration	SUBCUTANEOUS
--------------------------------	--------------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MOXIDECTIN (UNII: NGU5H31YO9) (MOXIDECTIN - UNII:NGU5H31YO9)	MOXIDECTIN	3.4 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
TRISTEARIN (UNII: P6OCJ2551R)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

Part 2 of 2

STERILE VEHICLE

inert injection

Product Information

Route of Administration SUBCUTANEOUS

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		17 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

PROHEART 6

moxidectin kit

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-3671
--------------	--------------------------	--------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-3671-1	1 in 1 CARTON		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	5 VIAL, MULTI-DOSE	88 mL
Part 2	5 VIAL, MULTI-DOSE	85 mL

Part 1 of 2

PROHEART

moxidectin injection

Product Information

Route of Administration	SUBCUTANEOUS
-------------------------	--------------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MOXIDECTIN (UNII: NGU5H31YO9) (MOXIDECTIN - UNII:NGU5H31YO9)	MOXIDECTIN	3.4 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
TRISTEARIN (UNII: P6OCJ2551R)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		17.6 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

Part 2 of 2**STERILE VEHICLE**

inert injection

Product Information**Route of Administration** SUBCUTANEOUS**Inactive Ingredients**

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		17 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

PROHEART 6

moxidectin kit

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-3672
---------------------	--------------------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-3672-1	1 in 1 CARTON		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	10 VIAL, MULTI-DOSE	176 mL
Part 2	10 VIAL, MULTI-DOSE	170 mL

Part 1 of 2

PROHEART

moxidectin injection

Product Information

Route of Administration	SUBCUTANEOUS
--------------------------------	--------------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MOXIDECTIN (UNII: NGU5H31YO9) (MOXIDECTIN - UNII:NGU5H31YO9)	MOXIDECTIN	3.4 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
TRISTEARIN (UNII: P6OCJ2551R)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		17.6 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
NADA	NADA141189	06/06/2001	

Part 2 of 2

STERILE VEHICLE

inert injection

Product Information

Route of Administration SUBCUTANEOUS

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		17 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141189	06/06/2001	

Labeler - Zoetis Inc. (828851555)