

**PAIN RELIEF- acetaminophen tablet**  
**Meijer Distribution Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Meijer Distribution, Inc. Pain Relief Drug Facts**

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 caplets every 6 hours while symptoms last</li><li>• do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

carnauba wax, corn starch\*, croscarmellose sodium\*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid

\*may contain one or more of these ingredients

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Compare to Extra Strength Tylenol<sup>®</sup> active ingredient

EXTRA STRENGTH

pain relief 500 mg Each

Acetaminophen | Pain Reliever | Fever Reducer | For Adults

100 Caplets

actual size

meijer

NDC 41250-484-78

Compare to ExtraStrength Tylenol® active ingredient\*\*

EXTRA STRENGTH

pain relief 500 mg Each

Acetaminophen | Pain Reliever | Fever Reducer | For Adults

100 Caplets



actual size

meijer

NDC 41250-484-78

Compare to ExtraStrength Tylenol® active ingredient\*\*

EXTRA STRENGTH

pain relief 500 mg Each

Acetaminophen | Pain Reliever | Fever Reducer | For Adults

100 Caplets



actual size

DO NOT USE, IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com



**Drug Facts**

**Active ingredient (in each caplet)** **Purpose**

Acetaminophen 500 mg.....Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - minor pain of arthritis
  - backache
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**User warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** if with any other drug containing acetaminophen (read label or compare tablets). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition.

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

do not take more than directed (see overdose warning)

- adults and children 12 years and over
  - take 2 caplets every 6 hours while symptoms last
  - do not take more than 6 caplets in 24 hours, unless directed by a doctor
  - do not use for more than 10 days unless directed by a doctor
- children under 12 years
  - ask a doctor

**Other information**

store at 21-25°C (68-77°F)

**Inactive ingredients** carmelumina wax, corn starch\*, or croscarmellose sodium\*, hypromellose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, sodium starch glycolate, stearic acid. \*may contain one or more of these ingredients

**Questions or comments?** 1-800-719-9280

\*\*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Tylenol®.



48478 6E C12

## PAIN RELIEF

acetaminophen tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-484
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

### Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	L484
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-484-71	1 in 1 CARTON	08/15/1987	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-484-78	1 in 1 CARTON	10/28/1992	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-484-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/13/1992	

4	NDC:41250-484-76	1 in 1 CARTON	03/20/1992	
4		120 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:41250-484-02	550 in 1 BOTTLE; Type 0: Not a Combination Product	06/28/2013	04/06/2016
6	NDC:41250-484-82	2 in 1 PACKAGE	10/28/1992	
6		1 in 1 CARTON		
6		100 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:41250-484-62	1 in 1 CARTON	06/16/2017	
7		24 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/15/1987	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 1/2019

Meijer Distribution Inc