

PAIN RELIEF- acetaminophen tablet
Meijer Distribution Inc

Meijer Distribution, Inc. Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain one or more of these ingredients

Questions or comments?

Principal Display Panel

meijer®

EXTRA STRENGTH

Pain Relief

COMPARE TO EXTRA STRENGTH TYLENOL® ACTIVE INGREDIENT



ACETAMINOPHEN

PAIN RELIEVER | FEVER REDUCER | FOR ADULTS

Actual Size

24 CAPLETS

500 mg each



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

OUR QUALITY GUARANTEE

My Meijer

www.meijer.com/ satisfaction

www2recycle.info

PAPER BOX
PLASTIC BOTTLE

WWW.MEIJER.COM

DIST. BY MEIJER DISTRIBUTION, N.C.
GRAND RAPIDS, MI 49544

This product is not manufactured or distributed by
Kenvue Inc., owner of the registered trademark Tylenol®.

meijer.


EXTRA STRENGTH

Pain Relief

COMPARE TO EXTRA STRENGTH TYLENOL® ACTIVE INGREDIENT**


NDC 41250-484-62

ACETAMINOPHEN
PAIN RELIEVER | FEVER REDUCER | FOR ADULTS



Actual Size

24
CAPLETS
500 mg each



Drug Facts

Active ingredient (in each caplet)
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

ask a doctor

children under 12 years


Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium*; hypromellose, croscarmellose sodium*; hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients

Questions or comments?
1-800-719-9280



7 08820 83345 6

PID 367268

48462 6E C6

PAIN RELIEF

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-484
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	L484
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-484-71	1 in 1 CARTON	08/15/1987	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-484-78	1 in 1 CARTON	10/28/1992	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-484-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/13/1992	
4	NDC:41250-484-76	1 in 1 CARTON	03/20/1992	06/23/2018
4		120 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:41250-484-02	550 in 1 BOTTLE; Type 0: Not a Combination Product	06/28/2013	04/06/2016
6	NDC:41250-484-82	2 in 1 PACKAGE	10/28/1992	
6		1 in 1 CARTON		
6		100 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:41250-484-62	1 in 1 CARTON	06/16/2017	
7		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/15/1987	

Labeler - Meijer Distribution Inc (006959555)

Revised: 1/2026

Meijer Distribution Inc