

**DRAMAMINE- dimenhydrinate tablet**  
**Select Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Dramamine®**

**Drug Facts**

**Active ingredient (in each tablet)**

Dimenhydrinate 50 mg

**Purpose**

Antiemetic

**Use**

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

**Warnings**

**Do not give to** children under 2 years of age unless directed by a doctor

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking** sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,** ask a doctor before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) immediately.

### Directions

- to prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 1 to 2 chewable tablets every 4-6 hours</li><li>• do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor</li></ul>
children 6 to under 12 years	<ul style="list-style-type: none"><li>• give 1/2 to 1 chewable tablet every 6-8 hours</li><li>• do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor</li></ul>
children 2 to under 6 years	<ul style="list-style-type: none"><li>• give 1/2 chewable tablet every 6-8 hours</li><li>• do not give more than 1½ chewable tablets in 24 hours, or as directed by a doctor</li></ul>

### Other information

- **Phenylketonurics:** contains phenylalanine 0.84 mg per tablet
- store at room temperature 20°- 25°C (68°-77°F)
- **do not use if pouch is opened**
- see bottom of this panel for lot number and expiration date

### Inactive ingredients

anhydrous citric acid, aspartame, FD&C yellow #6 aluminum lake, flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified starch, sorbitol

### Questions or comments?

call **1-800-382-7219**

Dist By:

Medtech Products, Inc. Tarrytown, NY 10591

Product Repackaged and Distributed

with permission of manufacturer by:

**Select Corporation** Carrollton, TX 75007

**PRINCIPAL DISPLAY PANEL - 50 mg Tablet Packet Carton**

#1

Pharmacist

Recommended

BRAND

DIMENHYDRINATE  
TABLETS / ANTIEMETIC

Dramamine®

motion sickness

CHEWABLE

Dual Action:

Prevents & Relieves Nausea,  
Dizziness and Vomiting

TO OPEN

PUSH IN TAB AND PULL OUT

25 Packets of 2 Orange Flavor Tablets  
(50 mg each)

DIMENHYDRINATE  
TABLETS / ANTIEMETIC

#1  
Pharmacist  
Recommended  
BRAND

# Dramamine®

**motion sickness**

**CHEWABLE**

**Dual Action:**

Prevents & Relieves Nausea,  
Dizziness and Vomiting

#1  
Pharmacist  
Recommended  
BRAND

DIMENHYDRINATE  
TABLETS / ANTIEMETIC

# Dramamine®

**motion sickness**

**CHEWABLE**

**Dual Action:**

Prevents & Relieves Nausea,  
Dizziness and Vomiting

#1  
Pharmacist  
Recommended  
BRAND

DIMENHYDRINATE  
TABLETS / ANTIEMETIC

# Dramamine®

**motion sickness**

**CHEWABLE**

**Dual Action:**

Prevents & Relieves Nausea,  
Dizziness and Vomiting



**TO OPEN  
PUSH IN TAB AND PULL OUT**



**25 Packets of 2 Orange Flavor Tablets  
(50 mg each)**

### Drug Facts

**Active Ingredient (In each tablet)**

Dimenhydrinate 50 mg

**Purpose**

Antiemetic

**Use**

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

**Warnings**

Do not give to children under 2 years of age unless directed by a doctor

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers****When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a doctor before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.

**Directions**

- to prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

adults and children 12 years and over	• take 1 to 2 chewable tablets every 4-6 hours • do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	• give 1/2 to 1 chewable tablet every 6-8 hours • do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	• give 1/2 chewable tablet every 6-8 hours • do not give more than 1½ chewable tablets in 24 hours, or as directed by a doctor

### Drug Facts (continued)

**Other Information**

- Phenylketonurics: contains phenylalanine 0.84 mg per tablet
- store at room temperature 20° - 25°C (68° - 77°F)
- do not use if pouch is opened
- see bottom of this panel for lot number and expiration date

**Inactive Ingredients**

anhydrous citric acid, aspartame, FD&amp;C yellow #6 aluminum lake, flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified starch, sorbitol

**Questions or comments? call 1-800-382-7219**


©2021 List By:  
Medtech Products, Inc. Tarrytown, NY 10591  
A Prestige Consumer Healthcare Company  
©2021 Trade dress is owned by Medtech Products Inc. All rights reserved.  
www.dramamine.com

Product Repackaged and Distributed  
with permission of manufacturer by:  
Select Corporation Carrollton, TX 75007

Select  
Corporation



2101962239



## DRAMAMINE

dimenhydrinate tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52904-962(NDC:63029-901)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIMENHYDRINATE</b> (UNII: JB937PER5C) (chlortheophylline - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>silicon dioxide</b> (UNII: ETJ7Z6XBU4)	
<b>croscarmellose sodium</b> (UNII: M28OL1HH48)	
<b>lactose, unspecified form</b> (UNII: J2B2A4N98G)	
<b>magnesium stearate</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52904-962-04	1 in 1 BLISTER PACK	01/15/2012	
1		2 in 1 POUCH; Type 0: Not a Combination		

1		Product		
2	NDC:52904-962-25	25 in 1 CARTON	01/15/2012	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	01/15/2012	

**Labeler** - Select Corporation (053805599)

Revised: 4/2022

Select Corporation