

SINUS AND ALLERGY MAXIMUM STRENGTH- chlorpheniramine maleate, phenylephrine hcl tablet

Chain Drug Consortium

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Premier Value 44-462

Active ingredients (in each tablet)

Chlorpheniramine maleate 4 mg

Phenylephrine HCl 10 mg

Purpose

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes
 - sinus congestion and pressure
 - nasal congestion

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- **do not exceed recommended dose**
- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.
- children under 12 years: do not use this product in children under 12 years of age

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, lactose, magnesium stearate, microcrystalline cellulose, silica gel, stearic acid

Questions or comments?

1-800-426-9391

Principal Display Panel

**Premier
Value®**

NDC 68016-687-24

***COMPARE TO THE ACTIVE INGREDIENTS IN
SUDAFED PE® SINUS + ALLERGY**

**Maximum Strength
Sinus & Allergy**

Chlorpheniramine maleate 4 mg - **ANTIHISTAMINE**
Phenylephrine HCl 10 mg - **NASAL DECONGESTANT**

- Sneezing • Itchy eyes • Runny nose
- Sinus pressure + congestion

24 Tablets

INDEPENDENTLY TESTED

PV

SATISFACTION GUARANTEED

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Sudafed PE® Sinus + Allergy.

50844 ORG061246208

DISTRIBUTED BY: CHAIN CONSORTIUM, LLC

UPARC, BLDG. A3, SUITE 338

1020 WILLIAM PITT WAY

PITTSBURGH, PA 15238

www.chaindrugconsortium.com

If for any reason you are not satisfied with
this product, please return it to the store
where purchased for a full refund.



Maximum Strength Sinus & Allergy



Maximum Strength Sinus & Allergy

Chlorpheniramine maleate 4 mg- **ANTIHISTAMINE**
Phenylephrine HCl 10 mg- **NASAL DECONGESTANT**

- Sneezing • Itchy eyes • Runny nose
- Sinus pressure + congestion

24 Tablets

NDC 68016-687-24

*COMPARE TO THE ACTIVE INGREDIENTS IN
SUDAFED PE® SINUS + ALLERGY

OMIT W

TAMPER EVIDENT: DO NOT USE IF
CARTON IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING



Premier Value
Maximum Strength
Sinus & Allergy



B-1590-462-08
DRG061246208



DISTRIBUTED BY: CHAM DRUG CONSORTIUM, LLC
UPARC, BLDG. A3, SUITE 338
1020 WILLIAM PITT WAY
PITTSBURGH, PA 15238
www.chamdrugconsortium.com

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Sudafed PE Sinus + Allergy.
50844 DRG061246208

| | |
|--|-------------------|
| Drug Facts | |
| Active ingredients (in each tablet) Chlorpheniramine maleate 4 mg.....Antihistamine Phenylephrine HCl 10 mg.....Nasal decongestant | |
| Uses Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ sinus congestion and pressure ■ nasal congestion | |
| Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ high blood pressure ■ heart disease ■ thyroid disease ■ diabetes ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. When using this product ■ do not exceed recommended dose ■ excitability may occur, especially in children | |
| Directions ■ adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours. ■ children under 12 years: do not use this product in children under 12 years of age | |
| Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number | |
| Inactive ingredients croscarmellose sodium, lactose, magnesium stearate, microcrystalline cellulose, silica gel, stearic acid | |
| Questions or comments? 1-800-426-9391 | |
| Drug Facts (continued) | Drug Facts |
| Stop use and ask a doctor if ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery | Drug Facts |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. | Drug Facts |
| Directions ■ adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours. ■ children under 12 years: do not use this product in children under 12 years of age | Drug Facts |
| Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number | Drug Facts |
| Inactive ingredients croscarmellose sodium, lactose, magnesium stearate, microcrystalline cellulose, silica gel, stearic acid | Drug Facts |
| Questions or comments? 1-800-426-9391 | Drug Facts |

Premier Value 44-462

SINUS AND ALLERGY MAXIMUM STRENGTH
chlorpheniramine maleate, phenylephrine hcl tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68016-687 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-----------------------------|----------|
| CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U) | CHLORPHENIRAMINE MALEATE | 4 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | WHITE | Score | no score |
| Shape | ROUND | Size | 7mm |
| Flavor | | Imprint Code | 44;462 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:68016-687-24 | 1 in 1 CARTON | 06/09/2005 | |
| 1 | | 24 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part341 | 06/09/2005 | |

Labeler - Chain Drug Consortium (101668460)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867894 | MANUFACTURE(68016-687) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 832867837 | PACK(68016-687) |

Revised: 6/2019

Chain Drug Consortium