

ACETAMINOPHEN- acetaminophen tablet, coated
Strategic Sourcing Services LLC

SMK - 1001 - 2018-1230

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 4 to 6 hours while symptoms last▪ do not take more than 10 tablets in 24 hours▪ do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">▪ take 1 tablet every 4 to 6 hours while symptoms last▪ do not take more than 5 tablets in 24 hours▪ do not use for more than 5 days unless directed by a doctor
children under 6 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate, stearic acid

PRINCIPAL DISPLAY PANEL

sunmark®

COMPARE TO TYLENOL® REGULAR STRENGTH TABLETS

ACTIVE INGREDIENT†

NDC 49348-973-10

pain relief

Regular Strength

Pain reliever & Fever reducer

ACETAMINOPHEN

- Contains No Aspirin

100 TABLETS 325 mg EACH

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



F100102SMK Rt

DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING



This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength tablets.

NC

Drug Facts

Active ingredient (in each tablet) **Purpose**
Acetaminophen 325 mg.....Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ backache
■ minor pain of arthritis ■ toothache ■ muscular aches
■ premenstrual and menstrual cramps
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease. ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur ■ redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years and over
■ take 2 tablets every 4 to 6 hours while symptoms last
■ do not take more than 10 tablets in 24 hours
■ do not use for more than 10 days unless directed by a doctor

children 6-11 years
■ take 1 tablet every 4 to 6 hours while symptoms last
■ do not take more than 5 tablets in 24 hours
■ do not use for more than 5 days unless directed by a doctor

children under 6 years
■ ask a doctor

Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate, stearic acid

McKesson

Another Quality Product
Distributed By McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
Please visit us at www.sunmarkbrand.com

NC



ACETAMINOPHEN

acetaminophen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-973
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	M2A3;57344
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-973-10	1 in 1 CARTON	09/01/2017	09/30/2025
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:49348-973-16	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/01/2017	07/31/2022

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/13/2012	09/30/2025

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 10/2023

Strategic Sourcing Services LLC