

**RAPID RELIEF PLANTAR FASCIITIS PAIN RELIEF PATCH- lidocaine  
menthol patch  
RAPID AID VIET NAM CO., LTD**

-----

**Active Ingredient(s)**

Lidocaine 4%  
Menthol 1%

**Purpose**

Topical anesthetic  
Topical analgesic

**Use**

For temporary relief of pain.

**Warnings**

For external use only.

**Do not use**

More than 1 patch at a time . On wounds or damaged skin . With a heating pad such as electric heating pad, as this may result in excessive skin irritation or skin burn . If you are allergic to any ingredients of this product . In large quantities, particularly over raw surfaces or blistered areas . More than one week without consulting a doctor.

**When using this product:**

Use only as directed . Do not apply to wounds or damaged, broken, or irritated skin . Avoid contact with eye mucous membranes or rashes. If contact occurs, rinse thoroughly . Do not bandage tightly. Do not use at the same time as other topical analgesics.

**Stop use and ask a doctor if:**

. Localized skin reactions occur, such as rash, itching, redness, irritation, pain and blistering as rare but serious burns can occur . Condition worsens . Symptoms persist for more than 7 days . Symptoms clear up and occur again within a few days.

If pregnant or breast feeding, ask a physician before use.

## KEEP OUT OF REACH OF CHILDREN

If swallowed, get medical help or contact a Poison Control Center right away.

## Directions

Adults and children 12 years of age and older: Clean and dry affected area · Remove film from patch and apply sticky side to desired area · Apply patch to affected area not more than 3 times daily. Do not leave on skin for more than a total of 8 hours per day. Can be cut for custom sizing of pain area . Children under 12 years of age: Consult a doctor . Supervise children when they use this product.

## STORAGE AND HANDLING

Store between 15-30°C (59-86°F)

## Inactive ingredients

Acrylamide/Sodium acryloyldimethyltaurate copolymer, Dihydroxyaluminum aminoacetate, DMDM hydantoin, Edetic acid, Glycerin, Laurocapram, Polyacrylic acid, Propylene glycol, Sodium polyacrylate, Tartaric acid, Titanium dioxide, Water.

## Package Label - Principal Display Panel

 <p><b>RAPID RELIEF</b> Stay in Motion</p> <p><b>Plantar Fasciitis Pain Relief Patch</b> <b>Lidocaine 4% + Menthol 1%</b> Topical Anesthetic &amp; Analgesic</p> <p>Numbs aggravated nerves Temporary relief from pain associated with</p> <ul style="list-style-type: none"><li>• Heel &amp; arch</li><li>• Ball of foot</li><li>• Arthritis</li></ul> <p><b>USE UP TO 8 HOURS</b></p> <p><b>8 Patches</b></p> <p>2.5" x 4" (6.4cm x 10.2cm)</p>	 <ul style="list-style-type: none"><li>• Numbs aggravated nerves</li><li>• Portable and convenient</li><li>• No messy residue</li></ul> <p>Manufactured by: Rapid Aid Vietnam Co., Ltd. Cau Ha A Hamlet, An Quang Commune, Hai Phong City, Viet Nam, 180000 Made in Vietnam</p> <p>Imported by: Rapid Aid Corp., Mississauga, ON L5L 5Z3 CANADA</p> <p>MPN250080808 0 11 64317 25008 4</p>	<p><b>Do Not Reuse</b>      Consult instructions for use</p> <table border="1"><tr><td><b>Drug Facts</b></td></tr><tr><td><b>Active ingredient</b> Lidocaine 4% ..... Topical anesthetic Menthol 1% ..... Topical analgesic</td></tr><tr><td><b>Uses</b> For temporary relief of pain.</td></tr><tr><td><b>Warnings</b> For external use only.</td></tr><tr><td><b>Do not use:</b> • More than 1 patch at a time · On wounds or damaged skin · With a heating pad such as electric heating pad, as this may result in excessive skin irritation or skin burn · If you are allergic to any ingredients of this product · In large quantities, particularly over raw surfaces or blistered areas · More than one week without consulting a doctor.</td></tr><tr><td><b>When using this product:</b> • Use only as directed · Do not apply to wounds or damaged, broken, or irritated skin · Avoid contact with eyes, mucous membranes or rashes. If contact occurs, rinse thoroughly · Do not bandage tightly · Do not use at the same time as other topical analgesics.</td></tr><tr><td><b>Stop use and ask a doctor if:</b> • Localized skin reactions occur, such as rash, itching, redness, irritation, pain and blistering as rare but serious burns can occur · Condition worsens · Symptoms persist for more than 7 days · Symptoms clear up and occur again within a few days.</td></tr><tr><td><b>If pregnant or breast feeding,</b> ask a physician before use. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</td></tr><tr><td><b>Directions</b> <b>Adults and children 12 years of age and older:</b> • Clean and dry affected area · Remove film from patch and apply sticky side to desired area · Apply patch to affected area not more than 3 times daily. Do not leave on skin for more than a total of 8 hours per day · Can be cut for custom sizing of pain area · <b>Children under 12 years of age:</b> Consult a doctor · Supervise children when they use this product.</td></tr><tr><td><b>Other information</b> • Store between 15-30°C (59-86°F)</td></tr><tr><td><b>Inactive ingredients</b> Acrylamide/Sodium acryloyldimethyltaurate copolymer, Dihydroxyaluminum aminoacetate, DMDM hydantoin, Edetic acid, Glycerin, Laurocapram, Polyacrylic acid, Propylene glycol, Sodium polyacrylate, Tartaric acid, Titanium dioxide, Water.</td></tr></table> <p>Questions or Comments?      1-800-COLD-HOT      www.rapidaid.com</p>	<b>Drug Facts</b>	<b>Active ingredient</b> Lidocaine 4% ..... Topical anesthetic Menthol 1% ..... Topical analgesic	<b>Uses</b> For temporary relief of pain.	<b>Warnings</b> For external use only.	<b>Do not use:</b> • More than 1 patch at a time · On wounds or damaged skin · With a heating pad such as electric heating pad, as this may result in excessive skin irritation or skin burn · If you are allergic to any ingredients of this product · In large quantities, particularly over raw surfaces or blistered areas · More than one week without consulting a doctor.	<b>When using this product:</b> • Use only as directed · Do not apply to wounds or damaged, broken, or irritated skin · Avoid contact with eyes, mucous membranes or rashes. If contact occurs, rinse thoroughly · Do not bandage tightly · Do not use at the same time as other topical analgesics.	<b>Stop use and ask a doctor if:</b> • Localized skin reactions occur, such as rash, itching, redness, irritation, pain and blistering as rare but serious burns can occur · Condition worsens · Symptoms persist for more than 7 days · Symptoms clear up and occur again within a few days.	<b>If pregnant or breast feeding,</b> ask a physician before use. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	<b>Directions</b> <b>Adults and children 12 years of age and older:</b> • Clean and dry affected area · Remove film from patch and apply sticky side to desired area · Apply patch to affected area not more than 3 times daily. Do not leave on skin for more than a total of 8 hours per day · Can be cut for custom sizing of pain area · <b>Children under 12 years of age:</b> Consult a doctor · Supervise children when they use this product.	<b>Other information</b> • Store between 15-30°C (59-86°F)	<b>Inactive ingredients</b> Acrylamide/Sodium acryloyldimethyltaurate copolymer, Dihydroxyaluminum aminoacetate, DMDM hydantoin, Edetic acid, Glycerin, Laurocapram, Polyacrylic acid, Propylene glycol, Sodium polyacrylate, Tartaric acid, Titanium dioxide, Water.
<b>Drug Facts</b>													
<b>Active ingredient</b> Lidocaine 4% ..... Topical anesthetic Menthol 1% ..... Topical analgesic													
<b>Uses</b> For temporary relief of pain.													
<b>Warnings</b> For external use only.													
<b>Do not use:</b> • More than 1 patch at a time · On wounds or damaged skin · With a heating pad such as electric heating pad, as this may result in excessive skin irritation or skin burn · If you are allergic to any ingredients of this product · In large quantities, particularly over raw surfaces or blistered areas · More than one week without consulting a doctor.													
<b>When using this product:</b> • Use only as directed · Do not apply to wounds or damaged, broken, or irritated skin · Avoid contact with eyes, mucous membranes or rashes. If contact occurs, rinse thoroughly · Do not bandage tightly · Do not use at the same time as other topical analgesics.													
<b>Stop use and ask a doctor if:</b> • Localized skin reactions occur, such as rash, itching, redness, irritation, pain and blistering as rare but serious burns can occur · Condition worsens · Symptoms persist for more than 7 days · Symptoms clear up and occur again within a few days.													
<b>If pregnant or breast feeding,</b> ask a physician before use. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.													
<b>Directions</b> <b>Adults and children 12 years of age and older:</b> • Clean and dry affected area · Remove film from patch and apply sticky side to desired area · Apply patch to affected area not more than 3 times daily. Do not leave on skin for more than a total of 8 hours per day · Can be cut for custom sizing of pain area · <b>Children under 12 years of age:</b> Consult a doctor · Supervise children when they use this product.													
<b>Other information</b> • Store between 15-30°C (59-86°F)													
<b>Inactive ingredients</b> Acrylamide/Sodium acryloyldimethyltaurate copolymer, Dihydroxyaluminum aminoacetate, DMDM hydantoin, Edetic acid, Glycerin, Laurocapram, Polyacrylic acid, Propylene glycol, Sodium polyacrylate, Tartaric acid, Titanium dioxide, Water.													

## RAPID RELIEF PLANTAR FASCIITIS PAIN RELIEF PATCH

lidocaine menthol patch

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83569-007
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>MENTHOL</b> (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	1 g in 100 g
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>POLYACRYLIC ACID (300000 MW)</b> (UNII: A8371R0U5J)	
<b>POLYACRYLIC ACID (800000 MW)</b> (UNII: D0I6NSZ87U)	
<b>DIHYDROXYALUMINUM AMINOACETATE</b> (UNII: DO250MG0W6)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>DMDM HYDANTOIN</b> (UNII: BYR0546TOW)	
<b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM)	
<b>SODIUM POLYACRYLATE (8000 MW)</b> (UNII: 285CYO341L)	
<b>ACRYLAMIDE/SODIUM ACRYLOYLDIMETHYLTAURATE COPOLYMER (120000 MPA.S AT 1%)</b> (UNII: 5F4963KLHS)	
<b>TARTARIC ACID</b> (UNII: W4888I119H)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>LAUROCAPRAM</b> (UNII: 1F3X9DRV9X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:83569-007-01	1 in 1 BOX	07/10/2025	
1		6 in 1 POUCH		
1		5 g in 1 PATCH; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M017	07/10/2025	

**Labeler** - RAPID AID VIET NAM CO., LTD (673067008)

**Registrant** - RAPID AID VIET NAM CO., LTD (673067008)

**Establishment**

Name	Address	ID/FEI	Business Operations
RAPID AID VIET NAM CO., LTD		673067008	manufacture(83569-007)

Revised: 10/2025

RAPID AID VIET NAM CO., LTD