

ESOMEPRAZOLE MAGNESIUM- esomeprazole magnesium capsule, delayed release
Thirty Madison Inc

Drug Facts

Active ingredient(s) in each capsule

*Esomeprazole 20 mg

(Each delayed-release capsule corresponds to 20.7 mg esomeprazole magnesium, USP)

Purpose

Acid reducer

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadness
- frequent **chest pain.**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus or mycophenolate mofetil (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- each capsule contains: magnesium, 15 mg
- read directions and warnings before use
- keep the carton. It contains important information
- store at 20 to 25°C (68 to 77°F)

Inactive ingredients

acetyl tributyl citrate, dibutyl sebacate, FD&C Blue #1, ferrous ferric oxide, gelatin, glyceryl monostearate, hypromellose, magnesium oxide, magnesium stearate, methacrylic acid copolymer dispersion, poloxamer, polysorbate 80, povidone, potassium hydroxide, propylene glycol, shellac, sugar, talc, titanium dioxide

Questions or comments

call toll-free weekdays 8 AM to 10 PM EST at 1-888-375-3784

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Container Label

Container Label:

evens

Esomeprazole Magnesium
Delayed-Release Capsules
USP, 20mg* / Acid Reducer,
24 hour

Treats frequent heartburn
May take 1 to 4 days for full effect
14 Capsules
One 14-day course of treatment

TAMPER EVIDENT-FEATURES: Do not use if foil seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or black band around the center of each capsule is broken or missing.

KEEP THE CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule)
Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 20.7 mg esomeprazole magnesium, USP)

Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Distributed by Thirty Madison, Inc.
New York, NY 10016
Made in India

LOT/EXP 150075941

Drug Facts (continued)

- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or dicitazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information ■ each capsule contains: magnesium, 15 mg


- read the directions and warnings before use
- keep the carton. It contains important information: ■ store at 20 to 25°C (68 to 77°F)

Inactive ingredients
acetyl tributyl citrate, dibutyl sebacate, FD&C Blue #1, ferrous ferric oxide, gelatin, glyceryl monostearate, hypromellose, magnesium oxide, magnesium stearate, methacrylic acid copolymer dispersion, poloxamer, polysorbate 80, povidone, potassium hydroxide, propylene glycol, shellac, sugar, talc, titanium dioxide

Questions or comments?
1-888-375-3784

call toll-free weekdays 8 AM to 10 PM EST at

Carton label



ESOMEPRAZOLE MAGNESIUM

esomeprazole magnesium capsule, delayed release

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:71713-202 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| Esomeprazole Magnesium (UNII: R6DXU4WAY9) (Esomeprazole - UNII:N3PA6559 FT) | Esomeprazole | 20 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| acetyltributyl citrate (UNII: 0ZBX0N59RZ) | |
| Dibutyl Sebacate (UNII: 4W5IH7FLNY) | |
| Ferrosoferric Oxide (UNII: XM0M87F357) | |
| Gelatin (UNII: 2G86QN327L) | |
| Glyceryl Monostearate (UNII: 230OU9XXE4) | |
| Hypromelloses (UNII: 3NXW29V3WO) | |
| Magnesium Oxide (UNII: 3A3U0GI71G) | |
| Magnesium Stearate (UNII: 70097M6I30) | |
| Methacrylic Acid - Ethyl Acrylate Copolymer (1:1) Type A (UNII: NX76LV5T8J) | |
| Poloxamer 188 (UNII: LQA7B6G8JG) | |
| Polysorbate 80 (UNII: 6OZP39ZG8H) | |
| Povidone K30 (UNII: U725QWY32X) | |
| Talc (UNII: 7SEV7J4R1U) | |
| Titanium Dioxide (UNII: 15FIX9V2JP) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |

| | |
|---|--|
| Potassium Hydroxide (UNII: WZH3C48M4T) | |
| Shellac (UNII: 46N107B71O) | |
| Sucrose (UNII: C151H8M554) | |
| Fd&C Blue No. 1 (UNII: H3R47K3TBD) | |

Product Characteristics

| | | | |
|-----------------|---|---------------------|----------|
| Color | BLUE (light blue cap) , BLUE (dark blue body) | Score | no score |
| Shape | CAPSULE | Size | 16mm |
| Flavor | | Imprint Code | RDY;327 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:71713-202-01 | 1 in 1 CARTON | 05/31/2019 | |
| 1 | | 14 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 2 | NDC:71713-202-03 | 3 in 1 CARTON | 05/31/2019 | |
| 2 | | 14 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA207673 | 05/31/2019 | |

Labeler - Thirty Madison Inc (080774087)