

ACETAMINOPHEN- acetaminophen tablet
MEIJER DISTRIBUTION INC

Acetaminophen Extended-release tablets USP, 650 mg
Pain Reliever/Fever Reducer
For The Temporary Relief of Minor Arthritis Pain

Contains No Aspirin

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

Active ingredient (in each caplet)

Acetaminophen USP, 650 mg

Purpose

Pain reliever/ fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache
- toothache ■ temporarily reduces fever

Liver Warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions**■ do not take more than directed (see overdose warning)**

adults ■ take 2 caplets every 8 hours with water

■ swallow whole; do not crush, chew, split or dissolve

■ do not take more than 6 caplets in 24 hours

■ do not use for more than 10 days unless directed by a doctor

under 18 years of age ■ ask a doctor

Other Information

■ store between 20-25°C (68-77°F)

■ **do not use if foil inner seal is broken or missing**

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid

contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

meijer.

See New Warning

8HR Arthritis Pain Relief

ACETAMINOPHEN
EXTENDED-RELEASE
TABLETS USP, 650 mg

Pain Reliever/Fever Reducer
For The Temporary Relief Of Minor
Arthritis Pain
Contains No Aspirin

150 Caplets**
650 mg each
(**Capsule-Shaped
Bi-Layer Tablets)

Actual Size

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN
NDC 79481-0183-5

COMPARE TO
TYLENOL® 8 HR
ARTHRITIS PAIN
ACTIVE INGREDIENT*

NOT FOR HOUSEHOLDS
WITH YOUNG CHILDREN

Important: Read all product information before using.

Do not use if foil inner seal is broken or missing.

Drug Facts

Active ingredient (in each caplet)

Acetaminophen USP, 650 mg. Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - minor pain of arthritis
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Always alert acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- itching
- skin rash
- skin peeling
- blistering
- swelling

If a skin reaction occurs, stop use and seek medical help right away.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

DIST. BY MEIJER DISTRIBUTION, INC. (CONTINUED ON BACK OF LABEL)

GRAND RAPIDS, MI 49544

www.meijer.com

FD 43128



*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark "Tylenol".

MADE IN INDIA

LOT
EXP
2000000004289
700000003292

Inside (adhesive side)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-525-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning).

Take 2 caplets every 8 hours with water.

Swallow whole. Do not crush, chew, split or dissolve.

Do not take more than 6 caplets in 24 hours.

Do not use for more than 10 days or as directed by a doctor.

Ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:

ask a doctor if:



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0183
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	

SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
HYDROXYETHYL CELLULOSE (140 CPS AT 5%) (UNII: 8136Y38GY5)	
STEARIC ACID (UNII: 4ELV7Z 65AP)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ 8WG20P6)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	G650
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0183-5	150 in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2023	
2	NDC:79481-0183-2	225 in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2023	
3	NDC:79481-0183-4	400 in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2023	
4	NDC:79481-0183-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	07/17/2023	

Labeler - MEIJER DISTRIBUTION INC (006959555)

Revised: 12/2025

MEIJER DISTRIBUTION INC