

**ACETAMINOPHEN- acetaminophen tablet**  
**MEIJER, INC.**

-----

**8HR MUSCLE ACHES & PAIN RELIEF**

**Acetaminophen Extended-release tablets USP, 650 mg**

**Pain Reliever/Fever Reducer**

**For up to 8 Hours Relief of Minor Muscle Aches & Pain**  
**Contains No Aspirin**

**See New Warning**

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

**Active ingredient (in each caplet)**

Acetaminophen USP, 650 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- muscular aches
- backache
- minor pain of arthritis
- toothache
- premenstrual and menstrual cramps
- headache
- the common cold
- temporarily reduces fever

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

## **Ask a doctor before use if you have**

liver disease

## **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

## **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

Keep out of reach of children.

## **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

### **■ do not take more than directed (see overdose warning)**

adults and children ■ take 2 caplets every 8 hours with water

■ swallow whole; do not crush, chew, split or dissolve

■ do not take more than 6 caplets in 24 hours

■ do not use for more than 10 days unless directed by a doctor 12 years of age and over





## ACETAMINOPHEN

acetaminophen tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0184
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>HYDROXYETHYL CELLULOSE (140 CPS AT 5%)</b> (UNII: 8136Y38GY5)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	G650
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0184-8	24 in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2023	
2	NDC:79481-0184-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	07/17/2023	

**Labeler** - MEIJER, INC. (006959555)