

**COLD AND HEAD CONGESTION SEVERE- acetaminophen, guaifenesin,
phenylephrine hcl tablet, film coated
DOLGENCORP, LLC**

Dollar General 44-527C-HC

Active ingredients (in each caplet)

Acetaminophen 325 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Expectorant
Nasal decongestant

Uses

- for the temporary relief of the following cold symptoms:
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- high blood pressure
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- heart disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-888-309-9030

Principal Display Panel

DG® | health

**Compare to
active ingredients
of Tylenol® COLD +
HEAD CONGESTION
SEVERE***

**Cold & Head
Congestion Severe**

Acetaminophen, Guaifenesin, Phenylephrine HCl
Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

Relieves:

- Fever/Headache • Body aches/Sore throat
- Nasal congestion • Mucus/Chest congestion

Pseudoephedrine Free

Non Drowsy

24 Caplets

Actual Caplet Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Johnson & Johnson Corporation,
owner of
the registered trademark Tylenol® COLD + HEAD CONGESTION SEVERE.
50844 REV0922D52708

DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
(888) 309-9030

*This product is not manufactured or distributed by Kenne Inc., owner of the registered trademark Tylenol® COLD + HEAD CONGESTION SEVERE. 50844 REV0922052708
 DISTRIBUTED BY OLD EAST MAIN CO., 100 MISSION RIDGE GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
 (888) 309-9030



Cold & Head Congestion Severe

Acetaminophen, Guaifenesin, Phenylephrine HCl
 Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

- Relieves:
- Fever/Headache
 - Body aches/Sore throat
 - Nasal congestion
 - Mucus/Chest congestion

Non Drowsy



24 Caplets
 Actual Caplet Size

Compare to active ingredients of Tylenol® COLD + HEAD CONGESTION SEVERE*

Drug Facts (continued)
 Other information ■ each caplet contains: sodium 3 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ see end flap for expiration date and lot number

Drug Facts (continued)
 Inactive ingredients com starch, croscollon, flavoc, magnesium stearate, maltodextrin, microcystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-888-309-9030

Drug Facts
 Active ingredients (in each caplet)
 Acetaminophen 325 mg Pain reliever/fever reducer
 Guaifenesin 200 mg Expectorant
 Phenylephrine HCl 5 mg Nasal decongestant

Purpose
 ■ for the temporary relief of the following cold symptoms:
 ■ headache ■ nasal congestion ■ minor aches and pains
 ■ sinus congestion and pressure ■ sore throat
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive ■ temporarily reduces fever

Drug Facts (continued)

Drug Facts (continued)
 Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ 3 or more alcoholic drinks every day while using this product
 ■ with other drugs containing acetaminophen
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin redness
 If a skin reaction occurs, stop use and seek medical help right away.
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
 Do not use
 ■ with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients
 Ask a doctor before use if you have
 ■ liver disease ■ diabetes
 ■ difficulty in urination due to enlargement of the prostate gland

Drug Facts (continued)
 Directions
 ■ do not take more than directed
 ■ adults and children 12 years and over
 ■ take 2 caplets every 4 hours
 ■ swallow whole - do not crush, chew, or dissolve
 ■ do not take more than 10 caplets in 24 hours
 ■ children under 12 years: ask a doctor

Drug Facts (continued)
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
 If pregnant or breast-feeding, ask a health professional before use.
 ■ cough that occurs with too much phlegm (mucus) ■ heart disease
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ high blood pressure ■ thyroid disease
 Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
 When using this product do not exceed recommended dosage.
 Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

B-0315-527C-08-HC
 REV0922052708



A0955
 1 50844 52708 6
 TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING
 No Print/No Varnish Lot and Expiration No.

Dollar General 44-527C

COLD AND HEAD CONGESTION SEVERE
 acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-927
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor	MENTHOL	Imprint Code	44;527
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-927-08	2 in 1 CARTON	08/06/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/06/2005	

Labeler - DOLGENCORP, LLC (068331990)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-927) , pack(55910-927)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-927)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-927)

Revised: 3/2025

DOLGENCORP, LLC