NICOTINE POLACRILEX- nicotine polacrilex gum, chewing Dr. Reddy's Laboratories Inc.

Nicotine Polacrilex Gum Coated Ice Mint 4 mg ANDA204833 - Equate

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex (equal 4 mg nicotine)

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece	1 piece	1 piece
every	every	every
1 to 2	2 to 4	4 to 8
hours	hours	hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

• each piece contains: calcium 120 mg, sodium 8 mg

- store at 20 25°C (68 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 aluminum lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol.

Questions or comments?

Calltoll-free 1-800-585-8682 weekdays (8:00 am - 8:00 pm EST

Principal Display Panel

*Compare to the active ingredient in Nicorrette® Gum

Habitrol

Nicotine Polacrilex Gum

USP, 4 mg (nicotine)

STOP SMOKING AID

Gum

4 mg

SUGAR FREE

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **WITHIN 30 MINUTES** OF WAKING UP.

If you smoke your first cigarette <u>MORE THAN</u> 30 MINUTES after waking up, use Habitrol 2 mg Gum

Coated Mint Flavor

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivate to quit

2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks

3. Use long enough - Use Nicotine Polacrilex Gum for the full 12 weeks

4. Use with a Support Program as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE
- PROOF OF AGE REQUIRED

• NOT FOR SALE IN VENDING MACHINE OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED.

††This product is not manufactured or distributed by Haleon group of companies, distributor Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

DISTRIBUTED BY:

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Package Label



erop uogduceranti may, reungen ve unestauctep voj en Bruto Bruto Bruto are not a sen ar

e vol controlled with medication. Vicotine can increase session and whole the president reaction of the president to a series of the president to the president of the president to the president of t

within the boar (however, do and confinementy use one place allow and the second second to complete transform, must be allowed to use the motion of second the places and second second to complete transform, the law year model to use the second second to complete transform, the law year and the use of a second s

HOA3 em 4 PIECES OST If you smoke your first cigarette MORE THAN 30 MINUTES after MORE THAN 30 MINUTES after More and a second and a second a second MINUTES and a second MINUTES a second OF WAKING UP. FOR THOSE WHO SMOKE THEIR FIRST CLOARETTE WITHIN 30 MINUTES Flavor Dateon Sugar-tree ann repering the angle of the property of the prop CD 8 191 SELICED JO 1000 ung AIM ONIMORIC JOIC

9202/90

Dr. Reddy's Laboratories, Inc. Princeton, NJ 08540

PLD-A6238 FC009480

Nicotine Polacrilex Gum USP, 4 mg (nicotine) STOP SMOKINE ALD	at extent-restricted det in extent-restricted and behave to be staff the needing, and we de staff the staff and a detaint and models. The staff the staff the staff of part and the staff the needing, and and and and the default and the staff the needing and and and and the default and and the staff the need of part and and the staff the staff of and and and and the staff the staff the default and and and and and and and and default and and and and and default and and and and and and default and and and and default and and and and default and and and and default and and and default and and and default and and and and default and and and default and and default and and default and and default	cturward a thace a do und aat on daale yoo undaast tyse buckase mail dinar yyyeen tyse gradee ge dinar yyyeen tyse gradee ge usanga usanga	a di chaque paya de la construction de la const construction de la construction de la constru	bes Aseto nor resets arruns egut eft bru / (stanim DC tuote) eftre ro, mug enticols eft 8 teit eft vir yab rog enorig	BUISTER PACKAGED FOR YOUR PROTECTION. Do not use 11 findividual seals are open or torn do not use 17 findividual seals are open of torn
Ηαριτιο	USB I solucia witridriveri graphare, including nicetine creving, associated with a solucing antiding arredung	Enou 2 of 1 Views every	5 JO 4 JOINTS	sinoria di + Viano scervi	
	Purpose (in the sech chewing piece) Purpose (in the prince) Purpose (in the prince) of the prince of	and have any order for the first the first of the first o	tunim DE nintiw attensio Se xloav ST privolici ort ci 9 of T solvelit	21 di Di Kaladi Subari 21 di Di Kaladi	where proof of age cancol be vehicled a not for eace in vescing machines or from any source a proof of required
mu6 *stranovi ni traibergni avtos ett of eregmo0*	spey brug	nuquoo) spoey brug	(pə		age to areay 61 value ecold of olde vol for a
NDC 12801-303-12					

HABITROL Nicotine Polacrilex Gum Coated Mint Flavor

Product Information	n				
Product Type	HUMAN OTC DR	UG Item Co	de (Source)	NDC:7	5907-303
Route of Administratio	n ORAL				
Active Ingredient/Ac	-				
Ir	ngredient Name		Basis of S	Strength	Strength
NICOTINE (UNII: 6M3C89ZY	6R) (NICOTINE - UNII:6M	13C89ZY6R)	NICOTINE		4 mg
Inactive Ingredients	5				
	Ingredier	nt Name			Strength
ACESULFAME POTASSIUM	(UNII: 230V73Q5G9)				
ACACIA (UNII: 5C5403N260)					
CARNAUBA WAX (UNII: R12					
D&C YELLOW NO. 10 (UNI					
HYDROXYPROPYL CELLUL		JNII: 9XZ8H6N6OH)			
MAGNESIUM OXIDE (UNII: 3					
SODIUM BICARBONATE (U					
SODIUM CARBONATE (UNI	: 45P3261C7T)				
TALC (UNII: 7SEV7J4R1U)					
TITANIUM DIOXIDE (UNII: 1					
XYLITOL (UNII: VCQ006KQ1	Ξ)				
Product Characteris	tico				
	yellow	5 e e # e			_
Color	SQUARE	Score		no score	2
Shape Flavor	MINT	Size		1411111	
Contains		Imprint Code			
contains					
Packaging					
			Marketing Sta	rt Mar	keting End
# Item Code	Package Descr		marketing Stal	rt mar	kelind ENO

1	NDC:75907-303- 15	150 in 1 CARTON	06/12/2025				
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product					
_	Marketing Information						
Μ	larketing	Information					
Μ	larketing Marketing Category	Information Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
	Marketing	Application Number or Monograph	-	-			

Labeler - Dr. Reddy's Laboratories Inc. (802315887)

Revised: 6/2025

Dr. Reddy's Laboratories Inc.