

# JATENZO- testosterone undecanoate capsule, liquid filled TOLMAR Inc.

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use JATENZO® safely and effectively. See full prescribing information for JATENZO.

JATENZO (testosterone undecanoate) capsules, for oral use CIII  
Initial U.S. Approval: 1953

## RECENT MAJOR CHANGES

Boxed Warning, Increases in Blood Pressure	Removed 07/2025
Indications and Usage (1)	07/2025
Contraindications, Men with “age related hypogonadism” (4)	Removed 07/2025
Warnings and Precautions, Venous Thromboembolism (5.2)	07/2025
Warnings and Precautions, Blood Pressure Increases (5.4)	07/2025
Warnings and Precautions, Cardiovascular Risk (5.3)	Removed 07/2025

## INDICATIONS AND USAGE

JATENZO (testosterone undecanoate) is an androgen indicated for testosterone replacement therapy in adult males for conditions associated with a deficiency or absence of endogenous testosterone (1).

Limitations of use:

- Safety and efficacy of JATENZO in men with “age-related hypogonadism” have not been established (1)
- Safety and efficacy of JATENZO in males less than 18 years old have not been established (1, 8.4).

## DOSAGE AND ADMINISTRATION

- Prior to initiating JATENZO, confirm the diagnosis of hypogonadism by ensuring that serum testosterone concentrations have been measured in the morning on at least two separate days and that these concentrations are below the normal range (2.1).
- Take JATENZO with food (2.2).
- Starting dose: 237 mg orally once in the morning and once in the evening.
- Adjust the dose to a minimum of 158 mg twice daily and a maximum of 396 mg twice daily based on serum testosterone drawn 6 hours after the morning dose at least 7 days after starting treatment or following dose adjustment and periodically thereafter (2.2).

## DOSAGE FORMS AND STRENGTHS

JATENZO (testosterone undecanoate) capsules for oral use are available in the following strengths: 158 mg, 198 mg, 237 mg (3).

## CONTRAINDICATIONS

- Men with breast cancer or known or suspected prostate cancer (4, 5.3).
- Women who are pregnant. Testosterone may cause fetal harm (4, 5.6, 8.1, 8.2).
- Known hypersensitivity to JATENZO or any of its ingredients (4).

## WARNINGS AND PRECAUTIONS

- Polycythemia: Monitor hematocrit approximately every 3 months to detect increased red blood cell mass and polycythemia (5.1).
- Venous thromboembolism (VTE): VTE, including deep vein thrombosis (DVT) and pulmonary embolism (PE), have been reported in patients using testosterone. Evaluate patients with signs or symptoms consistent with DVT or PE (5.2).
- Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer: Monitor patients with benign prostatic hyperplasia (BPH) for worsening of signs and symptoms of BPH (5.3).
- Blood Pressure Increases: JATENZO can increase blood pressure, which can increase cardiovascular risk over time. Measure blood pressure periodically. Not recommended for use in men with uncontrolled hypertension (5.4).
- Abuse of Testosterone: Testosterone has been subject to abuse, typically at doses higher than recommended for the approved indication and in combination with other anabolic androgenic steroids (5.5).
- Potential for Adverse Effects on Spermatogenesis: Exogenous administration of androgens may lead to azoospermia (5.7).
- Edema: Edema, with or without congestive heart failure, may occur in patients with pre-existing cardiac, renal, or hepatic disease (5.9, 6.1).
- Sleep Apnea: Sleep apnea may occur in those with risk factors (5.11).
- Monitor prostate specific antigen (PSA) and lipid concentrations periodically (5.3, 5.12).
- Risk of Depression and Suicide: Depression and suicidal ideation have occurred during clinical trials in patients treated with JATENZO (5.15).

## ADVERSE REACTIONS

Most common adverse reactions (incidence > 2%): polycythemia, diarrhea, dyspepsia, eructation, peripheral edema, nausea, increased hematocrit, headache, prostatomegaly, and hypertension (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Tolmar, Inc. at 1-844-4TO-LMAR or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## DRUG INTERACTIONS

- Androgens may decrease blood glucose and therefore may decrease insulin requirements in diabetic patients (7.1).

- Changes in anticoagulant activity may be seen with androgens. More frequent monitoring of International Normalized Ratio (INR) and prothrombin time is recommended in patients taking warfarin (7.2).
- Use of testosterone with corticosteroids may result in increased fluid retention. Use with caution, particularly in patients with cardiac, renal, or hepatic disease (7.3).
- Concomitant administration of medications that are known to increase blood pressure may lead to additional increases in blood pressure when used with JATENZO (7.4).

----- **USE IN SPECIFIC POPULATIONS** -----

- Geriatric Patients: There are insufficient long-term safety data to assess the potential risks of cardiovascular disease and prostate cancer (8.5).

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 9/2025**

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**RECENT MAJOR CHANGES**

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

2.1 Confirmation of Hypogonadism Before Initiation of JATENZO

2.2 Dosing and Dose Adjustment Information

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Polycythemia

5.2 Venous Thromboembolism

5.3 Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer

5.4 Blood Pressure Increases

5.5 Abuse of Testosterone and Monitoring of Testosterone Concentrations

5.6 Not for Use in Women

5.7 Potential for Adverse Effects on Spermatogenesis

5.8 Hepatic Adverse Effects

5.9 Edema

5.10 Gynecomastia

5.11 Sleep Apnea

5.12 Lipid Changes

5.13 Hypercalcemia

5.14 Decreased Thyroxine-binding Globulin

5.15 Risk of Depression and Suicide

**6 ADVERSE REACTIONS**

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

7.1 Insulin

7.2 Oral Vitamin K Antagonist Anticoagulants

7.3 Corticosteroids

7.4 Medications that May Also Increase Blood Pressure

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

**9 DRUG ABUSE AND DEPENDENCE**

9.1 Controlled Substance

9.2 Abuse

9.3 Dependence

**10 OVERDOSAGE**

**11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

## **14 CLINICAL STUDIES**

14.1 Clinical Trials in Hypogonadal Males

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

JATENZO (testosterone undecanoate) is an androgen indicated for testosterone replacement therapy in adult males for conditions associated with a deficiency or absence of endogenous testosterone:

- Primary hypogonadism (congenital or acquired): testicular failure due to conditions such as cryptorchidism, bilateral torsion, orchitis, vanishing testis syndrome, orchiectomy, Klinefelter syndrome, chemotherapy, or toxic damage from alcohol or heavy metals. These men usually have low serum testosterone concentrations and gonadotropins (follicle-stimulating hormone [FSH], luteinizing hormone [LH]) above the normal range.
- Hypogonadotropic hypogonadism (congenital or acquired): gonadotropin or luteinizing hormone-releasing hormone (LHRH) deficiency or pituitary-hypothalamic injury from tumors, trauma, or radiation. These men have low testosterone serum concentrations but have gonadotropins in the normal or low range.

#### **Limitations of use:**

- Safety and efficacy of JATENZO in men with “age-related hypogonadism” have not been established.
- Safety and efficacy of JATENZO in males less than 18 years old have not been established [see *Use in Specific Populations (8.4)*].

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Confirmation of Hypogonadism Before Initiation of JATENZO**

Prior to initiating JATENZO, confirm the diagnosis of hypogonadism by ensuring that serum testosterone concentrations have been measured in the morning on at least two separate days and that these testosterone concentrations are below the normal range.

#### **2.2 Dosing and Dose Adjustment Information**

Individualize the dosage of JATENZO based on the patient's serum testosterone concentration response to the drug. The recommended starting dose is 237 mg taken orally twice daily, once in the morning and once in the evening. Take JATENZO with food.

#### *Dose Adjustment*

To ensure proper dose adjustment, measure serum testosterone concentrations 6 hours after the morning dose in plain tubes, clotted at room temperature for 30 minutes prior to centrifugation. Adjust the JATENZO dose based on this serum testosterone measurement as shown in Table 1. Wait seven days after starting treatment or adjusting

the dose before checking the serum testosterone concentration. Thereafter, periodically monitor serum testosterone concentrations 6 hours after the morning dose.

Administer the same dose in the morning and evening. The minimum recommended dose is 158 mg twice daily. The maximum recommended dose is 396 mg (two 198 mg capsules) twice daily.

**Table 1: JATENZO Dose Adjustment Scheme**

<b>Testosterone Concentration in Serum From Plain Tube Drawn 6 hours After Morning Dose</b>	<b>Current JATENZO Dose (mg, twice daily)</b>	<b>New JATENZO Dose (mg, twice daily)</b>
Less than 425 ng/dL	158	198
	198	237
	237	316 (two 158 mg capsules)
	316 (two 158 mg capsules)	396 (two 198 mg capsules)
425 ng/dL – 970 ng/dL	No Dose Change	
More than 970 ng/dL	396 (two 198 mg capsules)	316 (two 158 mg capsules)
	316 (two 158 mg capsules)	237
	237	198
	198	158
	158	Discontinue Treatment

### 3 DOSAGE FORMS AND STRENGTHS

JATENZO capsules for oral use are available in three strengths:

- The 158 mg testosterone undecanoate capsules are opaque red and imprinted with “158” in white ink.
- The 198 mg testosterone undecanoate capsules are opaque white and imprinted with “198” in red ink.
- The 237 mg testosterone undecanoate capsules are opaque orange and imprinted with “237” in white ink.

### 4 CONTRAINDICATIONS

JATENZO is contraindicated in:

- Men with carcinoma of the breast or known or suspected carcinoma of the prostate [see *Warnings and Precautions (5.3)*].
- Women who are pregnant. Testosterone can cause virilization of the female fetus when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].
- Men with known hypersensitivity to JATENZO or any of its ingredients [see *Description (11)*].

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Polycythemia

Increases in hematocrit reflective of increases in red blood cell mass, may require lowering the dose or discontinuation of JATENZO. Check that hematocrit is not elevated prior to initiating JATENZO. Evaluate hematocrit approximately every 3 months while the patient is on JATENZO. If hematocrit becomes elevated, stop JATENZO until the hematocrit decreases to an acceptable concentration. If JATENZO is restarted and again causes hematocrit to become elevated, stop JATENZO permanently. An increase in red blood cell mass may increase the risk of thromboembolic events [see *Warnings and Precautions (5.2)*].

#### 5.2 Venous Thromboembolism

There have been postmarketing reports of venous thromboembolic events, including

deep vein thrombosis (DVT) and pulmonary embolism (PE), in patients using testosterone replacement products such as JATENZO.

In the Testosterone Replacement therapy for Assessment of long-term Vascular Events and efficacy ResponSE in hypogonadal men (TRAVERSE) Study, a randomized, double-blind, placebo-controlled, cardiovascular (CV) outcomes study, compared to placebo, topical testosterone gel was associated with a numerically higher incidence of VTE (1.7% vs 1.2%) which included DVT (0.6% vs 0.5%) and PE events (0.9% vs 0.5%) [see *Adverse Reactions (6.1)*].

Evaluate patients who report symptoms of pain, edema, warmth and erythema in the lower extremity for DVT and those who present with acute shortness of breath for PE. If a venous thromboembolic event is suspected, discontinue treatment with JATENZO and initiate appropriate workup and management [see *Adverse Reactions (6.2)*].

### **5.3 Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer**

Patients with BPH treated with androgens are at an increased risk for worsening of signs and symptoms of BPH. Monitor patients with BPH for worsening signs and symptoms.

Patients treated with androgens may be at increased risk for prostate cancer. Evaluate patients for prostate cancer prior to initiating and during treatment with androgens [see *Contraindications (4)*].

### **5.4 Blood Pressure Increases**

JATENZO can increase blood pressure. Ambulatory blood pressure monitoring (ABPM) demonstrated JATENZO increased systolic /diastolic BP by an average of 4.9/2.5 mm Hg from baseline after 4 months of treatment in a clinical trial [see *Adverse Reactions (6.1)*]. In patients with hypertension on antihypertensive therapy, JATENZO increased the mean systolic/diastolic BP by 5.4/3.2 mm Hg from baseline. Average blood pressures had not plateaued at the end of the trial.

The CV risk associated with topical testosterone gel was evaluated in TRAVERSE, a randomized, double-blind, placebo-controlled, CV outcomes study in men with a history of CV disease or multiple CV risk factors. In TRAVERSE, topical testosterone gel increased mean systolic blood pressure by 1.0 mm Hg from baseline to 36 months, whereas a mean decrease from baseline of 0.5 mm Hg was observed in the placebo group at this timepoint, for a mean between-group difference of 1.5 mm Hg. However, the incidences of major adverse cardiovascular events (MACE), including cardiovascular death, non-fatal myocardial infarction [MI] and non-fatal stroke, were similar between treatment groups (7% for topical testosterone gel vs 7.3% for placebo) [See *Adverse Reactions (6.1)*].

Monitor blood pressure periodically in men using JATENZO, especially men with hypertension. JATENZO is not recommended for use in patients with uncontrolled hypertension.

### **5.5 Abuse of Testosterone and Monitoring of Testosterone Concentrations**

Testosterone has been subject to abuse, typically at doses higher than recommended for the approved indication and in combination with other anabolic androgenic steroids. Anabolic androgenic steroid abuse can lead to serious cardiovascular and psychiatric adverse reactions [see *Drug Abuse and Dependence (9)*].

If testosterone abuse is suspected, check testosterone concentrations to ensure they are within therapeutic range [see *Dosage and Administration (2)*]. However, testosterone levels may be in the normal or subnormal range in men abusing synthetic testosterone derivatives. Counsel patients concerning the serious adverse reactions associated with abuse of testosterone and anabolic androgenic steroids. Conversely, consider the possibility of testosterone and anabolic androgenic steroid abuse in suspected patients who present with serious cardiovascular or psychiatric adverse events.

## **5.6 Not for Use in Women**

Due to lack of controlled studies in women and potential virilizing effects, JATENZO is not indicated for use in women [see *Contraindications (4) and Use in Specific Populations (8.1, 8.2)*].

## **5.7 Potential for Adverse Effects on Spermatogenesis**

With large doses of exogenous androgens, including JATENZO, spermatogenesis may be suppressed through feedback inhibition of pituitary FSH possibly leading to adverse effects on semen parameters including sperm count [see *Use in Specific Populations (8.3)*]. Patients should be informed of this possible risk when deciding whether to use or to continue to use JATENZO.

## **5.8 Hepatic Adverse Effects**

Prolonged use of high doses of orally active 17-alpha-alkyl androgens (e.g., methyltestosterone) has been associated with serious hepatic adverse effects (peliosis hepatis, hepatic neoplasms, cholestatic hepatitis, and jaundice). Peliosis hepatis can be a life-threatening or fatal complication. Long-term therapy with intramuscular testosterone enanthate has produced multiple hepatic adenomas. JATENZO is not known to cause these adverse effects. Nonetheless, patients should be instructed to report any signs or symptoms of hepatic dysfunction (e.g. jaundice). If these occur, promptly discontinue JATENZO while the cause is evaluated.

## **5.9 Edema**

Androgens, including JATENZO, may promote retention of sodium and water. Edema, with or without congestive heart failure, may be a serious complication in patients with pre-existing cardiac, renal, or hepatic disease. In addition to discontinuation of the drug, diuretic therapy may be required.

## **5.10 Gynecomastia**

Gynecomastia may develop and persist in patients being treated for hypogonadism.

## **5.11 Sleep Apnea**

The treatment of hypogonadal men with testosterone may potentiate sleep apnea in some patients, especially those with risk factors such as obesity or chronic lung disease.

## **5.12 Lipid Changes**

Changes in the serum lipid profile may require dose adjustment of lipid lowering drugs or discontinuation of testosterone therapy. Monitor the lipid profile periodically, particularly after starting testosterone therapy.

## **5.13 Hypercalcemia**

Androgens, including JATENZO, should be used with caution in cancer patients at risk of hypercalcemia (and associated hypercalciuria). Monitor serum calcium concentrations regularly during treatment with JATENZO in these patients.

## **5.14 Decreased Thyroxine-binding Globulin**

Androgens, including JATENZO, may decrease concentrations of thyroxin-binding globulin, resulting in decreased total T4 serum concentrations and increased resin uptake of T3 and T4. Free thyroid hormone concentrations remain unchanged, however, and there is no clinical evidence of thyroid dysfunction.

## **5.15 Risk of Depression and Suicide**

Depression and suicidal ideation has been reported in patients treated with JATENZO in clinical trials. Advise patients and caregivers to seek medical attention for manifestations of new onset or worsening depression, suicidal ideation or behavior, anxiety, or other mood changes [see *Adverse Events (6.1)*].

## 6 ADVERSE REACTIONS

### 6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of JATENZO was evaluated in a randomized, controlled clinical study with 166 patients treated with JATENZO twice daily with morning and evening meals for approximately 4 months. All patients were started on 237 mg twice daily, then the dose was titrated to 158 mg, 198 mg, 316 mg, or 396 mg twice daily to achieve testosterone concentrations in the eugonadal range.

Table 2 summarizes adverse reactions ( $\geq 2\%$ ) reported in this 4-month study.

**Table 2: Number (%) of Patients with Adverse Reactions  $\geq 2\%$  in a 4-Month Study with JATENZO**

Preferred Term	Overall (N = 166) n (%)
Headache	8 (4.8)
Hematocrit increased	8 (4.8)
Hypertension	6 (3.6)
High-density lipoprotein decreased	5 (3.0)
Nausea	4 (2.4)

Among the 569 patients who received JATENZO in all Phase 2 and 3 trials combined, the following adverse reactions were reported in  $>2\%$  of patients: polycythemia, diarrhea, dyspepsia, eructation, peripheral edema, nausea, increased hematocrit, headache, prostatomegaly, and hypertension.

Three of the 166 patients (1.8%) in the 4-month study experienced adverse reactions that led to premature discontinuation from the study, including rash (n=1) and headache (n=2).

#### *BP Increases*

In the 4-month clinical study, 24-hour ABPM was conducted on 166 patients. ABPM was conducted at baseline and at Day 139 of JATENZO therapy. A total of 135 patients had acceptable ABPM recordings at both time periods and were at least 85% compliant with study drug. In that group, the mean change in 24-hour systolic BP and diastolic BP from baseline to final on-treatment visit on Day 139 (n=135) was +4.9 mm Hg (95% CI 3.5, 6.4) and +2.5 mm Hg (95% CI 1.5, 3.6), respectively.

In patients with a history of hypertension who were receiving antihypertensive therapy, the mean ABPM systolic and diastolic BP increased by 5.4 mm Hg [95% CI 3.3, 7.6] and 3.2 mm Hg [95% CI 1.7, 4.7], respectively [n=67]. In patients with no history of hypertension, the mean ABPM systolic and diastolic blood pressure increased by 4.4 mm Hg [95% CI 2.3, 6.4] and 1.8 mm Hg [95% CI 0.2, 3.3], respectively [n=63].

Twelve (7.2%) patients on JATENZO either started antihypertensive medications or had their antihypertensive regimen increased during the study.

A total of 6 patients were reported to have an adverse reaction of hypertension (2 patients with hypertension and 4 patients with worsening hypertension), and 3 were reported to have an adverse reaction of increased blood pressure.

#### *HR Increases*

JATENZO increased mean heart rate by an average of 2.2 beats per minute (bpm) [95% CI (1.0, 3.3), N=135] during the study. Patients without a history of hypertension had a greater average increase in mean heart rate (2.7 bpm [95% CI (0.8, 4.6), N=63]) compared to patients with treated hypertension (1.9 bpm [95% CI (0.3, 3.5), N=67]).

### *Increases in Hematocrit*

Increases in hematocrit were reported in 8 of the 166 (4.8%) patients, which occurred in the second half of the study. None of these increases led to premature discontinuation of JATENZO.

### *Headaches*

Headaches were reported in 8 of the 166 patients (4.8%) of which three required treatment with analgesics or non-steroidal anti-inflammatory drugs and 2 led to premature discontinuation from the study. Five of these 8 patients had headache events that resolved within 1 to 2 days.

### *Depression and suicidal ideation*

Two of the 166 patients (1.2%) reported either worsening depression (n=1) or new-onset depression (n=1). One of the 569 patients (0.2%) in clinical trials had suicidal ideation. Each patient completed the study.

### *Increases in Serum PSA*

The mean increase from baseline in PSA was 0.2 ng/mL (n=161). Increases in serum PSA concentrations, defined as an increase from baseline of at least 1.4 ng/mL or PSA greater than 4 ng/mL, occurred in 3 (1.9%) of the patients at the final visit.

### Cardiovascular Outcomes

TRVERSE was a randomized, double-blind, cardiovascular outcomes study to assess the cardiovascular (CV) safety of topical testosterone gel compared to placebo in 5198 hypogonadal men aged 45 to 80 years with a history of CV disease or with multiple CV risk factors. The primary outcome was the incidence of the composite endpoint of major adverse cardiovascular events (MACE), consisting of CV death, non-fatal myocardial infarction (MI), and non-fatal stroke.

The mean duration of therapy was approximately 22 months. The mean duration of follow-up was 33 months. Approximately 61% of all patients discontinued topical testosterone gel or placebo therapy.

The mean patient age ( $\pm$ SD) was 63.3 (7.9) years, with 2452 patients aged 65 years or more (47%); 2847 (about 55%) patients had pre-existing cardiovascular disease, whereas 2357 patients (about 45%) had an elevated cardiovascular risk at baseline, and mean BMI was 35kg/m<sup>2</sup>. Approximately 80% of patients were White, 17% were Black, and 3% were of other races or ethnic groups. Approximately 69%, 84%, and 93% had diabetes mellitus, hyperlipidemia, and hypertension, respectively.

The mean serum testosterone concentration at baseline in patients receiving topical testosterone gel was 220.4 ng/dL (n=2596). The mean serum testosterone concentrations at 12 months, 24 months, 36 months, and 48 months in patients receiving topical testosterone gel were 440.5 ng/dL (n=1683), 420.9 ng/dl (n=1125), 428.7 ng/dL (n=731), and 365.2 ng/dL (n=220), respectively.

For patients treated with topical testosterone gel, the incidence of MACE was 7.0% (n=182 events) and for those receiving placebo, the incidence of MACE was 7.3% (n=190 events). The study demonstrated non-inferiority of topical testosterone gel versus placebo because the upper bound of 95% CI was less than the pre-specified risk margin, of 1.5 for MACE (Hazard Ratio 0.96 [95% CI: 0.78, 1.17]).

### Additional Adverse Reactions Reported in TRVERSE

Additional adverse reactions reported in TRVERSE at an incidence rate >2% in either treatment group and greater in topical testosterone gel versus placebo included: nonfatal arrhythmias warranting intervention (5.2% vs 3.3%), atrial fibrillation (3.5% vs 2.4%), acute kidney injury (2.3% vs 1.5%) and bone fracture (3.5% vs 2.5%). For the adverse reaction of bone fracture, each event was adjudicated by clinical review.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of testosterone. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Cardiovascular Disorders:** myocardial infarction, stroke [*see Warnings and Precautions (5.4), Clinical Trial Experience (6.1)*]

**Vascular Disorders:** Venous thromboembolism [*see Warnings and Precautions (5.1)*]

## **7 DRUG INTERACTIONS**

### **7.1 Insulin**

Changes in insulin sensitivity or glycemic control may occur in patients treated with androgens. In diabetic patients, the metabolic effects of androgens may decrease blood glucose and, therefore, may necessitate a decrease in the dose of anti-diabetic medication.

### **7.2 Oral Vitamin K Antagonist Anticoagulants**

Changes in anticoagulant activity may be seen with androgens; therefore, more frequent monitoring of international normalized ratio (INR) and prothrombin time are recommended in patients taking warfarin, especially at the initiation and termination of androgen therapy.

### **7.3 Corticosteroids**

The concurrent use of testosterone with corticosteroids may result in increased fluid retention and requires careful monitoring particularly in patients with cardiac, renal or hepatic disease.

### **7.4 Medications that May Also Increase Blood Pressure**

Some prescription medications and nonprescription analgesic and cold medications contain drugs known to increase blood pressure. Concomitant administration of these medications with JATENZO may lead to additional increases in blood pressure [*see Warnings and Precautions (5.4)*].

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

JATENZO is contraindicated in pregnant women. Testosterone is teratogenic and may cause fetal harm based on data from animal studies and its mechanism of action [*see Contraindications (4) and Clinical Pharmacology (12.1)*]. Exposure of a female fetus to androgens may result in varying degrees of virilization. In animal developmental studies, exposure to testosterone in utero resulted in hormonal and behavioral changes in offspring and structural impairments of reproductive tissues in female and male offspring. These studies did not meet current standards for nonclinical development toxicity studies.

#### Data

##### *Animal Data*

In developmental studies conducted in rats, rabbits, pigs, sheep and rhesus monkeys, pregnant animals received intramuscular injection of testosterone during the period of organogenesis. Testosterone treatment at doses that were comparable to those used for testosterone replacement therapy resulted in structural impairments in both female

and male offspring. Structural impairments observed in females included increased anogenital distance, phallus development, empty scrotum, no external vagina, intrauterine growth retardation, reduced ovarian reserve, and increased ovarian follicular recruitment. Structural impairments seen in male offspring included increased testicular weight, larger seminal tubular lumen diameter, and higher frequency of occluded tubule lumen. Increased pituitary weight was seen in both sexes.

Testosterone exposure in utero also resulted in hormonal and behavioral changes in offspring. Hypertension was observed in pregnant female rats and their offspring exposed to doses approximately twice those used for testosterone replacement therapy.

## **8.2 Lactation**

### Risk Summary

JATENZO is not indicated for use in women.

## **8.3 Females and Males of Reproductive Potential**

### Infertility

During treatment with large doses of exogenous androgens, including JATENZO, spermatogenesis may be suppressed through feedback inhibition of the hypothalamic-pituitary-testicular axis [see *Warnings and Precautions (5.7)*], possibly leading to adverse effects on semen parameters including sperm count. Reduced fertility is observed in some men taking testosterone replacement therapy. Testicular atrophy, subfertility, and infertility have also been reported in men who abuse anabolic androgenic steroids [see *Drug Abuse and Dependence (9.2)*]. With either type of use, the impact on fertility may be irreversible.

## **8.4 Pediatric Use**

The safety and efficacy of JATENZO in pediatric patients less than 18 years old have not been established. Improper use may result in acceleration of bone age and premature closure of epiphyses.

## **8.5 Geriatric Use**

There have not been sufficient numbers of geriatric patients involved in controlled clinical studies utilizing JATENZO to determine whether efficacy or safety in those over 65 years of age differs from younger subjects. No patients over 65 years of age were enrolled in the 4-month efficacy and safety clinical study utilizing JATENZO. Additionally, there is insufficient long-term safety data in geriatric patients utilizing JATENZO to assess the potentially increased risk of cardiovascular disease and prostate cancer.

Geriatric patients treated with androgens may also be at risk for worsening of signs and symptoms of BPH [see *Warnings and Precautions (5.3)*].

## **9 DRUG ABUSE AND DEPENDENCE**

### **9.1 Controlled Substance**

JATENZO contains testosterone undecanoate, which is a Schedule III controlled substance as defined under the Controlled Substances Act.

### **9.2 Abuse**

Drug abuse is intentional non-therapeutic use of a drug, even once, for its rewarding psychological and physiological effects. Abuse and misuse of testosterone are seen in male and female adults and adolescents. Testosterone, often in combination with other anabolic androgenic steroids (AAS), and not obtained by prescription through a pharmacy, may be abused by athletes and bodybuilders. There have been reports of misuse by men taking higher doses of legally obtained testosterone than prescribed and

continuing testosterone despite adverse events or against medical advice.

#### *Abuse-Related Adverse Reactions*

Serious adverse reactions have been reported in individuals who abuse anabolic androgenic steroids and include cardiac arrest, myocardial infarction, hypertrophic cardiomyopathy, congestive heart failure, cerebrovascular accident, hepatotoxicity, and serious psychiatric manifestations, including major depression, mania, paranoia, psychosis, delusions, hallucinations, hostility and aggression.

The following adverse reactions have also been reported in men: transient ischemic attacks, convulsions, hypomania, irritability, dyslipidemias, testicular atrophy, subfertility, and infertility.

The following additional adverse reactions have been reported in women: hirsutism, virilization, deepening of voice, clitoral enlargement, breast atrophy, male-pattern baldness, and menstrual irregularities.

The following adverse reactions have been reported in male and female adolescents: premature closure of bony epiphyses with termination of growth, and precocious puberty.

Because these reactions are reported voluntarily from a population of uncertain size and may include abuse of other agents, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

### **9.3 Dependence**

#### *Behaviors Associated with Addiction*

Continued abuse of testosterone and other anabolic steroids, leading to addiction is characterized by the following behaviors:

- Taking greater dosages than prescribed
- Continued drug use despite medical and social problems due to drug use
- Spending significant time to obtain the drug when supplies of the drug are interrupted
- Giving a higher priority to drug use than other obligations
- Having difficulty in discontinuing the drug despite desires and attempts to do so
- Experiencing withdrawal symptoms upon abrupt discontinuation of use

Physical dependence is characterized by withdrawal symptoms after abrupt drug discontinuation or a significant dose reduction of a drug. Individuals taking supratherapeutic doses of testosterone may experience withdrawal symptoms lasting for weeks or months which include depressed mood, major depression, fatigue, craving, restlessness, irritability, anorexia, insomnia, decreased libido and hypogonadotropic hypogonadism.

Drug dependence in individuals using approved doses of testosterone for approved indications has not been documented.

### **10 OVERDOSAGE**

There is a single report of acute overdosage with use of an approved injectable testosterone product: this subject had serum testosterone concentrations of up to 11,400 ng/dL which were implicated in a cerebrovascular accident.

One case of overdose with JATENZO was reported in clinical trials. This patient inadvertently took a higher dose than prescribed (474 mg twice daily, which is 20% higher than the maximum recommended dose). He did not report any adverse reactions associated with the overdose.


Treatment of overdosage consists of discontinuation of JATENZO and appropriate symptomatic and supportive care.

## 11 DESCRIPTION

JATENZO (testosterone undecanoate) for oral use is provided as a gelatin capsule containing testosterone undecanoate, a fatty-acid ester of testosterone. Testosterone undecanoate is a white to off-white yellow crystalline powder. Testosterone, an androgen, is formed by cleavage of the ester side chain of testosterone undecanoate.

Testosterone undecanoate is chemically described as 17 $\beta$ -hydroxyandrost-4-en-3-one undecanoate. It has the empirical formula of C<sub>30</sub>H<sub>48</sub>O<sub>3</sub> and the molecular weight of 456.7. The structural formula for testosterone undecanoate is presented in Figure 1.

**Figure 1: Testosterone Undecanoate**



JATENZO capsules are available in three strengths of 158 mg, 198 mg, and 237 mg.

The 158 mg strength is an opaque red capsule that contains 158 mg of testosterone undecanoate and is imprinted with “158” in white ink. The 198 mg strength is an opaque white capsule that contains 198 mg of testosterone undecanoate and is imprinted with “198” in red ink. The 237 mg strength is an opaque orange capsule that contains 237 mg of testosterone undecanoate and is imprinted with “237” in white ink. All capsule strengths also contain oleic acid, polyoxyl 40 hydrogenated castor oil (Cremophor RH 40), borage seed oil, peppermint oil, and butylated hydroxytoluene as inactive ingredients.

Gelatin capsule shells are composed of the following inactive ingredients: Gelatin, sorbitol, glycerin, and purified water in all strengths, iron oxide red in 158 mg, FD&C Yellow #6 in 158 and 237 mg, and titanium dioxide in 198 and 237 mg capsules.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Endogenous androgens, including testosterone and dihydrotestosterone (DHT), are responsible for the normal growth and development of the male sex organs and for maintenance of secondary sex characteristics. These effects include the growth and maturation of prostate, seminal vesicles, penis and scrotum; the development of male hair distribution, such as facial, pubic, chest and axillary hair; laryngeal enlargement, vocal cord thickening, alterations in body musculature and fat distribution.

Male hypogonadism, a clinical syndrome resulting from insufficient secretion of testosterone, has two main etiologies. Primary hypogonadism is caused by defects of the gonads, such as Klinefelter syndrome or Leydig cell aplasia, whereas secondary hypogonadism (also known as hypogonadotropic hypogonadism) is the failure of the hypothalamus (or pituitary) to produce sufficient gonadotropins (FSH, LH).

### 12.2 Pharmacodynamics

No specific pharmacodynamic studies were conducted using JATENZO.

### 12.3 Pharmacokinetics

#### **Absorption**

JATENZO delivers physiologic amounts of testosterone, producing testosterone

concentrations that approximate normal concentrations seen in healthy men.

JATENZO was taken orally at a starting dose of 237 mg twice per day with meals in a multicenter, open-label, randomized, 2-arm, active-controlled trial in hypogonadal males. The dose was adjusted, as needed, on Days 14 and 56 between a minimum of 158 mg twice per day and a maximum of 396 mg twice per day based on the average plasma testosterone concentration obtained over 24 hours after the morning dose. The average daily NaF-EDTA plasma testosterone concentration was 403 ( $\pm$  128) ng/dL at the end of treatment, where the normal eugonadal range in NaF-EDTA plasma was 252-907 ng/dL in this study. Note that the titration scheme for use in clinical practice is based on serum total testosterone [see *Dosage and Administration (2.2)*].

Table 3 summarizes the pharmacokinetic (PK) parameters for plasma total testosterone in patients completing at least 105 days of JATENZO treatment administered twice daily.


**Table 3: NaF-EDTA Plasma Testosterone  $C_{avg}$  and  $C_{max}$  at Final PK Visit**

PK Parameter		All Doses (N=151)
$C_{avg}$ (ng/dL)	Mean	403
	SD	128
$C_{max}$ (ng/dL)	Mean	1008
	SD	581

PK = pharmacokinetic;  $C_{avg}$  = 24-hour average concentration;  $C_{max}$  = maximum concentration

Figure 2 summarizes the mean plasma total testosterone profile for the patients at the final PK visit.

**Figure 2: Mean ( $\pm$ SEM) Concentration-Time Profile for NaF-EDTA Plasma Total Testosterone in JATENZO Treated Subjects at Final PK Visit**



SEM = standard error of the mean; T = testosterone

When JATENZO was dosed with different breakfasts containing various amounts of fat, the bioavailability with the 30 g fat, 45 g fat, and high-calorie high-fat breakfasts was comparable, but there was a food effect with the 15 g fat breakfast compared to the 30 g fat breakfast. The 15 g fat breakfast had a 25% decrease in testosterone exposure compared to the 30 g fat breakfast.

### Distribution

Circulating testosterone is primarily bound in serum to sex hormone-binding globulin (SHBG) and albumin. Approximately 40% of testosterone in plasma is bound to SHBG, 2% remains unbound (free) and the rest is loosely bound to albumin and other proteins.

### Metabolism

The androgenic activity of testosterone undecanoate occurs after the ester bond linking the testosterone to the undecanoic acid is cleaved by endogenous non-specific

esterases. Undecanoic acid is metabolized like all fatty acids via the beta-oxidation pathway.

Testosterone is metabolized to various 17-keto steroids through two different pathways. The major active metabolites of testosterone are dihydrotestosterone (DHT) and estradiol.

### Drug Interaction Studies

#### *Clinical Pharmacology Studies*

Concomitant administration of JATENZO at 237 mg dose with midazolam (CYP3A4 substrate), digoxin (P-gp substrate), and rosuvastatin (BCRP substrate) had no clinically meaningful effect on the concentration of these substrates.

The potential for JATENZO doses higher than 237 mg to inhibit CYP3A4, P-gp and BCRP is unknown.

### **Excretion**

About 90% of a dose of testosterone given intramuscularly is excreted in the urine as glucuronic and sulfuric acid conjugates of testosterone and its metabolites. About 6% of a dose is excreted in the feces, mostly in the unconjugated form. Inactivation of testosterone occurs primarily in the liver.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

Testosterone has been tested by subcutaneous injection and implantation in mice and rats. In mice, the implant induced cervical-uterine tumors, which metastasized in some cases. There is suggestive evidence that injection of testosterone into some strains of female mice increases their susceptibility to hepatoma. Testosterone is also known to increase the number of tumors and decrease the degree of differentiation of chemically induced carcinomas of the liver in rats.

#### Mutagenesis

Testosterone was negative in the *in vitro* Ames and in the *in vivo* mouse micronucleus assays.

#### Impairment of Fertility

The administration of exogenous testosterone suppresses spermatogenesis in the rat, dog and non-human primates, which was reversible on cessation of the treatment.

### **13.2 Animal Toxicology and/or Pharmacology**

JATENZO has been evaluated in 3- and 9-month repeat-dose oral toxicity studies in male eugonadal dogs. JATENZO caused exaggerated pharmacological effects on androgen-responsive tissues including testes, epididymis, prostate and adrenals at exposures to testosterone or testosterone undecanoate, comparable to the maximum human exposure based on AUC comparisons. Following a 4-week drug-free period, a reduced severity of these findings was observed, suggesting partial reversibility.

In adrenal glands, moderate to severe atrophy, characterized as thinning of the zona fasciculata, was observed with reduced adrenal weights and reduced circulating levels of cortisol in testosterone undecanoate-treated dogs after 3 months of treatment. Following 9-month treatment, there were dose-related decreases in adrenal weights in testosterone undecanoate-treated male dogs and moderate adrenal vacuolation in one testosterone undecanoate-treated male dog. The clinical significance of these adrenal and cortisol findings is unknown.

## 14 CLINICAL STUDIES

### 14.1 Clinical Trials in Hypogonadal Males

The efficacy and safety of JATENZO was evaluated in 166 adult hypogonadal males in an open-label study of approximately 4 months duration (NCT02722278). The study included a Screening Phase, a Treatment Titration Phase, and a Treatment Maintenance Phase.

JATENZO was taken orally at a starting dose of 237 mg twice per day with meals. The dose was adjusted on Days 21 and 56 between a minimum of 158 mg twice per day and a maximum of 396 mg twice per day on the basis of the average testosterone concentration obtained over 24 hours post-morning dose.

The primary endpoint was the percentage of patients with mean plasma total testosterone concentration ( $C_{avg}$ ) over 24-hours within the normal eugonadal range on the final PK visit of the study.

Secondary endpoints were the percentage of patients with a maximum total testosterone concentration ( $C_{max}$ ) above three predetermined limits: less than or equal to 1500 ng/dL, between 1800 and 2500 ng/dL, and greater than 2500 ng/dL.

One hundred and forty-five (87%) of the 166 hypogonadal men who received JATENZO had a mean total testosterone concentration ( $C_{avg}$ ) within the normal eugonadal range at the end of treatment.

The percentage of patients who received JATENZO and had  $C_{max}$  less than or equal to 1500 ng/dL, between 1800 and 2500 ng/dL, and greater than 2500 ng/dL at the final PK visit were 83%, 3%, and 3%, respectively. Note that the testosterone concentrations were not measured in serum but the effects of different sample preparation conditions were accounted for in data analysis of the results shown here. The titration scheme for use in clinical practice is based on serum total testosterone [see *Dosage and Administration* (2.2)].

## 16 HOW SUPPLIED/STORAGE AND HANDLING

JATENZO (testosterone undecanoate) capsules are available in three strengths of 158 mg, 198 mg, and 237 mg. Capsules are packaged as 120 units in wide-mouth, round, white HDPE bottles with white, polypropylene, child resistant caps and induction-sealed liner.

158 mg capsules are opaque red capsules imprinted with "158" in white ink and are supplied in bottles: NDC 69087-158-12.

198 mg capsules are opaque white capsules imprinted with "198" in red ink and are supplied in bottles: NDC 69087-198-12.

237 mg capsules are opaque orange capsules imprinted with "237" in white ink and are supplied in bottles: NDC 69087-237-12.

### Keep JATENZO out of reach of children.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F). Avoid exposing the capsules to moisture (store in a dry place).

## 17 PATIENT COUNSELING INFORMATION

### Advise the patient to read the FDA-approved patient labeling (Medication Guide).

#### Venous Thromboembolism

- Inform patients that JATENZO can cause venous thromboembolism. Advise patients of the signs and symptoms of venous thromboembolism, which may include the following: lower limb pain, edema, or erythema; and dyspnea or chest pain. Advise

patients to promptly report the signs and symptoms of venous thromboembolism, discontinue use of JATENZO, and seek urgent medical care.

### **Increased Blood Pressure**

- Inform patients that JATENZO can increase BP which can increase cardiovascular risk over time.
- Instruct patients about the importance of monitoring BP periodically while on JATENZO. If BP increases while on JATENZO, antihypertensive medications may need to be started, added, or adjusted to control BP, or JATENZO may need to be discontinued.

### **Other Adverse Reactions**

Inform patients that treatment with androgens may lead to adverse reactions which include:

- Changes in urinary habits related to effects on prostate size, such as increased urination at night, hesitancy, frequency, urinary urgency, having a urine accident, being unable to pass urine and weak urine flow
- Breathing disturbances that may reflect obstructive sleep apnea, including those associated with sleep, or excessive daytime sleepiness
- Too frequent or persistent erections of the penis
- Ankle swelling that may reflect peripheral edema
- Red blood cell count increase
- PSA increase
- Nausea and vomiting

Instruct patients to report any changes in their state of health, such as changes in urinary habits, breathing, sleep, and mood including new onset or worsening of depression, or suicidal ideation.

Keep JATENZO out of the reach of children.

Distributed by:

Tolmar, Inc. Fort Collins, CO 80526

133462 Rev. 1 09/2025

<b>MEDICATION GUIDE</b> <b>JATENZO® (juh-TEN-zoh)</b> <b>(testosterone undecanoate) capsules, for oral use CIII</b>
---

<b>What is JATENZO?</b>
-------------------------

JATENZO is a prescription medicine that contains testosterone. JATENZO is used to treat adult men who have low or no testosterone due to certain medical conditions. It is not known if JATENZO is safe or effective to treat men who have low testosterone due to aging.
---

It is not known if JATENZO is safe or effective in children younger than 18 years old. Improper use of JATENZO may affect bone growth in children.
--

JATENZO is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your JATENZO in a safe place to protect it. Never give your JATENZO to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.
---

JATENZO is not meant for use by women.
--

<b>Do not take JATENZO if you:</b>
------------------------------------

- |  |
|--|
| <ul style="list-style-type: none"><li>• have breast cancer.</li><li>• have or might have prostate cancer.</li><li>• are a woman who is pregnant. JATENZO may harm your unborn baby.</li><li>• are allergic to JATENZO or any ingredients in JATENZO. See the end of this Medication Guide for a complete list of ingredients in JATENZO.</li></ul> |
|--|

<b>Before you take JATENZO, tell your healthcare provider about all of your</b>
---

**medical conditions, including if you:**

- have high blood pressure or are treated for high blood pressure.
- have heart problems.
- have high red blood cell count (hematocrit) or high hemoglobin laboratory value.
- have urinary problems due to an enlarged prostate.
- have liver or kidney problems.
- have a history of mental health illness including suicidal thoughts or actions, depression, anxiety or mood disorder.
- have problems breathing while you sleep (sleep apnea).

**Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.**

Using JATENZO with certain other medicines can affect each other. Especially, tell your healthcare provider if you take:

- insulin
- medicines that decrease blood clotting (blood thinners)
- corticosteroids.
- medicines that increase blood pressure such as some cold medicines and pain medicines.

Know the medicines you take. Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

**How should I take JATENZO?**

- Take JATENZO exactly as your healthcare provider tells you take it.
- Take JATENZO by mouth two times daily. Take 1 time in the morning and take 1 time in the evening.
- Take JATENZO with food.
- Your healthcare provider may change your JATENZO dose. Do not change your JATENZO dose without talking to your healthcare provider.

**What are the possible side effects of JATENZO?****JATENZO may cause serious side effects including:**

- **Increase in blood pressure.**
  - JATENZO can increase your blood pressure, which can increase your risk of having a heart attack or stroke over time. Your risk may be greater if you already have heart disease or if you have had a heart attack or stroke.
  - You may need to start new blood pressure medicines or have your medicines adjusted to control your blood pressure while on JATENZO.
  - If your blood pressure cannot be controlled, JATENZO may need to be stopped.
  - Your healthcare provider should check your blood pressure while you use JATENZO.
- **Increase in red blood cell count (hematocrit) or hemoglobin.**
  - JATENZO increases red blood cell counts in some patients. High red blood cell counts increase the risk of blood clots, strokes, and heart attacks.
  - You may need to stop JATENZO if your red blood cell count increases.
  - Your healthcare provider should check your red blood cell count and hemoglobin while you use JATENZO.
- **If you already have an enlarged prostate, your signs and symptoms may worsen while using JATENZO.**

These may include:

  - increased urination at night
  - trouble starting your urine stream
  - urinating many times during the day
  - urge to go to the bathroom right away
  - a urine accident
  - inability to pass urine or weak urine flow
- **Increased risk of prostate cancer.** Your healthcare provider should check you

for prostate cancer or any other prostate problems before you start and while you use JATENZO.

- **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
- **Abuse.** Testosterone can be abused, when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects. Your healthcare provider should check you for signs of abuse before and during treatment with JATENZO.
- **In large doses JATENZO may lower your sperm count.**
- **Liver problems.** Symptoms of liver problems may include:
  - nausea or vomiting
  - yellowing of your skin or whites of your eyes
  - dark urine
  - pain on the right side of your stomach area (abdominal pain)
- **Swelling of your ankles, feet, or body (edema), with or without heart failure.**
- **Enlarged or painful breasts.**
- **Breathing problems while you sleep (sleep apnea).**
- **Changes in mood.** Talk to your healthcare provider if you have changes in mood or behavior including, new or worsening depression, or suicidal thoughts.

**Call your healthcare provider right away if you have any of the serious side effects listed above.**

**The most common side effects of JATENZO include:**

- increased red blood cell count
- diarrhea
- heart burn
- burping
- foot, leg and ankle swelling
- nausea
- enlarged prostate gland
- high blood pressure
- headache

**Other side effects include** more erections than are normal for you or erections that last a long time.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of JATENZO. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store JATENZO?**

- Store JATENZO at room temperature between 68°F to 77°F (20°C to 25°C).
- Store JATENZO in a dry place.

**Keep JATENZO and all medicines out of the reach of children.**

**General information about the safe and effective use of JATENZO**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use JATENZO for a condition for which it was not prescribed. Do not give JATENZO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about JATENZO that is written for health professionals.

**What are the ingredients in JATENZO?**

**Active ingredient:** testosterone undecanoate

**Inactive ingredients:** oleic acid, polyoxyl 40 hydrogenated castor oil (Cremophor RH

40), borage seed oil, peppermint oil, and butylated hydroxytoluene. The ingredients of the soft gelatin capsule shells are gelatin, sorbitol, glycerin, purified water, iron oxide red, FD&C Yellow #6, and titanium dioxide.

Distributed by:  
Tolmar, Inc. Fort Collins, CO 80526

For more information, go to [www.JATENZO.com](http://www.JATENZO.com) or call 1-844-4TO-LMAR.

This Medication Guide has been approved by the U.S. Food and Drug Administration Issued: 09/2025

## PRINCIPAL DISPLAY PANEL

### 158 mg Bottle Label



### 198 mg Bottle Label



### 237 mg Bottle Label



### 237 mg Bottle Sample Label



## JATENZO

testosterone undecanoate capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69087-158
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CIII

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TESTOSTERONE UNDECANOATE (UNII: H16A5VCT9C) (TESTOSTERONE - UNII:3XMK78S47O)	TESTOSTERONE UNDECANOATE	158 mg

### Inactive Ingredients

Ingredient Name	Strength
OLEIC ACID (UNII: 2UMI9U37CP)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
BORAGE OIL (UNII: F8XAG1755S)	
PEPPERMINT OIL (UNII: AV092KU4JH)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
GELATIN (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
GLYCERIN (UNII: PDC6A3C00X)	
WATER (UNII: 059QF0K00R)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

### Product Characteristics

<b>Color</b>	red (red (opaque))	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	158
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69087-158-12	12 in 1 BOX	08/01/2019	
1		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA206089	08/01/2019	

## JATENZO

testosterone undecanoate capsule, liquid filled

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:69087-198
Route of Administration	ORAL	DEA Schedule	CIII

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TESTOSTERONE UNDECANOATE (UNII: H16A5VCT9C) (TESTOSTERONE - UNII:3XMK78S470)	TESTOSTERONE UNDECANOATE	198 mg

Inactive Ingredients	
Ingredient Name	Strength
OLEIC ACID (UNII: 2UMI9U37CP)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
BORAGE OIL (UNII: F8XAG1755S)	
PEPPERMINT OIL (UNII: AV092KU4JH)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
GELATIN (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
GLYCERIN (UNII: PDC6A3C00X)	
WATER (UNII: 059QF0K00R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white (white (opaque))	Score	no score
Shape	CAPSULE (CAPSULE)	Size	23mm
Flavor		Imprint Code	198
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69087-198-12	12 in 1 BOX	08/01/2019	
1		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA206089	08/01/2019	

## JATENZO

testosterone undecanoate capsule, liquid filled

Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:69087-237	
Route of Administration	ORAL	DEA Schedule	CIII	
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
TESTOSTERONE UNDECANOATE (UNII: H16A5VCT9C) (TESTOSTERONE - UNII:3XMK78S470)		TESTOSTERONE UNDECANOATE	237 mg	
Inactive Ingredients				
Ingredient Name			Strength	
OLEIC ACID (UNII: 2UMI9U37CP)				
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)				
BORAGE OIL (UNII: F8XAG1755S)				
PEPPERMINT OIL (UNII: AV092KU4JH)				
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)				
GELATIN (UNII: 2G86QN327L)				
SORBITOL (UNII: 506T60A25R)				
GLYCERIN (UNII: PDC6A3C00X)				
WATER (UNII: 059QF0K00R)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	orange (orange (opaque))	Score	no score	
Shape	CAPSULE (CAPSULE)	Size	26mm	
Flavor		Imprint Code	237	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69087-237-12	12 in 1 BOX	08/01/2019	
1		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:69087-237-28	48 in 1 BOX	06/09/2025	
2		28 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA206089	08/01/2019		

**Labeler** - TOLMAR Inc. (791156578)