

HEALTHY ACCENTS MULTI SYMPTOM COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl suspension
DZA Brands LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DZA Brands, LLC Multi-Symptom Cold Drug Facts

Active ingredients (in each 5 mL = 1 teaspoon)

Acetaminophen 160 mg

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

Purpose

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
- minor aches and pains
- headache
- sore throat
- cough
- stuffy nose
- sneezing and runny nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Liver warning)
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4-6 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 36 lbs	under 4 years	do not use
36-47 lbs	4-5 years	do not use unless directed by a doctor
48-95 lbs	6-11 years	2 tsp or 10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each teaspoon contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, tribasic sodium phosphate, xanthan gum

Questions or comments?

1-866-322-2439

Principal Display Panel

children's

pain relief plus

multi-symptom cold

Fever Reducer - Pain Reliever - Antihistamine

Cough Suppressant - Nasal Decongestant

fever & sore throat (acetaminophen)

cough (dextromethorphan HBr)

stuffy nose (phenylephrine HCl)

sneezing & runny nose (chlorpheniramine maleate)

Compare to Children's Tylenol® Plus Multi-Symptom Cold active ingredients

grape flavor

alcohol free - oral suspension

healthy accents™

Children's
pain relief plus
multi-symptom
cold

MDC 55316-903-26

healthy accents™

Children's
pain relief plus
multi-symptom
cold

Fever Reducer • Pain Reliever • Antihistamine
Cough Suppressant • Nasal Decongestant

fever & sore throat
(acetaminophen)
cough
(dextromethorphan HBr)
stuffy nose
(phenylephrine HCl)
sneezing & runny nose
(chlorpheniramine maleate)

Compare to
Children's Tylenol® Plus
Multi-Symptom Cold
active ingredients



grape flavor



4 FL OZ (118ml)

alcohol free • oral suspension



7 25439 97107 2

LOT NO.

EXP.

: 10326 HA C2

Drug Facts

Active ingredients (in each 5 mL = 1 teaspoon)	Purpose
Acetaminophen 160 mg.....	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg.....	Antihistamine
Dextromethorphan HBr 5 mg.....	Cough suppressant
Phenylephrine HCl 2.5 mg.....	Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - stuffy nose
 - sneezing and runny nose
 - cough
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use


- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- persistent or chronic cough such as occurs with asthma
- diabetes
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a breathing problem such as chronic bronchitis



Multi-Symptom Cold Carton Image 1



Drug Facts (continued)

Ask a doctor or pharmacist before use if your child is

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Liver warning)
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4-6 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 36 lbs	under 4 years	do not use
36-47 lbs	4-5 years	do not use unless directed by a doctor
48-95 lbs	6-11 years	2 tsp or 10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Drug Facts (continued)

Other information

- each teaspoon contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, tribasic sodium phosphate, xanthan gum

Questions or comments?
1-866-322-2439

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Distributed by: DZA Brands, LLC
2110 Executive Drive
Salisbury, NC 28147

For product questions or concerns contact us at 1-866-322-2439. Please include UPC number and code from package.



Multi-Symptom Cold Carton Image 1

Multi-Symptom Cold Carton Image 2

HEALTHY ACCENTS MULTI SYMPTOM COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55316-903
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE (UNII: WAT0DDB505)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	PURPLE (opaque)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:55316-903-26	1 in 1 CARTON	06/06/2011	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/06/2011	

Labeler - DZA Brands LLC (090322194)

Revised: 12/2018

DZA Brands LLC