ANTI-DIARRHEAL- loperamide hydrochloride capsule, liquid filled CHAIN DRUG MARKETING ASSOCIATION INC.

Loperamide HCI Capsules - Quality Choice

Active ingredient (in each capsule)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCI

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- in children under 12 years of age

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
- not for use in children under 12 years of age
- adults and children 12 years and over: 2 capsules after the first loose stool; 1 capsule after each subsequent loose stool; but no more than 4 capsules in 24 hours

Other information

- store at 20°-25°C (68°-77°F). Protect from Light.
- avoid excessive heat above 40°C (104°F)
- do not use if carton or blister unit is open or torn
- see side panel for lot number and expiration date

Inactive ingredients

butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water

Questions or comments?

call toll free **1-888-235-2466**

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium $\ensuremath{\mathbb{R}}$ A-D

L0000568

Lot #

Expiration Date


R0322

Distributed by C.D.M.A., Inc.© 43157 W 9 Mile Rd Novi. MI 48375

www.qualitychoice.com

Questions: 248-449-9300

Principal Display Panel



Active ingredient (in each capsule) Loperamide HCI 2 mg.	Purpose Anti-diarrhea
Use controls symptoms of diarrhea, including Trave	lers' Diarrhea
Warnings Allergy alert: Do not use if you have ever had a rash o Heart alert: Taking more than directed can cause serio	
Do not use I if you have bloody or black stool I in o	children under 12 years of age
Ask a doctor before use if you have fever muc: a history of at	us in the stool a history of liver disease prormal heart rhythm
Ask a doctor or pharmacist before use if you are tak certain prescription drugs.	ing a prescription drug. Loperamide may interact with
When using this product tiredness, drowsiness, or dizzine	ss may occur. Be careful when driving or operating machinery
Stop use and ask a doctor if symptoms get worse you get abdominal sw	 diarrhea lasts for more than 2 days elling or bulging. These may be signs of a serious condition.
If pregnant or breast-feeding, ask a health profession Keep out of reach of children. In case of overdose, get	al before use. medical help or contact a Poison Control Center right away
Directions drink plenty of clear fluids to help prevent dehydra not for use in children under 12 years of age adults and children 12 years and over: 2 capsules after th but no more than 4 capsules in 24 hours	tion caused by diarrhea. he first loose stool; 1 capsule after each subsequent loose stoo
Other information store at 20°-25°C (68°-77°F). Protect from Light. avoid excessive heat above 40°C (104°F)	 do not use if carton or blister unit is open or torn see side panel for lot number and expiration date
Inactive ingredients butylated hydroxyanisole, e polyoxyl 40 hydrogenated castor oil, purified water	dible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate,

QC ®

QUALITY CHOICE

NDC 63868-876-24

*Compare to the Active Ingredient in Imodium[®] A-D

Anti-Diarrheal

Loperamide HCI Soft Gelatin Capsules, 2 mg

Controls the Symptoms of Diarrhea

Suitable for Adults and Children

12 Years and Over

[†]Each Liquid-Filled Capsule Contains 2 mg Loperamide HCI

24 Softgels [†]

psule, liquid filled							
Product Information							
HUMAN OTC DRUG	Item Code (Source)	NDC:63868-876					
ORAL							
Active Ingredient/Active Moiety							
	ORAL	HUMAN OTC DRUG Item Code (Source)					

Pa	ackaging								
	ontains								
	ontains				Imprint Code			LFZ	
	avor							LP2	
			Size			10 score			
0	lor		blue		Score			no score	
P	roduct Char	acteris	tics						
D	roduct Char	actoris	tics						
N	ATER (UNII: 0590	QF0KO0R)							
) (DLYOXYL 40 HY	DROGENA	ATED CASTOR O	DIL (UNII	: 7YC686GQ8F)				
	YCERYL CAPRY								
	· ·		· ·)					
βL	YCERIN (UNII: P	DC6A3C0C	DX)						
GE	ELATIN (UNII: 2G	86QN327L	.)						
= C	&C BLUE NO.	L (UNII: H3	R47K3TBD)						
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)									-
Ingredient Name									
r	active Ingre	edients							
ЛГ	III:6X9OC3H4II)					HY	DROCHLORID	E	
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE -						PERAMIDE		2 mg	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC. (011920774)

Registrant - Bionpharma Inc. (079637826)

Establishment								
Name	Address	ID/FEI	Business Operations					
Patheon Softgels Inc.		002193829	manufacture(63868-876)					

Revised: 12/2022