

CLINDAMYCIN PHOSPHATE AND BENZOYL PEROXIDE- clindamycin phosphate and benzoyl peroxide gel
Oceanside Pharmaceuticals

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Clindamycin Phosphate and Benzoyl Peroxide Gel safely and effectively. See full prescribing information for Clindamycin Phosphate and Benzoyl Peroxide Gel.

Clindamycin Phosphate and Benzoyl Peroxide Gel, 1.2%/2.5%, for topical use
Initial U.S. Approval: 2000

----- **INDICATIONS AND USAGE** -----

Clindamycin Phosphate and Benzoyl Peroxide Gel is a lincosamide antibiotic and benzoyl peroxide indicated for the topical treatment of acne vulgaris. (1)

----- **DOSAGE AND ADMINISTRATION** -----

- Apply a pea-sized amount of Clindamycin Phosphate and Benzoyl Peroxide Gel to the face once daily. (2)
- Not for oral, ophthalmic, or intravaginal use. (2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Clindamycin Phosphate and Benzoyl Peroxide Gel contains clindamycin phosphate 1.2% and benzoyl peroxide 2.5% in a topical gel in 50 gram pumps. (3)

----- **CONTRAINDICATIONS** -----

Clindamycin Phosphate and Benzoyl Peroxide Gel is contraindicated in:

- Patients who have demonstrated hypersensitivity (e.g., anaphylaxis) to clindamycin, benzoyl peroxide, any components of the formulation, or lincomycin. (4)
- Patients with a history of regional enteritis, ulcerative colitis, or antibiotic-associated colitis. (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Colitis: Orally and parenterally administered clindamycin has been associated with severe colitis, which may result in death. Diarrhea, bloody diarrhea, and colitis (including pseudomembranous colitis) have been reported with the use of topical and systemic clindamycin. Clindamycin Phosphate and Benzoyl Peroxide Gel should be discontinued if significant diarrhea occurs. (5.1)
- Ultraviolet Light and Environmental Exposure: Minimize sun exposure following drug application. (5.2)

----- **ADVERSE REACTIONS** -----

The following selected adverse reactions occurred in less than 0.2% of patients: application site pain (0.1%); application site exfoliation (0.1%); and application site irritation (0.1%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Valeant Pharmaceuticals North America LLC at 1-800-321-4576 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

----- **DRUG INTERACTIONS** -----

- Clindamycin Phosphate and Benzoyl Peroxide Gel should not be used in combination with erythromycin-containing products because of its clindamycin component. (7.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 11/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

4.1 Hypersensitivity

4.2 Colitis/Enteritis

5 WARNINGS AND PRECAUTIONS

5.1 Colitis

5.2 Ultraviolet Light and Environmental Exposure

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Erythromycin

7.2 Neuromuscular Blocking Agents

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanisms of Action

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Dispensing Instructions for the Pharmacist

16.3 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Clindamycin Phosphate and Benzoyl Peroxide Gel is indicated for the topical treatment of acne vulgaris in patients 12 years or older.

2 DOSAGE AND ADMINISTRATION

Apply a pea-sized amount of Clindamycin Phosphate and Benzoyl Peroxide Gel to the face once daily.

Use of Clindamycin Phosphate and Benzoyl Peroxide Gel beyond 12 weeks has not been evaluated.

Concomitant topical acne therapy should be used with caution because a possible cumulative irritancy effect may occur, especially with the use of peeling, desquamating, or abrasive agents.

Clindamycin Phosphate and Benzoyl Peroxide Gel is not for oral, ophthalmic, or intravaginal use.

3 DOSAGE FORMS AND STRENGTHS

Gel, 1.2%/2.5%

Each gram of Clindamycin Phosphate and Benzoyl Peroxide Gel contains 10 mg (1%) clindamycin as phosphate, and 25 mg (2.5%) benzoyl peroxide in a white to off-white, opaque, smooth gel.

4 CONTRAINDICATIONS

4.1 Hypersensitivity

Clindamycin Phosphate and Benzoyl Peroxide Gel is contraindicated in those individuals who have shown hypersensitivity to clindamycin, benzoyl peroxide, any components of the formulation, or lincomycin. Anaphylaxis, as well as allergic reactions leading to hospitalization, has been reported in postmarketing use with Clindamycin Phosphate and Benzoyl Peroxide Gel [see *Adverse Reactions (6.2)*].

4.2 Colitis/Enteritis

Clindamycin Phosphate and Benzoyl Peroxide Gel is contraindicated in patients with a history of regional enteritis, ulcerative colitis, or antibiotic-associated colitis [see *Warnings and Precautions (5.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Colitis

Systemic absorption of clindamycin has been demonstrated following topical use of clindamycin. Diarrhea, bloody diarrhea, and colitis (including pseudomembranous colitis) have been reported with the use of topical and systemic clindamycin. When significant diarrhea occurs, Clindamycin Phosphate and Benzoyl Peroxide Gel should be discontinued.

Severe colitis has occurred following oral and parenteral administration of clindamycin with an onset of up to several weeks following cessation of therapy. Antiperistaltic agents such as opiates and diphenoxylate with atropine may prolong and/or worsen severe colitis. Severe colitis may result in death.

Studies indicate toxin(s) produced by Clostridia is one primary cause of antibiotic-associated colitis. The colitis is usually characterized by severe persistent diarrhea and severe abdominal cramps and may be associated with the passage of blood and mucus. Stool cultures for *Clostridium difficile* and stool assay for *C. difficile* toxin may be helpful diagnostically.

5.2 Ultraviolet Light and Environmental Exposure

Minimize sun exposure including use of tanning beds or sun lamps following drug application [see *Nonclinical Toxicology (13.1)*].

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under prescribed conditions, adverse reaction rates observed in the clinical trial may not reflect the rates observed in practice. Because clinical trials are also conducted under widely varying conditions, adverse reactions observed in the clinical trials of a drug cannot always be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The following selected adverse reactions occurred in less than 0.2% of patients treated with Clindamycin Phosphate and Benzoyl Peroxide Gel: application site pain (0.1%); application site exfoliation (0.1%); and application site irritation (0.1%).

During clinical trials, subjects were assessed for local cutaneous signs and symptoms of erythema, scaling, itching, burning and stinging. Most local skin reactions increased and peaked around Week 4 and continually decreased over time reaching near baseline levels by Week 12. The percentage of subjects that had symptoms present before treatment, the maximum value recorded during treatment, and the percent with symptoms present at Week 12 are shown in Table 1.

Table 1: Local Skin Reactions - Percent of Subjects with Symptoms Present. Combined Results from the Two Phase 3 Trials (N = 773)

	Before Treatment (Baseline)			Maximum During Treatment			End of Treatment (Week 12)		
	Mild	Mod.*	Severe	Mild	Mod.*	Severe	Mild	Mod.*	Severe
Erythema	22	4	0	25	5	< 1	15	2	0
Scaling	8	< 1	0	18	3	0	8	1	0
Itching	10	2	0	15	2	0	6	< 1	0
Burning	3	< 1	0	8	2	0	2	< 1	0
Stinging	2	< 1	0	6	1	0	1	< 1	0

* Mod. = Moderate

6.2 Postmarketing Experience

Anaphylaxis, as well as allergic reactions leading to hospitalizations, has been reported in postmarketing use of products containing clindamycin/benzoyl peroxide.

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

7 DRUG INTERACTIONS

7.1 Erythromycin

Clindamycin Phosphate and Benzoyl Peroxide Gel should not be used in combination with topical or oral erythromycin-containing products due to its clindamycin component. *In vitro* studies have shown antagonism between erythromycin and clindamycin. The clinical significance of this *in vitro* antagonism is not known.

7.2 Neuromuscular Blocking Agents

Clindamycin has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. Therefore, Clindamycin Phosphate and Benzoyl Peroxide Gel should be used with caution in patients receiving such agents.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C.

There are no adequate and well-controlled studies in pregnant women treated with Clindamycin Phosphate and Benzoyl Peroxide Gel. Clindamycin Phosphate and Benzoyl Peroxide Gel should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Animal reproductive/developmental toxicity studies have not been conducted with Clindamycin Phosphate and Benzoyl Peroxide Gel or benzoyl peroxide. Developmental toxicity studies of clindamycin performed in rats and mice using oral doses of up to 600 mg/kg/day (240 and 120 times

amount of clindamycin in the highest recommended adult human dose based on mg/m², respectively) or subcutaneous doses of up to 200 mg/kg/day (80 and 40 times the amount of clindamycin in the highest recommended adult human dose based on mg/m², respectively) revealed no evidence of teratogenicity.

8.3 Nursing Mothers

It is not known whether clindamycin is excreted in human milk after topical application of Clindamycin Phosphate and Benzoyl Peroxide Gel. However, orally and parenterally administered clindamycin has been reported to appear in breast milk. Because of the potential for serious adverse reactions in nursing infants, a decision should be made whether to use Clindamycin Phosphate and Benzoyl Peroxide Gel while nursing, taking into account the importance of the drug to the mother.

8.4 Pediatric Use

Safety and effectiveness of Clindamycin Phosphate and Benzoyl Peroxide Gel in pediatric patients under the age of 12 have not been evaluated.

8.5 Geriatric Use


Clinical trials of Clindamycin Phosphate and Benzoyl Peroxide Gel did not include sufficient numbers of subjects aged 65 and older to determine whether they respond differently from younger subjects.

11 DESCRIPTION

Clindamycin Phosphate and Benzoyl Peroxide Gel, 1.2%/2.5% is a combination product with two active ingredients in a white to off-white, opaque, smooth, aqueous gel formulation intended for topical use. Clindamycin phosphate is a water-soluble ester of the semi-synthetic antibiotic produced by a 7(S)-chloro-substitution of the 7(R)-hydroxyl group of the parent antibiotic lincomycin.

The chemical name for clindamycin phosphate is Methyl 7-chloro-6,7,8-trideoxy-6-(1-methyl-trans-4-propyl-L-2-pyrrolidinecarboxamido)-1-thio-L-threo- α -D-galacto-octopyranoside 2-(dihydrogen phosphate). The structural formula for clindamycin phosphate is represented below:


Clindamycin phosphate:



Molecular Formula: C₁₈H₃₄ClN₂O₈PS Molecular Weight: 504.97

Benzoyl peroxide is an antibacterial and keratolytic agent. The structural formula for benzoyl peroxide is represented below:

Benzoyl peroxide:



Molecular Formula: C₁₄H₁₀O₄ Molecular Weight: 242.23

Clindamycin Phosphate and Benzoyl Peroxide Gel contains the following inactive ingredients: purified water, carbomer 980, propylene glycol, and potassium hydroxide. Each gram of Clindamycin Phosphate and Benzoyl Peroxide Gel contains 1.2% of clindamycin phosphate which is equivalent to 1% clindamycin.

12 CLINICAL PHARMACOLOGY

12.1 Mechanisms of Action

Clindamycin: Clindamycin is a lincosamide antibacterial [see *Clinical Pharmacology (12.4)*].

Benzoyl Peroxide: Benzoyl peroxide is an oxidizing agent with bacteriocidal and keratolytic effects but the precise mechanism of action is unknown.

12.3 Pharmacokinetics

The systemic absorption of clindamycin was investigated in an open-label, multiple-dose trial in 16 adult subjects with moderate to severe acne vulgaris treated with 1 gram of Clindamycin Phosphate and Benzoyl Peroxide Gel applied to the face once daily for 30 days. Twelve subjects (75%) had at least one quantifiable clindamycin plasma concentration above the lower limit of quantification (LOQ = 0.5 ng/mL) on Day 1 or Day 30. On Day 1, the mean (\pm standard deviation) peak plasma concentrations (C_{max}) was 0.78 ± 0.22 ng/mL (n=9 with measurable concentrations), and the mean AUC_{0-t} was 5.29 ± 0.81 h.ng/mL (n=4). On Day 30, the mean C_{max} was 1.22 ± 0.88 ng/mL (n=10), and the mean AUC_{0-t} was 8.42 ± 6.01 h.ng/mL (n=6). Clindamycin plasma concentrations were below LOQ in all subjects at 24 hours post-dose on the three tested days (Day 1, 15, and 30).

Benzoyl peroxide has been shown to be absorbed by the skin where it is converted to benzoic acid.

12.4 Microbiology

Clindamycin binds to the 50S ribosomal subunits of susceptible bacteria and prevents elongation of peptide chains by interfering with peptidyl transfer, thereby suppressing bacterial protein synthesis.

Clindamycin and benzoyl peroxide individually have been shown to have *in vitro* activity against *Propionibacterium acnes*, an organism which has been associated with acne vulgaris; however, the clinical significance of this activity against *P. acnes* is not known.

P. acnes resistance to clindamycin has been documented. Resistance to clindamycin is often associated with resistance to erythromycin.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity, mutagenicity and impairment of fertility testing of Clindamycin Phosphate and Benzoyl Peroxide Gel have not been performed.

Benzoyl peroxide has been shown to be a tumor promoter and progression agent in a number of animal

studies. Benzoyl peroxide in acetone at doses of 5 and 10 mg administered topically twice per week for 20 weeks induced skin tumors in transgenic Tg.AC mice. The clinical significance of this is unknown.

Carcinogenicity studies have been conducted with a gel formulation containing 1% clindamycin and 5% benzoyl peroxide. In a 2-year dermal carcinogenicity study in mice, treatment with the gel formulation at doses of 900, 2700, and 15000 mg/kg/day (1.8, 5.4, and 30 times amount of clindamycin and 3.6, 10.8, and 60 times amount of benzoyl peroxide in the highest recommended adult human dose of 2.5 g Clindamycin Phosphate and Benzoyl Peroxide Gel based on mg/m², respectively) did not cause any increase in tumors. However, topical treatment with a different gel formulation containing 1% clindamycin and 5% benzoyl peroxide at doses of 100, 500, and 2000 mg/kg/day caused a dose-dependent increase in the incidence of keratoacanthoma at the treated skin site of male rats in a 2-year dermal carcinogenicity study in rats. In an oral (gavage) carcinogenicity study in rats, treatment with the gel formulation at doses of 300, 900 and 3000 mg/kg/day (1.2, 3.6, and 12 times amount of clindamycin and 2.4, 7.2, and 24 times amount of benzoyl peroxide in the highest recommended adult human dose of 2.5 g Clindamycin Phosphate and Benzoyl Peroxide Gel based on mg/m², respectively) for up to 97 weeks did not cause any increase in tumors. In a 52-week dermal photocarcinogenicity study in hairless mice, (40 weeks of treatment followed by 12 weeks of observation), the median time to onset of skin tumor formation decreased and the number of tumors per mouse increased relative to controls following chronic concurrent topical administration of the higher concentration benzoyl peroxide formulation (5000 and 10000 mg/kg/day, 5 days/week) and exposure to ultraviolet radiation.

Clindamycin phosphate was not genotoxic in the human lymphocyte chromosome aberration assay. Benzoyl peroxide has been found to cause DNA strand breaks in a variety of mammalian cell types, to be mutagenic in *S. typhimurium* tests by some but not all investigators, and to cause sister chromatid exchanges in Chinese hamster ovary cells.

Fertility studies have not been performed with Clindamycin Phosphate and Benzoyl Peroxide Gel or benzoyl peroxide, but fertility and mating ability have been studied with clindamycin. Fertility studies in rats treated orally with up to 300 mg/kg/day of clindamycin (approximately 120 times the amount of clindamycin in the highest recommended adult human dose of 2.5 g Clindamycin Phosphate and Benzoyl Peroxide Gel, based on mg/m²) revealed no effects on fertility or mating ability.

14 CLINICAL STUDIES

The safety and efficacy of once-daily use of Clindamycin Phosphate and Benzoyl Peroxide Gel were assessed in two 12-week, multicenter, randomized, blinded trials in subjects 12 years and older with moderate to severe acne vulgaris. The two trials were identical in design and compared Clindamycin Phosphate and Benzoyl Peroxide Gel to clindamycin in the vehicle gel, benzoyl peroxide in the vehicle gel, and the vehicle gel alone.

The co-primary efficacy variables were:

1. Mean absolute change from baseline at Week 12 in
 1. Inflammatory lesion counts
 2. Non-inflammatory lesion counts
2. Percent of subjects who had a two grade improvement from baseline on an Evaluator's Global Severity (EGS) score.

The EGS scoring scale used in all of the clinical trials for Clindamycin Phosphate and Benzoyl Peroxide Gel is as follows:

Grade	Description
-------	-------------

Clear	Normal, clear skin with no evidence of acne vulgaris
Almost Clear	Rare non-inflammatory lesions present, with rare non-inflamed papules (papules must be resolving and may be hyperpigmented, though not pink-red)
Mild	Some non-inflammatory lesions are present, with few inflammatory lesions (papules/pustules only; no nodulocystic lesions)
Moderate	Non-inflammatory lesions predominate, with multiple inflammatory lesions evident: several to many comedones and papules/pustules, and there may or may not be one small nodulocystic lesion
Severe	Inflammatory lesions are more apparent, many comedones and papules/pustules, there may or may not be a few nodulocystic lesions
Very Severe	Highly inflammatory lesions predominate, variable number of comedones, many papules/pustules and many nodulocystic lesions

The results of Trial 1 at Week 12 are presented in Table 2:

Table 2: Trial 1 Results

Trial 1	Clindamycin Phosphate and Benzoyl Peroxide Gel N = 399	Clindamycin Gel N = 408	Benzoyl Peroxide Gel N = 406	Vehicle Gel N = 201
EGSS Clear or Almost Clear	115 (29%)	84 (21%)	76 (19%)	29 (14%)
2 grade reduction from baseline	131 (33%)	100 (25%)	96 (24%)	38 (19%)
Inflammatory Lesions: Mean absolute change	14.8	12.2	13.0	9.0
Mean percent (%) reduction	55.0%	47.1%	49.3%	34.5%
Non-Inflammatory Lesions: Mean absolute change	22.1	17.9	20.6	13.2
Mean percent (%) reduction	45.3%	38.0%	40.2%	28.6%

The results of Trial 2 at Week 12 are presented in Table 3:

Table 3: Trial 2 Results

Trial 2	Clindamycin Phosphate and Benzoyl Peroxide Gel N = 398	Clindamycin Gel N = 404	Benzoyl Peroxide Gel N = 403	Vehicle Gel N = 194
----------------	---	------------------------------------	---	--------------------------------

EGSS Clear or Almost Clear	113 (28%)	94 (23%)	94 (23%)	21 (11%)
2 grade reduction from baseline	147 (37%)	114 (28%)	114 (28%)	27 (14%)
Inflammatory Lesions:				
Mean absolute change	13.7	11.3	11.2	5.7
Mean percent (%) reduction	54.2%	45.3%	45.7%	23.3%
Non-Inflammatory Lesions:				
Mean absolute change	19.0	14.9	15.2	8.3
Mean percent (%) reduction	41.2%	34.3%	34.5%	19.2%

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Clindamycin Phosphate and Benzoyl Peroxide Gel is supplied as a 50 g pump (NDC 68682-132-50).

16.2 Dispensing Instructions for the Pharmacist

- Dispense Clindamycin Phosphate and Benzoyl Peroxide Gel with a 10-week expiration date.
- Specify "Store at room temperature up to 25°C (77°F). Do not freeze."

16.3 Storage and Handling

- PHARMACIST: Prior to Dispensing: Store in a refrigerator, 2° to 8°C (36° to 46°F).
- PATIENT: Store at room temperature at or below 25°C (77°F).
- Protect from freezing.
- Keep out of reach of children.
- Keep container tightly closed.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information).

- Patients who develop allergic reactions such as severe swelling or shortness of breath should discontinue use and contact their physician immediately.
- Clindamycin Phosphate and Benzoyl Peroxide Gel may cause irritation such as erythema, scaling, itching, or burning, especially when used in combination with other topical acne therapies.
- Excessive or prolonged exposure to sunlight should be limited. To minimize exposure to sunlight, a hat or other clothing should be worn. Sunscreen may also be used.
- Clindamycin Phosphate and Benzoyl Peroxide Gel may bleach hair or colored fabric.

Manufactured for:

Oceanside Pharmaceuticals, a division of Valeant Pharmaceuticals North America LLC
Bridgewater, NJ 08807 USA

By:

Valeant Pharmaceuticals International, Inc.

Laval, Quebec H7L 4A8, Canada

U.S. Patents 8,288,434; 8,663,699; 8,895,070; 9,078,870 and 9,693,988

© Valeant Pharmaceuticals North America LLC

9613800-20002089A

PATIENT INFORMATION

Clindamycin Phosphate and Benzoyl Peroxide Gel, 1.2%/2.5%

IMPORTANT: For use on skin only (topical use). Do not get Clindamycin Phosphate and Benzoyl Peroxide Gel in your mouth, eyes, or vagina, or on your lips.

Read the Patient Information that comes with Clindamycin Phosphate and Benzoyl Peroxide Gel before you start using it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your medical condition or your treatment.

What is Clindamycin Phosphate and Benzoyl Peroxide Gel?

Clindamycin Phosphate and Benzoyl Peroxide Gel is a prescription medicine used on the skin (topical) to treat acne vulgaris in people 12 years and older. Clindamycin Phosphate and Benzoyl Peroxide Gel contains clindamycin phosphate and benzoyl peroxide.

It is not known if Clindamycin Phosphate and Benzoyl Peroxide Gel is safe and effective for use longer than 12 weeks.

It is not known if Clindamycin Phosphate and Benzoyl Peroxide Gel is safe and effective in children under 12 years of age.

Who should not use Clindamycin Phosphate and Benzoyl Peroxide Gel?

Do not use Clindamycin Phosphate and Benzoyl Peroxide Gel if you have:

- Crohn's disease
- ulcerative colitis
- had inflammation of the colon (colitis), or severe diarrhea with past antibiotic use

Talk with your doctor if you are not sure if you have one of these conditions.

What should I tell my doctor before using Clindamycin Phosphate and Benzoyl Peroxide Gel?

Before using Clindamycin Phosphate and Benzoyl Peroxide Gel, tell your doctor about all of your medical conditions, including if you:

- **have any allergies.**
- **have any other medical conditions**
- **are pregnant or planning to become pregnant.** It is not known if Clindamycin Phosphate and Benzoyl Peroxide Gel will harm your unborn baby.
- **are breastfeeding or plan to breastfeed.** It is not known if Clindamycin Phosphate and Benzoyl Peroxide Gel passes into your breast milk. One of the medicines in Clindamycin Phosphate and Benzoyl Peroxide Gel contains clindamycin. Clindamycin when taken by mouth or by injection has been reported to appear in breast milk. You and your doctor should decide whether you will use Clindamycin Phosphate and Benzoyl Peroxide Gel while breastfeeding.

Tell your doctor about all the medicines and skin products you use. Especially tell your doctor if you will have surgery with general anesthesia. One of the medicines in Clindamycin Phosphate and Benzoyl Peroxide Gel (clindamycin) can affect how certain medicines work when used in general anesthesia.

- Clindamycin Phosphate and Benzoyl Peroxide Gel should not be used with products that contain erythromycin.
- Other skin and topical acne products may increase the irritation of your skin when used with

Clindamycin Phosphate and Benzoyl Peroxide Gel.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.


How should I use Clindamycin Phosphate and Benzoyl Peroxide Gel?

- Use Clindamycin Phosphate and Benzoyl Peroxide Gel exactly as prescribed.
- Your doctor will tell you how long to use Clindamycin Phosphate and Benzoyl Peroxide Gel.
- Throw away (discard) any unused Clindamycin Phosphate and Benzoyl Peroxide Gel.

Instructions for applying Clindamycin Phosphate and Benzoyl Peroxide Gel

- Apply Clindamycin Phosphate and Benzoyl Peroxide Gel to your face one time each day as prescribed.
- Before you apply Clindamycin Phosphate and Benzoyl Peroxide Gel, wash your face gently with a mild soap, rinse with warm water, and pat your skin dry.
- To apply Clindamycin Phosphate and Benzoyl Peroxide Gel to your face, use the pump to dispense one pea-sized amount of Clindamycin Phosphate and Benzoyl Peroxide Gel onto your fingertip. See Figure 1. One pea-sized amount of Clindamycin Phosphate and Benzoyl Peroxide Gel should be enough to cover your entire face.

Figure 1



-
1. Dot the one pea-sized amount of Clindamycin Phosphate and Benzoyl Peroxide Gel onto six areas of your face (chin, left cheek, right cheek, nose, left forehead, right forehead). See Figure 2.

Figure 2



1. After applying the Clindamycin Phosphate and Benzoyl Peroxide Gel this way, spread the gel over your face and gently rub it in. It is important to spread the gel over your whole face.
2. Wash your hands with soap and water after applying Clindamycin Phosphate and Benzoyl Peroxide Gel.
3. If your doctor tells you to put Clindamycin Phosphate and Benzoyl Peroxide Gel on other areas of your skin with acne, be sure to ask how much you should use.
4. Do not get Clindamycin Phosphate and Benzoyl Peroxide Gel in your mouth, eyes, or nose, or on your lips. If this occurs, rinse the affected area with warm water and call your doctor right away if the area becomes very red, itchy, tender, or swollen.
5. Do not get Clindamycin Phosphate and Benzoyl Peroxide Gel on cuts or open wounds.
6. Do not use more Clindamycin Phosphate and Benzoyl Peroxide Gel than prescribed.

What should I avoid while using Clindamycin Phosphate and Benzoyl Peroxide Gel?

- Limit your time in sunlight. Avoid using tanning beds or sun lamps. If you have to be in sunlight, wear a wide-brimmed hat or other protective clothing, and a sunscreen with SPF 15 rating or higher. Your doctor can give you more information about why this is important.
- Do not wash your face more than 2 to 3 times a day. Washing your face too often or scrubbing it may make your acne worse.
- Avoid getting Clindamycin Phosphate and Benzoyl Peroxide Gel in your hair or on colored fabric. Clindamycin Phosphate and Benzoyl Peroxide Gel may bleach hair or colored fabric.

What are possible side effects with Clindamycin Phosphate and Benzoyl Peroxide Gel?

Clindamycin Phosphate and Benzoyl Peroxide Gel can cause serious side effects including:

- **Inflammation of the colon (colitis).** Stop using Clindamycin Phosphate and Benzoyl Peroxide Gel and call your doctor right away if you have severe watery diarrhea, or bloody diarrhea.
- **Allergic reactions.** Stop using Clindamycin Phosphate and Benzoyl Peroxide Gel, call your doctor and get help right away if you have any of the following symptoms:
 - severe itching
 - swelling of your face, eyes, lips, tongue or throat
 - trouble breathing

Common side effects with Clindamycin Phosphate and Benzoyl Peroxide Gel include:

- **Skin irritation.** Stop using Clindamycin Phosphate and Benzoyl Peroxide Gel and call your doctor if you have a skin rash or your skin becomes very red, itchy or swollen.

Talk to your doctor about any side effect that bothers you or that does not go away.

These are not all the possible side effects with Clindamycin Phosphate and Benzoyl Peroxide Gel. Ask your doctor or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

You may also report side effects to Valeant Pharmaceuticals North America LLC at 1-800-321-4576.

How should I store Clindamycin Phosphate and Benzoyl Peroxide Gel?

- Store Clindamycin Phosphate and Benzoyl Peroxide Gel at room temperature at or below 25°C (77°F).
- The expiration date of Clindamycin Phosphate and Benzoyl Peroxide Gel is 10 weeks from the date you fill your prescription.
- Safely throw away expired Clindamycin Phosphate and Benzoyl Peroxide Gel.
- Do not freeze.
- Keep the container tightly closed.

Keep Clindamycin Phosphate and Benzoyl Peroxide Gel and all medicines out of reach of children.

General information about Clindamycin Phosphate and Benzoyl Peroxide Gel

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information leaflets. Do not use Clindamycin Phosphate and Benzoyl Peroxide Gel for a condition for which it was not prescribed. **Do not give Clindamycin Phosphate and Benzoyl Peroxide Gel to other people, even if they have the same condition you have. It may harm them.**

This leaflet summarizes the most important information about Clindamycin Phosphate and Benzoyl Peroxide Gel. If you would like more information, talk with your doctor. You can also ask your doctor or pharmacist for information about Clindamycin Phosphate and Benzoyl Peroxide Gel that is written for healthcare professionals.

For more information about Clindamycin Phosphate and Benzoyl Peroxide Gel, call 1-800-321-4576.

What are the ingredients in Clindamycin Phosphate and Benzoyl Peroxide Gel?

Active ingredients: clindamycin phosphate 1.2% and benzoyl peroxide 2.5%

Inactive ingredients: purified water, carbomer 980, propylene glycol, and potassium hydroxide

Manufactured for:

Oceanside Pharmaceuticals, a division of Valeant Pharmaceuticals North America LLC
Bridgewater, NJ 08807 USA

By:

Valeant Pharmaceuticals International, Inc.
Laval, Quebec H7L 4A8, Canada

U.S. Patents 8,288,434; 8,663,699; 8,895,070; 9,078,870 and 9,693,988

© Valeant Pharmaceuticals North America LLC

This Patient Information has been approved by the U.S. Food and Drug Administration.
Revised 11/2017

9613800-20002089A

Package/Label Display Panel - 50 gram Pump Label

NDC 68682-132-50


Rx only

Clindamycin Phosphate and Benzoyl Peroxide Gel, 1.2%/2.5%

FOR TOPICAL USE ONLY

One premixed 50-gram pump dispenser

OCEANSIDE Pharmaceuticals



CLINDAMYCIN PHOSPHATE AND BENZOYL PEROXIDE

clindamycin phosphate and benzoyl peroxide gel

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68682-132
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
Clindamycin Phosphate (UNII: EH6D7113I8) (Clindamycin - UNII:3U02EL437C)		Clindamycin	10 mg in 1 g	
Benzoyl Peroxide (UNII: W9WZN9A0GM) (Benzoyl Peroxide - UNII:W9WZN9A0GM)		Benzoyl Peroxide	25 mg in 1 g	
Inactive Ingredients				
Ingredient Name			Strength	
Water (UNII: 059QF0KO0R)				
CARBOMER HOMO POLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)				
propylene glycol (UNII: 6DC9Q167V3)				
potassium hydroxide (UNII: WZHBC48M4T)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68682-132-50	1 in 1 CARTON	07/02/2018	
1		50 g in 1 BOTTLE, PUMP; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA authorized generic	NDA050819	07/02/2018		

Labeler - Oceanside Pharmaceuticals (832011691)

Establishment

Name	Address	ID/FEI	Business Operations
Valeant Pharmaceuticals International, Inc		245141858	MANUFACTURE(68682-132)

Establishment

Name	Address	ID/FEI	Business Operations
Contract Pharmaceuticals Limited Canada		248761249	MANUFACTURE(68682-132)

Revised: 11/2017

Oceanside Pharmaceuticals