

**SEVERE TUSSIN CF- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution**

**CVS Pharmacy**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**CVS Pharmacy, Inc. Severe Tussin CF Drug Facts**

**Active ingredients (in each 20 mL)**

Acetaminophen, USP 650 mg

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 400 mg

Phenylephrine HCl, USP 10 mg

**Purposes**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- temporarily relieves these symptoms occurring with a cold or flu:
- cough due to minor throat and bronchial irritation
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

### **When using this product**

do not use more than directed

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

**Other information**

- **each 20 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

Compare to the active ingredients in Robitussin<sup>®</sup> Severe Multi-Symptom Cough Cold + Flu

MAXIMUM STRENGTH

Non-Drowsy

Severe

Tussin CF

ACETAMINOPHEN

Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr

Cough suppressant

GUAIFENESIN

Expectorant

PHENYLEPHRINE HCl

Nasal decongestant

Adult Cough & Cold

Relieves:

Cough, sore throat

Body aches, fever

Nasal congestion

Chest congestion

For Ages 12 & Over

Dosage cup provided

Package Contains One Bottle

4 FL OZ (118 mL)



## SEVERE TUSSIN CF

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-565
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SORBITOL (UNII: 506T60A25R)	

**Product Characteristics**

Color	RED	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-565-26	1 in 1 CARTON	06/25/2018	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69842-565-34	1 in 1 CARTON	06/25/2018	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/25/2018	

