

## **MAX STR.NASAL RELIEF SEVERE CONGESTION- oxymetazoline hydrochloride**

**0.05% liquid**

**Ride Aid**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredient**

Oxymetazoline Hydrochloride 0.05%

#### **Purpose**

Nasal decongestant

### **Uses**

- for the temporary relief of nasal congestion due to the common cold, hay fever or other upper respiratory allergies

### **Warnings**

#### **Ask a doctor before use if you have**

- heart disease • high blood pressure • thyroid disease
- diabetes • trouble urinating due to enlarged prostate gland

#### **When using this product**

**Do not use** this product if you have heart disease • high blood pressure • thyroid disease • diabetes • or difficulty in urination due to enlargement of the prostate gland

When using this product • Do not exceed recommended dosage. • This product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge. • The use of this container by more than one person may spread infection. • Do not use for more than 3 days. Use only as directed. • Frequent or prolonged use may cause nasal congestion to recur or worsen.

**Stop use and ask doctor if** symptoms persist, consult a doctor.

**Keep out of reach of children.** If product is swallowed, get medical help or contact a PoisonControl Center right away.

### **Directions**

Before using the first time, remove the protective cap from the tip and prime pump by depressing pump firmly several times.

To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses within any 24-hour period

- children under 6 years of age: consult a doctor

**Inactive ingredients**

Benzalkonium Chloride, Benzyl Alcohol, Camphor, Edetate Disodium, Eucalyptol, Menthol, Microcrystalline Cellulose, Polyethylene Glycol, Povidone, Propylene Glycol, Purified Water, Sodium Carboxymethyl Cellulose, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic.



nasal relief  
**severe  
congestion  
nasal spray**

oxymetazoline hydrochloride 0.05%



\*Compare to the active ingredient in Afrin® No-Drip®

MAXIMUM STRENGTH  
nasal relief  
**severe  
congestion  
nasal spray**

oxymetazoline  
hydrochloride 0.05%

NASAL DECONGESTANT

**12 HOUR**

won't drip from nose  
or down throat

fast, powerful  
congestion relief

colds; allergies

no drip pump spray

1 FL OZ (30 mL)

CAP REMOVAL  
INSTRUCTIONS:

To remove cap squeeze cap at base on opposite sides on the two smooth impressions with fingers. While squeezing, twist cap counterclockwise to remove. Refer to directions on top of cap.

To reapply cap turn cap on pump clockwise until it locks.

DISTRIBUTED BY:  
RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011



ACTUAL SIZE



# nasal relief severe congestion nasal spray

oxymetazoline hydrochloride 0.05%

**DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING**

## Drug Facts

**Active Ingredient** Oxymetazoline hydrochloride 0.05%.....**Purpose** Nasal decongestant

### Uses

For the temporary relief of nasal congestion due to the common cold • hay fever • other upper respiratory allergies.

### Warnings

**Do not use this product if you have** heart disease • high blood pressure • thyroid disease • diabetes • or difficulty in urination due to enlargement of the prostate gland

**When using this product** • Do not exceed recommended dosage. • This product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge. • The use of this container by more than one person may spread infection. • Do not use for more than 3 days. Use only as directed. • Frequent or prolonged use may cause nasal congestion to recur or worsen.

**Stop use and ask doctor** if symptoms persist, consult a doctor.

**Keep out of reach of children. If product is swallowed,** get medical help or contact a PoisonControl Center right away.

### Directions

**Shake well before use.** Before using the first time, remove the protective cap from the tip and prime pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. ▶



# nasal relief severe congestion nasal spray

## Drug Facts (continued)

Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after each use.

- **adults and children 6 to under 12 years of age** (with adult supervision): 2 or 3 sprays in each nostril, not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- **children under 6 years of age:** consult a doctor

## Other information

- store at room temperature

## Inactive ingredients

Benzalkonium Chloride, Benzyl Alcohol, Camphor, Edetate Disodium, Eucalyptol, Menthol, Microcrystalline Cellulose, Polyethylene Glycol, Povidone, Propylene Glycol, Purified Water, Sodium Carboxymethyl Cellulose, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic.

\*This product is not manufactured or distributed by MSD Consumer Care, the distributor of Afrin®.

**IMPORTANT:  
KEEP THIS CARTON  
FOR FUTURE REFERENCE  
ON FULL LABELING**

## MAX STR.NASAL RELIEF SEVERE CONGESTION

oxymetazoline hydrochloride 0.05% liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:118 22-1232
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>Oxymetazoline Hydrochloride</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	Oxymetazoline Hydrochloride	0.05 g in 100 mL
---	-----------------------------	------------------

### Inactive Ingredients

Ingredient Name	Strength
<b>Benzalkonium Chloride</b> (UNII: F5UM2KM3W7)	
<b>Benzyl Alcohol</b> (UNII: LKG8494WBH)	
<b>CAMPHOR (SYNTHETIC)</b> (UNII: 5TJD82A1ET)	
<b>Edetate Disodium</b> (UNII: 7FLD91C86K)	
<b>Eucalyptol</b> (UNII: RV6J6604TK)	
<b>Menthol</b> (UNII: L7T10EIP3A)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOLS</b> (UNII: 3WJQ0SDW1A)	
<b>Povidone</b> (UNII: FZ989GH94E)	
<b>Propylene Glycol</b> (UNII: 6DC9Q167V3)	
<b>Water</b> (UNII: 059QF0KO0R)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM</b> (UNII: K679OBS311)	
<b>SODIUM PHOSPHATE, DIBASIC, ANHYDROUS</b> (UNII: 22ADO53M6F)	
<b>SODIUM PHOSPHATE, MONOBASIC, ANHYDROUS</b> (UNII: KH7I04HPUU)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1232-1	1 in 1 CARTON	05/09/2014	
1		15 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/09/2014	

**Labeler** - Ride Aid (014578892)

**Registrant** - Product Quest Mfg, LLC (927768135)

### Establishment

Name	Address	ID/FEI	Business Operations
Product Quest Mfg, LLC		927768135	manufacture(11822-1232) , label(11822-1232)