

**HAN-PEN- penicillin g potassium powder, for solution**  
**US Vet Inc**

-----  
**Han-Pen (penicillin G potassium)**

**DESCRIPTION**

**0.5 BU Front Panel**

**Penicillin G Potassium**

Antibiotic for drinking water

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For oral use in turkeys only

For Veterinary use only

**Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).**

Keep out of the reach of children.

Restricted drug (California) -Use only as directed

**Active Ingredient:**

0.500 Billion Units of penicillin G potassium

Nonsterile

**Net Weight: 313 grams (11 ounces)**

Approved by FDA under ANADA #200-372

**INDICATIONS AND USAGE**

For the treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

**DOSAGE AND ADMINISTRATION**

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

**Directions**

Combine contents and approximately 1-1/2 pints (710 mL) of water in a glass or plastic container. Stir to dissolve. Allow concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour after preparation.

Automatic Watering Systems - Pour the concentrated solution into a glass or plastic container then add enough water to make 2.6 gallons (9.9 liters) of stock solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water]. The

automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions daily.

Gravity Flow Watering Systems - Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

## **WARNINGS**

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment. Do not use in turkeys producing eggs for human consumption.

## **PRECAUTIONS**

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian.

## **DESCRIPTION**

### **1 BU Front Panel**

#### **Penicillin G Potassium**

Antibiotic for drinking water

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For oral use in turkeys only

For Veterinary use only

**Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).**

Keep out of the reach of children.

Restricted drug (California) -Use only as directed

#### **Active Ingredient:**

1 Billion Units of penicillin G potassium

Nonsterile

**Net Weight: 626 grams (22 ounces)**

Approved by FDA under ANADA #200-372

## **INDICATIONS AND USAGE**

For the treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

## **DOSAGE AND ADMINISTRATION**

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

### **Directions**

Combine contents and approximately 3 pints (1420 mL) of water in a glass or plastic container. Stir to dissolve. Allow concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour after preparation.

Automatic Watering Systems - Pour the concentrated solution into a glass or plastic container then add enough water to make 5.2 gallons (19.7 liters) of stock solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water]. The automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions daily.

Gravity Flow Watering Systems - Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

## **WARNINGS**

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment. Do not use in turkeys producing eggs for human consumption.

## **PRECAUTIONS**

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian.

Manufactured for:

**US VET Inc.**

Liverpool, NY 13088

Rev. 08/2024

## **PACKAGE/LABEL PRINCIPAL DISPLAY PANEL**

NDC 86108-372-01

# Han-Pen™

## (penicillin G potassium)

Antibiotic for drinking water

**CAUTION: Federal (USA) law restricts this drug to use  
by or on the order of a licensed veterinarian.**

For oral use in turkeys only

For Veterinary use only

Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).

Keep out of the reach of children.

Restricted drug (California)- Use only as directed

**Active Ingredient:**

0.500 Billion Units penicillin G potassium  
Nonsterile

**Net Weight: 313 grams (11 ounces)**

Approved by FDA under ANADA # 200-372



NDC 86108-372-01

Han-Pen™

(penicillin G potassium)

Antibiotic for drinking water

### PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 86108-373-01

# Han-Pen™

## (penicillin G potassium)

Antibiotic for drinking water

**CAUTION: Federal (USA) law restricts this drug to use  
by or on the order of a licensed veterinarian.**

For oral use in turkeys only

For Veterinary use only

Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).

Keep out of the reach of children.

Restricted drug (California)- Use only as directed

**Active Ingredient:**

1 Billion Units penicillin G potassium  
Nonsterile

**Net Weight: 626 grams (22 ounces)**

Approved by FDA under ANADA # 200-372



NDC 86108-373-01  
Han-Pen <sup>TM</sup>  
(penicillin G potassium)  
Antibiotic for drinking water

## HAN-PEN

penicillin g potassium powder, for solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:86108-372
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Penicillin G Potassium</b> (UNII: VL775ZTH4C) (Penicillin G - UNII:Q42T66VG0C)	Penicillin G	500000000 [iU] in 313 g

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86108-372-01	313 g in 1 BAG		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200372	06/07/2021	

## HAN-PEN

penicillin g potassium powder, for solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:86108-373
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Penicillin G Potassium</b> (UNII: VL775ZTH4C) (Penicillin G - UNII:Q42T66VG0C)	Penicillin G	1000000000 [iU] in 626 g

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86108-373-01	626 g in 1 BAG		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANADA	ANADA200372		06/07/2021	

**Labeler -** US Vet Inc (117160633)

Establishment			
Name	Address	ID/FEI	Business Operations
GC Hanford Manufacturing Company		002238863	MANUFACTURE

Establishment			
Name	Address	ID/FEI	Business Operations
Inner Mongolia Changsheng Pharmaceutical Co. Ltd.		545265563	API MANUFACTURE

Establishment			
Name	Address	ID/FEI	Business Operations
Fersinsa GB, S.A. de C.V.		814622007	API MANUFACTURE