

BACTOSHIELD CHG- chlorhexidine gluconate 2% solution
Xttrium Laboratories, Inc.

XTT COMP SCJ 4 oz 2% CHG Bactoshield

Drug Facts

Active ingredient

Chlorhexidine gluconate 2% solution

Purpose

Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **patient preoperative skin preparation:** preparation of the patient's skin prior to surgery
- **skin wound and general skin cleansing**

Warnings

For external Use Only

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area
- as a preoperative skin preparation of the head or face

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if

placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.

- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms under running water for 30 seconds. Clean fingernails using a nailstick or similar cleaner.
- scrub for 1.5 minutes with about 8 mL of product with or without a wet brush paying close attention to the nails, cuticles, and skin between the fingers
- rinse thoroughly under running water for 30 seconds
- wash for an additional 1.5 minutes with 8 mL of product and rinse under running water for 30 seconds
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Questions or comments?

call 1-800-587-3721

Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

Packaging**SCJ Professional**

Healthcare

Bactoshield**CHG 2%**

NDC 11084-038-03

Surgical Scrub**Chlorhexidine Gluconate 2% Solution Antiseptic****Non-Sterile Solution****Surgical Scrub****2% CHG****Antimicrobial**

REORDER#

1322-39

SAP # 4000011062

1322-52K(123)

2SCJ04BTLLBLA

deb

SKIN CARE

For Single Use

Net Contents:

4 fl oz (118 mL)



Lift Here

Drug Facts	Purpose
Active ingredient Chlorhexidine gluconate 2% solution	Antiseptic
Uses ■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care ■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease	

Drug Facts (continued) ■ patient preoperative skin preparation: preparation of the patient's skin prior to surgery ■ skin wound and general skin cleansing
Warnings For external use only Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficult breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.
Do not use ■ if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product ■ in contact with meninges ■ in the genital area ■ as a preoperative skin preparation of the head or face
When using this product ■ keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums. ■ if solution should contact these areas, rinse out promptly and thoroughly with water ■ wounds which involve more than the superficial layers of the skin should not be routinely treated ■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
Surgical hand scrub: ■ wet hands and forearms under running water for 30 seconds. Clean fingernails using a nailstick or similar cleaner. ■ scrub for 1.5 minutes with about 8 mL of product with or without a wet brush paying close attention to the nails, cuticles, and skin between the fingers ■ rinse thoroughly under running water for 30 seconds ■ wash for an additional 1.5 minutes with 8 mL of product and rinse under running water for 30 seconds ■ dry thoroughly

Drug Facts (continued)**Healthcare personnel handwash:**

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Questions or comments?

call 1-800-587-3721
Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

Manufactured for: SCJ Professional USA, Inc.
Charlotte, NC 28217
1-866-783-0422 www.scjp.com

 **WARNING:** Cancer – www.P65Warnings.ca.gov.
Keep from freezing.

Manufactured for: SCJ Professional USA, Inc. Charlotte, NC 28217

1-866-783-0422 www.scjp.com

WARNING: Cancer -- www.P65Warnings.ca.gov. Keep from freezing.

BACTOSHIELD CHG

chlorhexidine gluconate 2% solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0116-0038
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	2 mg in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LAURAMINE OXIDE (UNII: 4F6FC4MI8W)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0KO0R)	
COCO DIETHANOLAMIDE (UNII: 92005F972D)	
HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0116-0038-03	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/16/2022	01/31/2027

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019422	11/16/2022	01/31/2027

Labeler - Xttrium Laboratories, Inc. (007470579)

Revised: 12/2025

Xttrium Laboratories, Inc.