

**NAPROXEN SODIUM CAPLETS- naproxen sodium tablet, coated**  
**Dr Reddy's Laboratories Inc**

-----  
**NAPROXEN SODIUM**  
**TABLETS USP, 220 mg**

**PAIN RELIEVER / FEVER REDUCER (NSAID)**

**Drug Facts**

***Active ingredient (in each tablet/caplet)***

Naproxen sodium USP, 220 mg  
(naproxen USP, 200 mg) (NSAID)<sup>1</sup>

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

***Warnings***

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAID's (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech

- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> <li>▪ take 1 tablet/caplet every 8 to 12 hours while symptoms last</li> <li>▪ for the first dose you may take 2 tablets/caplets within the first hour</li> <li>▪ do not exceed 2 tablets/caplets in any 8 to 12 hour period</li> <li>▪ do not exceed 3 tablets/caplets in a 24-hour period</li> </ul>
Children under 12 years	<ul style="list-style-type: none"> <li>▪ ask a doctor</li> </ul>

***Other information***

- **each tablet/caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

***Inactive ingredients***

FD &C Blue # 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

***Questions or comments?***

call toll-free weekdays (8 AM to 10 PM EST) at **1-888-375-3784**

Distributed by:

The Kroger Co.

Cincinnati, OH 45202

**Made in India**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL SECTION**

Naproxen Sodium Caplets, 220 mg

HealthCare Aisle
NDC 43598-058-29

## Naproxen Sodium

### Tablets USP, 220 mg (NSAID)

Pain reliever / fever reducer  
STRENGTH TO LAST 12 HOURS

**200 Caplets**  
*(Capsule-Shaped Tablets)*

**Compare to Aleve® Caplets active ingredient\*\***

**TAMPER EVIDENT:** Do not use, if foil seal under cap printed with "sealed for your protection" is broken or missing.  
**IMPORTANT:** Read the directions and warnings before use.

**Drug Facts**

**Active ingredient (in each caplet)** Purposes  
Naproxen sodium USP, 220 mg (naproxen USP, 200 mg) (NSAID) \*..Pain reliever/fever reducer \* nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - headache
  - menstrual cramps
  - toothache
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Peel Here

3 43598 05829 0

LOT 150084832

EXP

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®

Distributed by:  
Dr. Reddy's Laboratories, Inc.  
Princeton, NJ 08540

Rev:05/21

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better

**NAPROXEN SODIUM CAPLETS**

naproxen sodium tablet, coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43598-058
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Naproxen Sodium</b> (UNII: 9TN87S3A3C) (Naproxen - UNII:57Y76R9ATQ)	Naproxen Sodium	220 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>hypromelloses</b> (UNII: 3NXW29V3WO)	
<b>magnesium stearate</b> (UNII: 70097M6I30)	
<b>cellulose, microcrystalline</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>povidone</b> (UNII: FZ989GH94E)	
<b>talc</b> (UNII: 7SEV7J4R1U)	
<b>titanium dioxide</b> (UNII: 15FIX9V2JP)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	R;273
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-058-29	200 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075168	08/04/2021	

**Labeler** - Dr Reddy's Laboratories Inc (802315887)

Revised: 12/2022

Dr Reddy's Laboratories Inc