

**ANTACID DOUBLE STRENGTH- aluminum hydroxide, magnesium hydroxide, simethicone liquid**

**Rij Pharmaceutical Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 5 mL) (One teaspoonful)**

Aluminum Hydroxide (Equi. to Dried Gel, USP) 400 mg

Magnesium Hydroxide 400 mg

Simethicone 40 mg

**Purpose**

Antacid

Antigas

**Uses**

relieves

- heartburn
- acid indigestion
- sour stomach
- associated symptoms of gas and upset stomach
- overindulgence of food and drink

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if**

symptoms last for more than 2 weeks

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

**Directions**

- **shake well before each use**
- adults and children 12 years, take 2 - 4 teaspoonsful between meals, at bedtime or as directed by a physician
- do not take more than 12 teaspoonsful in 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

#### **Other information**

- TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.
- Store at room temperature 15° - 30°C (59° - 86°F)
- keep tightly closed
- protect from freezing
- each teaspoon (5 mL) contains: magnesium 165 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

#### **Inactive ingredients**

flavors, hypromellose, parabens, saccharin, sodium hypochlorite, sorbitol, xanthan gum and water

#### **Principal Display Panel**

NDC 53807-158-12

Compare to the active ingredient in  
Extra Strength Mylanta®†



EXTRA STRENGTH

# ANTACID

ANTACID • ANTI-GAS

*Magnesia & Alumina*  
Oral Suspension  
*with Simethicone*

- *Non-Constipating*
- *Pleasant Tasting*

**ORIGINAL FLAVOR**



12 FL. OZ. (355 mL)

## EXTRA STRENGTH ANTACID

### Drug Facts

#### Active Ingredients

(In each 5 ml) (One teaspoonful)	Purpose
Aluminum Hydroxide (Equi. to Dried Gel, USP) 400 mg .....	Antacid
Magnesium Hydroxide 400 mg .....	Antacid
Simethicone 40 mg .....	Antigas

**Uses** relieves ■ heartburn ■ acid indigestion ■ sour stomach  
■ associated symptoms of gas and upset stomach ■ overindulgence  
of food and drink

#### Warnings

Ask a doctor before use if you have ■ kidney disease  
■ a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**  
taking a prescription drug. Antacids may interact with certain  
prescription drugs.

**Stop use and ask a doctor if** ■ symptoms last more than 2  
weeks

**Keep out of reach of children.** In case of accidental overdose,  
get medical help or contact a Poison Control Center immediately.

#### Directions ■ shake well before each use

- adults and children over 12 years: take 2-4 teaspoonful between  
meals, at bedtime or as directed by a physician
- do not take more than 12 teaspoonfuls in a 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

#### Other information

- **TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON CAP  
IS BROKEN OR MISSING.**
- store at room temperature 15°-30°C (59°-86°F)
- keep tightly closed ■ protect from freezing
- each teaspoon (5 mL) contains: magnesium 165 mg
- each teaspoon (5 mL) contains: sodium 1 mg
- does not meet USP requirements for preservative effectiveness

**Inactive ingredients:** flavors, hypromellose, parabens,  
saccharin, sodium hypochlorite, sorbitol, xanthan gum and water.

†This product is not manufactured or distributed by Johnson & Johnson • Merck, owner of  
the registered trademark MYLANTA® Maximum Strength.

All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from  
retailer where purchased.

Rev.06/12

Manufactured By:  
RIJ PHARMACEUTICAL CORP.  
40 Commercial Avenue  
Middletown, NY 10941

Distributed By:  
Kinray Inc.,  
Whitestone, NY 11357



# ANTACID DOUBLE STRENGTH

aluminum hydroxide, magnesium hydroxide, simethicone liquid

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53807-158
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
BUTYLPARABEN (UNII: 3QPIU3FV8)	
SACCHARIN (UNII: FST467XS7D)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	
SORBITOL (UNII: 506T60A25R)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
LEMON OIL (UNII: I9GRO824LL)	

## Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53807-158-05	148 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	
2	NDC:53807-158-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part332	03/16/1999	

**Labeler** - Rij Pharmaceutical Corporation (144679156)

**Establishment**

Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-158)

Revised: 4/2018

Rij Pharmaceutical Corporation