GELATO TOPICAL ANESTHETIC- benzocaine gel Mycone Dental Supply Co., Inc DBA Keystone Industries and Deepak Products Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose:

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings:

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dos age. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Keep out of reach of children.

Other Information:

Store at 59°-86°F (15°-30°C). Protect from freezing.

Inactive Ingredients:

Flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow#5 (tartrazine), yellow #6, as a color additive.

Questions or comments?


1-800-333-3131 or

keystoneindustries.com



mouth symptoms do not improve f swelling, rash or fever develops, If pregnant or breast feeding, Do not exceed recommended in children under 2 years of age Keep out of reach of children. Poison Control Center right away. (tartrazine), yellow #6, as a color contact occurs, flush with water. blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, Apply only amount needed to Questions or comments? dosage. If more than used for children under 2 years of age: sodium saccharin. May contain edness persists or worsens; or pain is accidentally swallowed, directed by a physician. If sore Store at 59° - 86° F (15 - 30°C). the oral mucosa to prevent or get medical help or contact a flavoring, PEG 3350, PEG 400, see your physician promptly. When using this product in 7 days; irritation, pain, or Avoid contact with eyes. If ask a physician before use. **nactive** ingredients Other information keystoneindustries.com Protect from freezing. -800-333-3131 or for teething Directions relieve pain. do not use additive.

may cause methemoglobinemia, a serious condition that must be attention if you or a child in your Benzocaine 200 mg (in each g) occur even if you have used this pale, gray or blue colored skin reduces the amount of oxygen dizziness or lightheadedness warning: Use of this product Allergy Alert: Do not use on carried in the blood. This can product before. Stop use and for more than 7 days unless For oral mucosal use only, as treated promptly because it allergies to local anesthetics temporary relief of pain due to minor dental procedures. such as procaine, butacaine, directed by dentist. For the benzocaine or other "caine" patients with a history of fatigue or lack of energy Methemoglobinemia seek immediate medical Active ingredients shortness of breath · rapid heart rate **Drug Facts** Oral Anesthetic care develops: Warnings Purpose anesthetics. Do not use headache (cyanosis)



directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

Benzocaine 200 mg (in each g)

Active ingredients

Drug Facts

• for teething
• in children under 2 years of age
When using this product
Avoid contact with eyes. If
contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away. If pregnant or breast feeding

may cause methemoglobinemia a serious condition that must be

warning: Use of this product

Methemoglobinemia

Warnings

ask a physician before use.

Keep out of reach of children.

Directions

 Apply only amount needed to the oral mucosa to prevent or

attention if you or a child in your

pale, gray or blue colored skin

care develops:

occur even if you have used this

product before. Stop use and

seek immediate medical

reduces the amount of oxygen

treated promptly because it

carried in the blood. This can

relieve pain.
• children under 2 years of age: do not use

Other information

dizziness or lightheadedness

rapid heart rate
 shortness of breath

headache

cvanosis

Allergy Alert: Do not use on

patients with a history of

fatigue or lack of energy

allergies to local anesthetics

such as procaine, butacaine benzocaine or other "caine"

Store at 59° - 86° F (15 - 30° C). Protect from freezing. Inactive ingredients
flavoring, PEG 3350, PEG 400,
sodium saccharin. May contain
blue #1, green #3, green #5, red
#3, red #28, red #40, yellow #5,
(tartrazine), yellow #6, as a color

anesthetics.

Do not use

Do not use • for more than 7 days unless

Questions or comments?

additive.

keystoneindustries.com

-800-333-3131 or

GELATO TOPICAL ANESTHETIC

For oral mucosal use only, as

Oral Anesthetic

Purpose

directed by dentist. For the

temporary relief of pain due

to minor dental procedures.

benzocaine gel

 Product Information

 Product Type
 HUMAN OTC DRUG
 Item Code (Source)
 NDC:68400-356

Route of Administration DENTAL

Active Ingredient/Active Moiety

Ingredient Name
Basis of Strength

Ingredient NameBasis of StrengthStrengthBENZO CAINE (UNII: U3RS Y48 JW5) (BENZO CAINE - UNII: U3RS Y48 JW5)BENZO CAINE200 mg in 1 g

Inactive Ingredients

Ingredient Name

Strength

POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)

·	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C RED NO. 28 (UNII: 767IP0 Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics		
Color	yellow	Score
Shape		Size
Flavor	PINEAPPLE (Pina Colada)	Imprint Code
Contains		

ı	Packaging				
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
ı	1 NDC:68400-356-30	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	07/01/2013	

Labeler- Mycone Dental Supply Co., Inc DBA Keystone Industries and Deepak Products Inc (014769301)

Establishment			
Name	Address	ID/FEI	Business Operations
Keystone Industries		014769301	manufacture(68400-356), label(68400-356)

Revised: 11/2018 Mycone Dental Supply Co., Inc DBA Keystone Industries and Deepak Products Inc