

ACETAMINOPHEN- acetaminophen tablet
Cardinal Health 107, LLC

0220K

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/ Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use □

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **This Unit Dose package is not child resistant and is Intended for Institutional Use Only.**

Directions

- **do not take more than directed (see overdose warning)**
- **adults and children 12 years and over:**
 - take 2 tablets every 4 to 6 hours while symptoms last
 - do not take more than 10 tablets in 24 hours, unless directed by a doctor
 - do not use for more than 10 days unless directed by a doctor
- **children 6 years to under 12 years**
 - o take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor
- **children under 6 years:** ask a doctor

Other information

Store in a dry place at 15° - 30°C (59° - 86°F).

Povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?

1-800-231-4670

.

Keep this carton for complete product information

Tamper Evident: Do not use if sealed blister units are broken or damaged.

Distributed by: MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

(800) 616-2471

www.majorpharmaceuticals.com

Overbagged with 10 tablets per bag, NDC 55154-7899-0

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

Distributed By:

Cardinal Health

Dublin, OH 43017

L5509849-10524 / L5509849-20524

Principal Display Panel


NDC 55154-7899-0

ACETAMINOPHEN

Regula Strength Tablets 325mg

Pain Reliever/Fever Reducer

10 TABLETS



X119

NDC 55154-7899-0

ACETAMINOPHEN Regular Strength Tablets 325mg Pain Reliever/Fever Reducer

10 TABLETS

Drug Facts

Active ingredient (In each tablet)
Acetaminophen 325 mg

Purpose
Pain reliever/ Fever
reducer

Inactive ingredients Povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Uses Temporarily relieves minor aches and pains due to:

- headache ■ muscular aches ■ backache ■ minor pain of arthritis
- the common cold ■ toothache ■ premenstrual and menstrual cramps
- temporarily reduces fever

Directions

- do not take more than directed (see overdose warning) adults and children 12 years and over:
- take 2 tablets every 4 to 6 hours while symptoms last
- do not take more than 10 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- children 6 years to under 12 years
- take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor
- children under 6 years: ask a doctor

Other information

Store in a dry place at 15° - 30°C (59° - 86°F).

WARNINGS: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Tamper Evident: Do not use if sealed blister units are broken or damaged.

Questions or comments? 1-800-231-4670

Distributed By: MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268
(800) 616-2471
www.majorpharmaceuticals.com

Distributed by Cardinal Health
Dublin, OH 43017

L5509849-10524

Lot: Exp:

Distributed by Cardinal Health
Dublin, OH 43017
L5509849-20524

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children.

Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product
Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur ■ redness or swelling is present
- These could be signs of a serious condition.

ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55154-7899(NDC:0904-6773)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	

STARCH, CORN (UNII: O8232NY3SJ)

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	CPC;220
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-7899-0	1 in 1 BAG	10/23/2018	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/23/2018	

Labeler - Cardinal Health 107, LLC (118546603)

Revised: 9/2025

Cardinal Health 107, LLC