

**INFANT GAS RELIEF DYE FREE- simethicone suspension**  
**AmerisourceBergen (Good Neighbor Pharmacy) 46122**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredient (in each 0.3 mL)**

Simethicone 20 mg

**Purpose**

Antigas

**Uses**

relieves the discomfort of infant gas frequently caused by air swallowing or certain formulas or foods

**Warnings**

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddenings
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- shake well before using
- mL=milliliter
- measure only with enclosed syringe provided. Do not use any other dosing device
- keep dosing device with product
- replace cap tightly to maintain child resistance
- remove cap and insert syringe into hole at top of the bottle and turn upside down
- pull back syringe until filled to the prescribed level. If you pass the prescribed level, simply push syringe back until have reached the desired level. Slowly dispense the liquid into your child's mouth

(towards inner cheek)

- dosage can also be mixed with 1 oz, cool water, infant formula, or other suitable liquids
- clean syringe well after use
- all dosages may be repeated as needed, after meals and at bedtime, or as directed by a physician. Do not exceed 12 doses per day
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	0.3 mL
24	over 2 years	0.6 mL

### **Other information**

- store between 20-25°C (68-77°F)
- do not freeze

### **Inactive ingredients**

citric acid, carboxymethylcellulose sodium, flavors, maltitol, microcrystalline cellulose, purified water, sodium benzoate, sodium citrate, xanthan gum

### **Questions or comments?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to Mylicon® Gas Relief active ingredient\*

Infant Gas Relief

Simethicone 20mg - Antigas

Dye-Free

- Alcohol Free
- Saccharin Free
- Non-Staining

FL OZ (mL)

Use only with enclosed syringe

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND THE BOTTLE IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by McNeil Consumer Healthcrae, distributor of Mylicon® Gas Relief.

Distributed by AmerisourceBergen

1300 Morris Drive

Chesterbrook, PA 19087

Visit us at [www.mygnp.com](http://www.mygnp.com)

### **Package Label**

**Drug Facts**

<b>Active ingredient</b> (in each 0.3 mL) Simethicone 20 mg	<b>Purpose</b> Antigas
---	---------------------------

**Uses** relieves the discomfort of infant gas frequently caused by air swallowing or certain formulas or foods

**Warnings**  
Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- shake well before using
- mL = milliliter
- measure only with enclosed syringe provided. Do not use any other dosing device.
- keep dosing device with product
- replace cap tightly to maintain child resistance
- remove cap and insert syringe into hole at top of bottle and turn upside down
- pull back syringe until filled to the prescribed level. If you pass the prescribed level, simply push syringe back until you have reached the desired level. Slowly dispense the liquid into your child's mouth (towards inner cheek).
- dosage can also be mixed with 1oz. cool water, infant formula, or other suitable liquids
- clean syringe well after each use
- all dosages may be repeated as needed, after meals and at bedtime, or as directed by a physician. Do not exceed 12 doses per day.
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
under 24	under 2 years	0.3 mL
over 24	over 2 years	0.6 mL

**Drug Facts (continued)**

**Other information**

- store between 20-25°C (68-77°F)
- do not freeze

**Inactive ingredients** anhydrous citric acid, carboxymethylcellulose sodium, flavor, maltitol, microcrystalline cellulose, purified water, sodium benzoate, sodium citrate, xanthan gum

**Questions or comments?**  
Call 1-877-753-3936 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Mylicon® Gas Relief.

Compare to Mylicon® Gas Relief active ingredient\*

**GOOD NEIGHBOR PHARMACY®** NDC 46122-051-83

**Infant Gas Relief**  
Simethicone 20mg - Antigas

**Dye-Free**

- Alcohol Free
- Saccharin Free
- Non-Staining

1 fl oz (30 mL)



Use only with enclosed syringe

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR PRINTED SAFETY SEAL AROUND THE BOTTLE IS BROKEN OR MISSING.



Distributed By  
AmerisourceBergen  
1300 Morris Drive  
Chesterbrook, PA 19087  
Questions or Concerns?  
Visit us at [www.mygnp.com](http://www.mygnp.com)

ABC# 10095734



0 87701 40849 6

PLD-8272B F0004143

Lot No:  
Exp. Date:

**GOOD NEIGHBOR PHARMACY Infant Gas Relief**

<b>INFANT GAS RELIEF DYE FREE</b>			
simethicone suspension			
<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46 122-051
Route of Administration	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 0.3 mL
<b>Inactive Ingredients</b>			
	<b>Ingredient Name</b>		<b>Strength</b>
	ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
	CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)		

<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-051-03	1 in 1 BOX	06/30/2014	12/31/2022
1		30 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part332	06/30/2014	12/31/2022

**Labeler** - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)

Revised: 11/2019

AmerisourceBergen (Good Neighbor Pharmacy) 46122