

KANK-A MOUTH PAIN- benzocaine liquid
Blistex Inc.

Kank-A®
Mouth Pain Liquid

Drug Facts

Active ingredient

Benzocaine 20.0% (w/w)

Purpose

Oral anesthetic/analgesic

Uses

- for the temporary relief of pain due to canker sores, minor irritation of the mouth and gums caused by dentures or orthodontic appliances, or minor injury of the mouth or gums

Warnings

METHEMOGLOBINEMIA WARNING

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

When using this product

- do not use this product for more than 7 days unless directed by a dentist or doctor.

If sore mouth symptoms do not improve in 7 days; if irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly.

- do not exceed recommended dosage.

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older:
 - to assure formation of a long-lasting film coating, dry affected area and apply medication undiluted with applicator
 - allow a few seconds for coating to form
 - use up to 4 times daily, or as directed by a dentist or doctor
 - children under 12 years of age should be supervised in the use of this product
 - children under 2 years of age: do not use

Other information

- do not purchase if package has been opened
- cap tightly after use to avoid evaporation
- avoid contact with the eyes
- avoid contact with clothing and household/furniture surfaces to prevent possible staining
- this is a personal care item, and should be used by one individual only

Inactive ingredients

benzyl alcohol, cetylpyridinium chloride, compound benzoin tincture, dimethyl isosorbide, ethylcellulose, flavor, octylacrylamide/acrylates/butylaminoethyl methacrylate copolymer, oleth-10, polyethylene glycol, propylene glycol, ricinus communis (castor) seed oil, SD alcohol 38-B (29.6% v/v), sodium saccharin, sucralose, tannic acid

PRINCIPAL DISPLAY PANEL - 9.75 mL Bottle Package

MAXIMUM STRENGTH

KANK-A®

MOUTH PAIN LIQUID

ORAL ANESTHETIC

Maximum Pain Relief

Forms Protective Coating

For Sores Inside the Mouth

Canker Sores

Denture Abrasions

Brace Irritation

ADA
Accepted
American
Dental
Association ®

CONVENIENT
CONTROL-TIP
APPLICATOR

Net 0.33 fl. oz. (9.75 mL)

MAXIMUM STRENGTH

KANKA[®]

MOUTH PAIN LIQUID

ORAL ANESTHETIC

Maximum Pain Relief

Forms Protective Coating


For Sores Inside the Mouth

**Canker Sores
Denture Abrasions
Brace Irritation**

**ADA
Accepted**

American
Dental
Association [®]

Net 0.33 fl. oz. (9.75 mL)



**CONVENIENT
CONTROL-TIP
APPLICATOR**

MAXIMUM STRENGTH

ADA

Helps temporarily relieve

KANK-A



relieve temporary relieve
discomfort of mouth sores

Drug Facts

Active ingredient **Purpose**
Benzocaine 20.0% (w/w) Oral anesthetic/
analgesic

Uses

- for the temporary relief of pain due to canker sores, minor irritation of the mouth and gums caused by dentures or orthodontic appliances, or minor injury of the mouth or gums

Warnings

METHEMOGLOBINEMIA WARNING:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert:

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use:

- for teething
- in children under 2 years of age

When using this product

- do not use this product for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; if irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see ▶

Drug Facts (continued)

- your dentist or doctor promptly.
- do not exceed recommended dosage.

Keep out of reach of children.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older:
 - to assure formation of a long-lasting film coating, dry affected area and apply medication undiluted with applicator
 - allow a few seconds for coating to form
 - use up to 4 times daily, or as directed by a dentist or doctor
- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: do not use

Other information

- do not purchase if package has been opened
- cap tightly after use to avoid evaporation
- avoid contact with the eyes
- avoid contact with clothing and household/furniture surfaces to prevent possible staining
- this is a personal care item, and should be used by one individual only

Inactive ingredients

benzyl alcohol, cetylpyridinium chloride, compound benzoin tincture, dimethyl isosorbide, ethylcellulose, flavor, octylacrylamide/acrylates/butylaminoethyl methacrylate copolymer, oleth-10, polyethylene glycol, propylene glycol, ricinus communis (castor) seed oil, SD alcohol 38-B (29.6% v/v), sodium saccharin, sucralose, tannic acid

©2024 Blistex Inc., P.O. Box 5392
Oak Brook, IL 60522-5392

♻️ Card is 100% Recyclable. #C010363



KANK-A MOUTH PAIN

benzocaine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10157-9477
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

Benzocaine (UNII: U3RSY48JW5) (Benzocaine - UNII:U3RSY48JW5)		Benzocaine	200 mg	in 1 mL
Inactive Ingredients				
Ingredient Name			Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)				
CETYLPIRIDINIUM CHLORIDE (UNII: D9OM4SK49P)				
compound benzoin tincture (UNII: UJZ8IA4D1U)				
DIMETHYL ISOSORBIDE (UNII: SA6A6V432S)				
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)				
ACRYLATE/BUTYLAMINOETHYL METHACRYLATE/METHYL ACRYLATE/METHYL METHACRYLATE/OCTYLACRYLAMIDE COPOLYMER (40000 WAMW) (UNII: 8RZ43KFB5K)				
OLETH-10 (UNII: JD797EF70J)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
CASTOR OIL (UNII: D5340Y2I9G)				
ALCOHOL (UNII: 3K9958V90M)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
TANNIC ACID (UNII: 28F9E0DJY6)				
Product Characteristics				
Color	ORANGE	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10157-9477-1	9.75 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product	12/01/2011	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph drug	M022		12/01/2011	

Labeler - Blistex Inc. (005126354)

Establishment

Name	Address	ID/FEI	Business Operations
Blistex Inc.		005126354	MANUFACTURE(10157-9477)

