

KANK-A MOUTH PAIN- benzocaine liquid

Blis tex Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Kank-A[®] Mouth Pain Liquid

Drug Facts

Active ingredient

Benzocaine 20.0% (w/w)

Purpose

Oral anesthetic/analgesic

Uses

- for the temporary relief of pain due to canker sores, minor irritation of the mouth and gums caused by dentures or orthodontic appliances, or minor injury of the mouth or gums

Warnings

METHEMOGLOBINEMIA WARNING

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

When using this product

- do not use this product for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; if irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your doctor or dentist promptly.
- do not exceed recommended dosage.

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help

or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older:
 - to assure formation of a long-lasting film coating, dry affected area and apply medication undiluted with applicator
 - allow a few seconds for coating to form
 - use up to 4 times daily, or as directed by a dentist or doctor
 - children under 12 years of age should be supervised in the use of this product
 - children under 2 years of age: do not use

Other information

- do not purchase if package has been opened
- cap tightly after use to avoid evaporation
- avoid contact with the eyes
- avoid contact with clothing and household/ furniture surfaces to prevent possible staining
- this is a personal care item, and should be used by one individual only

Inactive ingredients

benzyl alcohol, cetylpyridinium chloride, compound benzoin tincture, dimethyl isosorbide, ethylcellulose, flavor, octylacrylamide/acrylates/butylaminoethyl methacrylate copolymer, oleth-10, PEG-6, propylene glycol, ricinus communis (castor) seed oil, SD alcohol 38B (29.6% v/v), sodium saccharin, sucralose, tannic acid

PRINCIPAL DISPLAY PANEL - 9.75 mL Bottle Package

MAXIMUM STRENGTH

Blis tex[®]

KANK+A[®]

MOUTH PAIN LIQUID
ORAL ANESTHETIC

Maximum Pain Relief

Forms Protective Coating

For Sores Inside the Mouth

Canker Sores

Denture Abrasions

Brace Irritation

ADA

Accepted

American

Dental

Association

®

**CONVENIENT
CONTROL-TIP**

APPLICATOR

Net 0.33 fl. oz. (9.75 mL)

MAXIMUM STRENGTH

Blistex

KANKA

MOUTH PAIN LIQUID


ORAL ANESTHETIC

Maximum Pain Relief

Forms Protective Coating

For Sores Inside the Mouth

Canker Sores
Denture Abrasions
Brace Irritation



Net 0.33 fl. oz. (9.75 mL)

CONVENIENT CONTROL-TIP APPLICATOR

MAXIMUM STRENGTH KANKA
MOUTH PAIN LIQUID

SEE INSIDE FOR SAVINGS ON **KANKA SoftBrush**

1.5625

LIFT PANEL TO VIEW COUPON

ADA Accepted logo: Helps temporarily relieve pain due to mouth sores

| | |
|--|--|
| <p>Drug Facts</p> <p>Active ingredient Purpose Benzocaine 20.0% (w/v) Oral anesthetic/analgesic</p> <p>Uses</p> <ul style="list-style-type: none"> for the temporary relief of pain due to canker sores, minor irritation of the mouth and gums caused by dentures or orthodontic appliances, or minor injury of the mouth or gums <p>Warnings</p> <p>METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> pale, gray, or blue colored skin (cyanosis) headache rapid heart rate shortness of breath dizziness or lightheadedness fatigue or lack of energy <p>Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.</p> | <p>Drug Facts (continued)</p> <p>Do not use:</p> <ul style="list-style-type: none"> for teething in children under 2 years of age <p>When using this product</p> <ul style="list-style-type: none"> do not use this product for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; if irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly. do not exceed recommended dosage. <p>Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> adults and children 2 years of age and older: <ul style="list-style-type: none"> to assure formation of a long-lasting film coating, dry affected area and apply medication undiluted with applicator allow a few seconds for coating to form use up to 4 times daily, or as directed by a dentist or doctor <p>©2018 Blistex Inc., P.O. Box 5392 Oak Brook, IL 60522-5392 Satisfaction GUARANTEED Card is 100% Recyclable. #C010152</p>  <p>0 41388 20391 1</p> |
|--|--|

MANUFACTURER'S COUPON EXPIRES 12/31/22

SAVE \$2.00 ON KANKA SoftBrush

0041388-024064



To the Retailer: Blistex® Inc. will reimburse you for the face value of this coupon plus 8¢ if submitted as part payment for KANKA® SoftBrush® in compliance with our redemption policy. Coupons available upon request. Void if copied, prohibited or regulated. Coupon may not be assigned or transferred by you. Cash value 1/100 cent. Send to Blistex 1324, NCH Marketing Services, P.O. Box 683001, Ft. Worth, TX 76168-0001. Good only on KANKA® SoftBrush®, any other use constitutes fraud. Limit one coupon per purchase.

| | |
|---|--|
| <p>Drug Facts (continued)</p> <ul style="list-style-type: none"> children under 12 years of age should be supervised in the use of this product children under 2 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> do not purchase if package has been opened cap tightly after use to avoid evaporation avoid contact with the eyes avoid contact with clothing and household/furniture surfaces to prevent possible staining | <p>Drug Facts (continued)</p> <ul style="list-style-type: none"> this is a personal care item, and should be used by one individual only <p>Inactive ingredients benzyl alcohol, cetylpyridinium chloride, compound benzoic tincture, dimethyl isosorbide, ethylcellulose, flavor, octylacrylamide/acrylates/butylaminoethyl methacrylate copolymer, oleth-10, PEG-6, propylene glycol, nicotiana glauca (castor) seed oil, SD alcohol 38-B (29.6% w/v), sodium saccharin, sucralose, tannic acid </p> |
|---|--|

MAXIMUM STRENGTH KANKA

Maximum Relief from Mouth Pain!

- Maximum strength formula relieves pain fast

For Tooth & Gum Pain, try:

KANKA SoftBrush

Tooth & Gum Pain Gel



Strong Relief With A Soft Touch

- Fast, Effective Relief... combines two analgesics for dual relief.
- Ultra Soft Brush Tip... gently gets between teeth and gums.
- Unique Applicator...

- Forms protective coating to shield against further irritation
- Convenient control tip for better application



Helps temporarily relieve pain due to mouth sores

easy to apply in hard-to-reach areas.

Over-the-counter remedies are no substitute for practicing good dental hygiene and regular visits to a dentist. If sore mouth symptoms do not improve in 7 days; if irritation, pain or redness persists or worsens; or if swelling, rash or fever develops, see your doctor or dentist promptly.

KANK-A MOUTH PAIN

benzocaine liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|-----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:10 157-9477 |
| Route of Administration | BUCCAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------------|
| Benzocaine (UNII: U3RSY48JW5) (Benzocaine - UNII:U3RSY48JW5) | Benzocaine | 200 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| CETYLPIRIDINIUM CHLORIDE (UNII: D9OM4SK49P) | |
| DIMETHYL ISOSORBIDE (UNII: SA6A6V432S) | |
| ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B) | |
| POLYOXYL-10 OLEYL ETHER (UNII: JD797EF70J) | |
| POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| CASTOR OIL (UNII: D5340Y2I9G) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| TANNIC ACID (UNII: 28F9E0DJY6) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|--|
| Color | ORANGE | Score | |
| Shape | | Size | |
| Flavor | | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-------------------|---|----------------------|--------------------|
| 1 | NDC:10 157-9477-1 | 9.75 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product | 12/01/2011 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part346 | 12/01/2011 | |

Labeler - Blistex Inc. (005126354)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------|---------|-----------|-------------------------|
| Blistex Inc. | | 005126354 | MANUFACTURE(10157-9477) |

Revised: 9/2019

Blistex Inc.