ANTI-DIARRHEAL- loperamide hci tablet Lil' Drug Store Products, Inc.

ANTI-DIARRHEAL loperamide hcl tablet

Drug Facts

Active Ingredient (in each tablet)

Loperamide HCl, USP 2 mg

Purpose

Anti-diarrheal

Use

Controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

Heart alert

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this producttiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

■ symptoms get worse

- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ find right dose on chart. If possible, use weight to dose; otherwise use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hour
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

Other Information

Other information

■ store between 68-77°F (20-25°C)

Inactive Ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions or Comments?

Call 1-800-351-2000 (8:30 AM - 5:00 PM CST)

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc, McNeil Consumer Healthcare Divison, distributor of Imodium® A-D

Distributed by: Lil' Drug Store Products, Inc., Cedar Rapids, IA 52404

2 mg Loperamide Tablet

Compare to the active ingredient in Imodium® A-D*


Diamode

Anti-Diarrheal Relief

Loperamide HCl
Tablets, USP 2 mg
[tablet image]
actual size

1 CAPLET IN EACH PACKET

READ DRUG FACTS AND INSTRUCTIONS ON PACKET



ANTI-DIARRHEAL

loperamide hci tablet

_			
Drod		Intorm	STIAN
FIUU	ULL	Inform	Ialivii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:29485-9286

Route of Administration ORAL

Active Ingredient/Active Moiety

ingredient Name		Basis of Strength	Strength

LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - LOPERAMIDE - HYDROCHLORIDE 2 mg

Inactive Ingredients

Strength

Product Characteristics			
Color	green	Score	no score
Shape	OVAL	Size	9mm
Flavor		Imprint Code	123
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29485- 9286-2	25 in 1 BOX	07/27/2016	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:29485- 9286-3	30 in 1 BOX	07/27/2016	
2		1 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA074091	07/27/2016	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Establishment				
Name	Address	ID/FEI	Business Operations	
Ohm Laboratories Inc.		184769029	manufacture(29485-9286)	